Erratum to: Practical application of acid dissociation in monitoring patients treated with adalimumab

Francisca Llinares-Tello · José Rosas-Gómez de Salazar · José Miguel Senabre-Gallego · Gregorio Santos-Soler · Carlos Santos-Ramírez · Esteban Salas-Heredia · Xavier Barber-Vallés · Juan Molina-García · AIRE-MB Group

Published online: 13 June 2014
© Springer-Verlag Berlin Heidelberg 2014

Erratum to: Rheumatol Int DOI 10.1007/s00296-014-3032-0

Unfortunately, one of the units in the below sentence was incorrectly published under the section “Patients and methods”. The correct sentence should read as:

Briefly, 20 μ L of serum was mixed with 100 μ L of 300 mM acetic acid to a final pH of 3.0.

The online version of the original article can be found under doi:[10.1007/s00296-014-3032-0](https://doi.org/10.1007/s00296-014-3032-0).

F. Llinares-Tello (✉) · J. Molina-García
Laboratory Department, Marina Baixa Hospital,
Av Jaime Botella Mayor nº7, 03570 Villajoyosa,
Alicante, Spain
e-mail: paquillinares@eresmas.com

J. Rosas-Gómez de Salazar · J. M. Senabre-Gallego ·
G. Santos-Soler · E. Salas-Heredia
Rheumatology Department, Marina Baixa Hospital,
Villajoyosa, Alicante, Spain

C. Santos-Ramírez
Rheumatology Department, Marina Salud Hospital,
Denia, Alicante, Spain

X. Barber-Vallés
CIO-Universidad Miguel Hernández, Elche, Alicante, Spain