Jocelyne Letarte · Ecaterina Simion · Mai Miner ·
Ken J. Kasha

Arabinogalactans and arabinogalactan-proteins induce embryogenesis in wheat (*Triticum aestivum* L.) microspore culture

Published online: 4 July 2006
© Springer-Verlag 2006

Plant Cell Reports (2006) 24: 691–698

In Material and methods, the subsections “Microspore induction culture” and “Plant differentiation and regeneration” contain errors. Please find the correct sentences below.

Microspore induction culture

The induction medium, MMS4, (Kasha et al. 2003a, b) was modified MS medium (Murashige and Skoog 1962) containing 2 mg l⁻¹ PAA, 0.5 mg l⁻¹ kinetin, 90 g l⁻¹ maltose, 355 mg l⁻¹ U2.5 amino acid mix (Comeau et al. 1992), and 975 mg l⁻¹ glutamine.

Plant differentiation and regeneration

Once ELS reached a size of about 2 mm, they were transferred onto differentiation medium: MMS5 (Kasha et al. 2003a, b) containing 0.5 mg l⁻¹ PAA, 0.2 mg l⁻¹ kinetin, 0.5 mg l⁻¹ GA₃, 30 g l⁻¹ maltose, 355 mg l⁻¹ Sigma U2.5 amino acid mix, 10 μM CuSO₄ at 22 °C during a 16 h light period.

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00299-005-0013-5>

J. Letarte · E. Simion · M. Miner · K. J. Kasha (✉)
Department of Plant Agriculture, University of Guelph,
Ontario, Canada, N1G 2W1
e-mail: kkasha@uoguelph.ca
Tel.: +519-824-4120
Fax: +519-763-8933