Influence of IBA and aphidicolin on DNA synthesis and adventitious root regeneration from Malus ‘Jork 9’ stem discs

Monika Sedira · Margareta Welander ·
Thomas Geier

Published online: 28 April 2007
© Springer-Verlag 2007

Erratum to: Plant Cell Rep DOI 10.1007/s00299-006-0249-8

In Materials and methods, the subsection “Rooting experiments with different concentrations of aphidicolin” contains an error. Please find the correct sentence below.

The basal rooting medium (BM) consisted of macronutrients in half strength and micronutrients according to Lepoivre (Quoirin et al. 1972): 200 mg/l NH_4NO_3 ,

900 mg/l KNO_3 , 600 mg/l $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$, 180 mg/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 135 mg/l KH_2PO_4 , 40 mg/l Fe–Na–EDTA, 8.6 mg/l $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 6.2 mg/l H_3BO_3 , 1 mg/l $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 0.0025 mg/l $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.008 mg/l KJ, 0.025 mg/l $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 0.0025 mg/l $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, supplemented with Walkey vitamins (Walkey 1972), 100 mg/l proline, 20 g/l sucrose, 5.3 g/l mannitol, 7 g/l agar (Difco Bacto).

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00299-006-0249-8>.

M. Sedira · M. Welander (✉)
Department of Crop Science,
Swedish University of Agricultural Sciences,
P.O. Box 44, 23053 Alnarp, Sweden
e-mail: margareta.welander@vv.slu.se

T. Geier
Fachgebiet Botanik,
Forschungsanstalt Geisenheim,
65366 Geisenheim, Germany