The rice pathogen-related protein 10 (JIOsPR10) is induced by abiotic and biotic stresses and exhibits ribonuclease activity

Sun Tae Kim · Seok Yu · Young Hyun Kang ·
Sang Gon Kim · Jae-Yean Kim · Sun-Hyung Kim ·
Kyu Young Kang

Published online: 3 September 2008
© Springer-Verlag 2008

Erratum to: Plant Cell Rep (2008) 27:593–603
DOI 10.1007/s00299-007-0485-6

Unfortunately, the presentation of Fig. 6 was wrong.
The correct Fig. 6 is given below.

The online version of the original article can be found under
doi:[10.1007/s00299-007-0485-6](https://doi.org/10.1007/s00299-007-0485-6).

S. T. Kim · J.-Y. Kim · K. Y. Kang (✉)
Environmental Biotechnology National Core Research Center,
Gyeongsang National University, Jinju 660-701, Korea
e-mail: kykang@gnu.ac.kr

S. Yu · J.-Y. Kim · K. Y. Kang
Plant Molecular Biology and Biotechnology Research Center,
Gyeongsang National University, Jinju 660-701, Korea

Y. H. Kang · S. G. Kim · J.-Y. Kim · K. Y. Kang
Division of Applied Life Science (BK21 Program),
Gyeongsang National University, Jinju 660-701, Korea

S.-H. Kim
Department of Environmental Horticulture,
University of Seoul, Seoul 130-743, Korea

Fig. 6 Immunolocalization of JIOsPR10 proteins during leaf and coleoptiles senescence. For immunohistochemical analysis, sections from senescent leaf and coleoptiles obtained with a microtome were incubated with purified JIOsPR10 antibody (dilution 1:100). Transverse (**a** and **b**) and longitudinal (**c** and **d**) sections of the senescent leaf and all transverse (**e**, **f**, **g**, and **h**) sections of the coleoptiles were

incubated with anti-JIOsPR10. Tissues were treated with preserum as a negative control (**a**, **e**, and **f**). The sections were incubated with alkaline phosphatase-conjugated anti-rabbit IgG antibody (dilution 1:300) for 1 h and visualized after the addition of NBT/BCIP solution for 20 min