

John F. Bruzzi
Alan C. Moss
Darren D. Brennan
Padraic MacMathuna
Helen M. Fenlon

Efficacy of IV Buscopan as a muscle relaxant in CT colonography

Published online: 4 February 2004
© Springer-Verlag 2004

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00330-003-2012-7>

J. F. Bruzzi · D. D. Brennan · H. M. Fenlon (✉)
Department of Radiology,
Mater Misericordiae Hospital,
Eccles Street, Dublin 7, Ireland
e-mail: hfenlon@eircom.net
Tel.: +353-1-8032274

A. C. Moss · P. MacMathuna
Department of Gastroenterology,
Mater Misericordiae Hospital,
Eccles Street, Dublin 7, Ireland

Eur Radiol (2003) 13:2264–2270

In the Results section (last paragraph, third sentence) and in Table 5, the numbers of polyps of sizes >9 mm and 6–9 mm were given incorrectly. The sentence should read “Conventional colonoscopy detected 6 polyps or masses larger than 9 mm, 7 polyps between 6 and 9 mm in diameter, and 34 polyps smaller than 6 mm (Figs. 2, 5; Table 2).”

The corrected table is given here. The change to the number of polyps does not affect the published sensitivity and specificity data, which were calculated on the correct figures.

Table 5 Number of polyps detected at conventional colonoscopy

Polyp size (mm)	No. of polyps Buscopan group	No. of polyps Non-Buscopan group	Total no. of polyps
>9	2	4	6
6–9	1	6	7
<6	15	19	34