A. S. Borthne
T. Reiseter
M. Abdelnoor
J. Rugtveit
G. Perminov
N.-E. Kløw

Bowel magnetic resonance imaging of pediatric patients with oral mannitol MRI compared to endoscopy and intestinal ultrasound

Published online: 23 May 2006
© Springer-Verlag 2006

Eur Radiol (2006) 16:207–214

The second sentence of the Abstract should read:

A median volume of 300 ml of mannitol in a 15% watery solution was ingested by 43 children prior to examination.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00330-005-2793-y>

A. S. Borthne (✉) · T. Reiseter
Department of Pediatric Radiology,
Ullevaal University Hospital,
Kirkeveien 166,
0407 Oslo, Norway
e-mail: aborth@broadpark.no
Tel.: +47-2-2119365
Fax: +47-2-3016588

M. Abdelnoor
Section of Epidemiology and Biostatistics,
Centre of Clinical Research, Ullevaal University Hospital,
Kirkeveien 166,
0407 Oslo, Norway

J. Rugtveit
Department of Pediatric Gastroenterology,
Ullevaal University Hospital,
Kirkeveien 166,
0407 Oslo, Norway

G. Perminov
Department of Pediatrics,
Akershus County University Hospital,
1473 Nordbyhagen, Norway

N.-E. Kløw
Department of Cardiovascular Radiology,
Ullevaal University Hospital,
Kirkeveien 166,
0407 Oslo, Norway