ERRATUM

Erratum to: Aiming for a shorter rheumatoid arthritis MRI protocol: can contrast-enhanced MRI replace T2 for the detection of bone marrow oedema?

Wouter Stomp¹ · Annemarie Krabben² · Désirée van der Heijde² · Tom W. J. Huizinga² · Johan L. Bloem¹ · Annette H. M. van der Helm-van Mil² · Monique Reijnders¹

Published online: 23 June 2016
© European Society of Radiology 2016

Erratum to: Eur Radiol
DOI 10.1007/s00330-014-3272-0

Due to a transcription error, an incorrect Fig. 3b was mistakenly included. Please find the correct Fig. 3b below:
The authors apologise for this error.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00330-014-3272-0>.

✉ Wouter Stomp
w.stomp@lumc.nl

¹ Department of Radiology, Leiden University Medical Center,
P.O. Box 9600, 2300 RC Leiden, The Netherlands

² Department of Rheumatology, Leiden University Medical Center,
P.O. Box 9600, 2300 RC Leiden, The Netherlands

Fig. 3 Correlation between total BME scores scored on T1Gd and T2 sequences. Scores for observer 1 in early arthritis (a) and advanced RA (b), and for observer 2 in early arthritis (c) and advanced RA (d). Scores

on T2 on the horizontal axis and scores on T1Gd on the vertical axis. Solid line represents the linear regression line; dashed line indicates the best possible (1:1) correlation