Aldona Bieniek

Archaeobotanical analysis of some early Neolithic settlements in the Kujawy region, central Poland, with potential plant gathering activities emphasised

Published online: 15 April 2004
© Springer-Verlag 2004

Veget Hist Archaeobot (2002) 11:33–40

Due to an unfortunate error Table 1 was printed incorrect.
The correct table is printed as follows:

The online version of the original article can be found at <http://dx.doi.org/10.1007/s003340200004>

A. Bieniek (✉)
W. Szafer Institute of Botany,
Polish Academy of Sciences,
Lubicz 46, 31-512 Kraków, Poland
e-mail: bieniek@ib-pan.krakow.pl

Table 1

Sites Cultures Taxa	Z1 L	G2 LP	G2 L	M4 LP	M4 L	O1 L	M4a L	K1 L	K1a L
Cultivated plants									
Hulled wheats - spikelet basis (x2) and glume basis	141	18	26		4448	177	449	81	47
<i>Triticum monoccum</i> - caryopsis	2				4		1		
<i>T. dicoccon</i> type - caryopsis	1			1	41	1	5		
<i>T. aestivum</i> s.l. - caryopsis					1?	5?	2?		
rachis internodes					1				
<i>Hordeum vulgare</i> ssp. <i>vulgare</i> - caryopsis		1							
cf. <i>Hordeum vulgare</i> - caryopsis									1
rachis internodes							2	1	2
<i>Hordeum</i> / <i>Triticum</i> - caryopsis	1				99		5		
rachis internodes									3
<i>Triticum</i> sp. – caryopsis					25	30	3	1	1
Cerealia indet. - caryopsis		5	5		12	17	33	10	
<i>Linum usitatissimum</i>					1				
<i>Pisum sativum</i> / <i>Vicia sativa</i>					1	1	1		
Wild plants									
<i>Avena</i> sp. – awn fragments		1							
<i>Bromus</i> sp.					12	6	14	1	
<i>Bromus hordaceus</i>				1	4				
<i>Bromus racemosus/arvensis</i>	4				8	4	8		
<i>Bromus sterilis/tectorum</i>						1	3		
<i>Campanula</i> sp.							6		
Caryophyllaceae indet.							1		
Chenopodiaceae indet.		1					1	6	
<i>Chenopodium album</i> type				50		12	39	5	15
<i>Corylus avellana</i>							1		
Cyperaceae indet.							3		
<i>Echinochloa crus-galli</i>					1		4		
<i>Fallopia convolvulus</i>		2		2	32	6	34	4	1
<i>Galium aparine</i>					1				
<i>Galium mollugo/sylvaticum</i>							3		
<i>Galium spurium</i>					9		3		
<i>Galium verum</i> type					2	1	12		
<i>Hierochloë</i> cf. <i>australis</i>							15		
<i>Juncus</i> sp.									1
Lamiaceae indet.	1						1	1	
<i>Melandrium album</i> type					2			1	
cf. <i>Mentha</i> sp.							1		
<i>Origanum vulgare</i>							1		
<i>Poa annua</i> type									1
<i>Poa nemoralis</i> type							2		
Poaceae indet. small		1			2	3	5		1
Poaceae indet. large		4			14		6	6	
<i>Polygonum aviculare</i>							2		
<i>Polygonum lapathifolium</i> s.l.				2					
<i>Polygonum minus</i>		2							
<i>Potentilla</i> sp.					1		2		
<i>Schoenoplectus tabernaemontani</i> / <i>triqueter</i>							2		
<i>Setaria pumila</i>							1		
<i>Setaria viridis/verticillata</i>			1			1	2		
<i>Stipa pennata</i> s.l. – awn fragments		1			122	179	761	519	665
<i>Stipa pennata</i> s.l. – caryopsis							4	6	7
Trifolieae indet.		1					2		
cf. <i>Vaccinium vitis-idaea</i>					1				
<i>Verbascum</i> sp.							2		
<i>Veronica</i> / <i>Plantago</i>									3
<i>Vicia tetrasperma/hirsuta</i>							1		
No. of features	1	2	1	2	6	6	9	13	3
No. of samples	1	2	1	2	8	9	16	20	3
total volume of soil (dm ³)	70	16	8	16	72	81	128	20	24