

Parkinson's disease mouse models in translational research

Paul M. A. Antony · Nico J. Diederich ·
Rudi Balling

Received: 24 February 2011 / Accepted: 14 April 2011 / Published online: 11 May 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract Animal models with high predictive power are a prerequisite for translational research. The closer the similarity of a model to Parkinson's disease (PD), the higher is the predictive value for clinical trials. An ideal PD model should present behavioral signs and pathology that resemble the human disease. The increasing understanding of PD stratification and etiology, however, complicates the choice of adequate animal models for preclinical studies. An ultimate mouse model, relevant to address all PD-related questions, is yet to be developed. However, many of the existing models are useful in answering specific questions. An appropriate model should be chosen after considering both the context of the research and the model properties. This review addresses the validity, strengths, and limitations of current PD mouse models for translational research.

Introduction

Age-related neurodegenerative diseases are an increasing burden to an aging population. Parkinson's disease (PD), the most common neurodegenerative movement disorder, frequently occurs in an idiopathic form without clearly defined cause. Familial forms of PD, arising from mono-genetic mutations, account for a minority of PD cases. The

clinical course of both idiopathic and familial forms of PD is believed to be influenced by environmental and genetic risk factors. Mouse models are powerful tools in PD translational research. However, an appropriate model should be chosen after considering both the model characteristics and the research context.

The Parkinson's disease puzzle

PD is a progressive neurodegenerative disorder clinically characterized by the cardinal symptoms of resting tremor, bradykinesia, cogwheel rigidity, and postural instability (Jankovic 2008). Responsiveness to L-3,4-dihydroxyphenylalanine (L-DOPA) and brain imaging distinguish PD from other disorders (D'Costa et al. 1995; Isaias and Antonini 2010). The pathological hallmarks of PD are loss of dopaminergic cells in the substantia nigra pars compacta and subsequent loss of dopamine innervation in the striatum (Forno 1996). Motor symptoms are the most obvious consequence of this nigrostriatal neurodegeneration. However, not only the basal ganglia but also other parts of the central nervous system as well as the autonomic nervous system are affected. A wide range of resulting non-motor symptoms can affect the patient's quality of life (Langston 2006; Lees 2009). It has been assumed that at least some subforms of PD start not with motor but non-motor symptoms such as hyposmia, rapid eye movement sleep behavior disorder (RBD), constipation, depression, reduced cardiac [¹²³I]metaiodobenzylguanidine (MIBG) uptake, and, possibly, reduced color discrimination (Chaudhuri et al. 2006; Fujishiro et al. 2008; Hawkes 2008; Katzenschlager and Lees 2004). There is also a broad consensus that neurodegenerative processes in PD start many years before the actual onset of clinical

P. M. A. Antony (✉) · N. J. Diederich · R. Balling
Luxembourg Centre for Systems Biomedicine (LCSB),
University of Luxembourg, 162, a Avenue de la Faïencerie,
1511 Luxembourg, Luxembourg
e-mail: paul.antony@uni.lu

N. J. Diederich
Department of Neuroscience, Centre Hospitalier de
Luxembourg, 4, rue Ernest Barblé, 1210 Luxembourg,
Luxembourg

manifestations (Savica et al. 2010; Tolosa et al. 2009). The not yet fully understood sequential plan underlying PD pathogenesis seems to correlate with clinical and pathological features (Hawkes et al. 2010). Lewy bodies, the histological hallmarks of PD, are α -synuclein-positive intracellular inclusions that are already present during the prodromal phase (Fearnley and Lees 1991). A yet asymptomatic, incidental Lewy body disease has been proposed (Markesberry et al. 2009). However, it is not yet fully understood if Lewy bodies have more protective or more destructive effects. Motor symptoms start only in the late phase of nigrostriatal degeneration, when 50–80% of nigral dopaminergic neurons are already lost (Bezard et al. 2001; Braak et al. 2003; Greford et al. 2006). At this stage, nonmotor features remain frequent and clinically significant because they largely contribute to impaired quality of life and shortened life expectancy. Neuroprotective treatments are not yet available but are one of the priorities in current PD research. However, reliable biomarkers for early diagnosis and disease state identification are not yet available (Gasser 2009a; Schapira et al. 2009).

Consequently, at present, the diagnosis of PD is based on the above-mentioned characteristic clinical motor features. Importantly, the currently recognized monogenic familial forms of PD do not demonstrate the same clinical features as those of sporadic late-onset forms of PD. Genetic forms can present more pronounced psychiatric features and lack nonmotor symptoms. Nevertheless, the phenotypical overlapping between familial and idiopathic PD is sufficient to dissect the commonly involved pathways. These include mitochondrial dysfunction, oxidative stress, protein misfolding, protein degradation, protein aggregation, and inflammation (Schapira and Tolosa 2010). Evidently, a better understanding of the molecular basis of PD will open new routes to diagnosis and therapy (Pienaar et al. 2008; Robinson 2010). On the other hand, it will influence the choice of adequate models for translational research.

Animal models for translational research in PD

Progress in understanding the etiology of PD has provided candidate targets for neuroprotective and neurorescue interventions. However, until now, no potential protective treatment has received regulatory approval (Rascol 2009). One reason for this is the lack of good PD animal models for preclinical translational research. Animal models should fulfill specific requirements for the testing of neuroprotective therapies for PD. First, the model should induce reproducible nigral lesions. Second, the loss of dopaminergic neurons should be stable, without spontaneous recovery. Third, the model should provide a time window for the application of potential neuroprotective

therapies (Erborg 2004). Ideally, a successful neuroprotective agent should prevent further behavioral changes, progressive biochemical deficits, and neurodegeneration. Thus, neuroprotection requires intervention when there are still cells left to be protected. Later interventions should be regarded as restorative strategies.

Current status and remaining challenges

Arvid Carlsson's research led to the development of L-DOPA-based treatments (Andersen 2009). Today, L-DOPA remains the gold standard treatment for PD. However, the drug has a wide range of adverse effects, most notably motor fluctuations and dyskinesias. Pharmacological strategies that avoid pulsatile dopaminergic stimulation have a substantial ameliorative effect on these syndromes. Despite these advances, disease progression remains unaffected (Lang and Obeso 2004). Finally, deep brain stimulation is the most recent symptomatic treatment that continuously and substantially reduces parkinsonian motor symptoms, whereas gait deterioration and nonmotor features continue to progress (Hamani et al. 2010).

Despite these therapeutic advances, PD patients continue to suffer from a severe reduction in quality of life. The development of interventions to stop or slow disease progression is therefore a major goal (Schapira and Tolosa 2010). The progress of this endeavor requires a better understanding of disease etiology and subtype stratification (Rajput et al. 2009). Importantly, early diagnosis enables the initiation of neuroprotective or other therapeutic interventions at less advanced stages of the disease.

Current mouse models for PD

PD mouse models are expected to closely match human pathology. The most important signs to replicate are motor symptoms, Lewy body formation, neuronal cell loss in the basal ganglia, age-related disease progression, and non-motor symptoms. None of the current models satisfies all of these criteria. However, many models fulfill a subset. Empirical choices of adequate models to answer specific questions in translational research need to consider, case by case, the model-specific advantages and limitations.

Pharmacological and neurotoxic PD models

Reserpine

One of the earliest developed PD models was the pharmacological reserpine model, which prevents the storage of

dopamine in presynaptic terminals, resulting in dopamine depletion in the striatum (Carlsson et al. 1957; Steg 1964). Reserpine rabbit models helped to elucidate the critical role of dopamine in the pathogenesis of PD. This finding led to the discovery of dopaminergic drugs such as L-DOPA. Despite this groundbreaking success, the reserpine model has significant limitations. Reserpine does not exclusively deplete dopamine and norepinephrine but all monoamines in a transient fashion. Another limitation of this model is that transient depletion does not mirror the progressive dopamine depletion in PD. In addition, the reserpine model does not reflect the histopathology of PD.

6-Hydroxydopamine (6-OHDA)

A few years after publication of the reserpine model, in the 1960s, nigrostriatal pathology was recognized as a primary factor for PD. In this context, the first neurotoxic models, aiming to reproduce nigrostriatal pathology, were established (Table 1). One neurotoxin found to induce PD was the dopamine analog 6-OHDA (Ungerstedt et al. 1974). 6-OHDA can be taken up into dopaminergic terminals via dopamine transporters. Inside the cell, it is metabolized, resulting in the production of hydrogen peroxide and free radicals, which induce neuronal death via mitochondrial dysfunction. 6-OHDA cannot cross the blood–brain barrier and therefore needs to be directly delivered to the brain. Ungerstedt et al. (1968) studied unilateral 6-OHDA lesions of the nigrostriatal pathway. By varying the position and extent of the lesion, different stages of human PD could be modeled. However, the 6-OHDA model has considerable limitations as well. It is a dopamine transporter-specific model and does not affect any other neurotransmitter.

Therefore, it does not replicate the full spectrum of clinical signs of human PD, i.e., the degeneration of the locus caeruleus with local depletion of norepinephrine is not replicated. In addition, the 6-OHDA-induced acute and local degeneration does not mirror the slow and extensive progression of neurodegeneration of the human disease. Finally, although lesioned animals show some of the cellular and behavioral deficits seen with the human disease, 6-OHDA-treated animals fail to develop Lewy bodies or α -synuclein-positive inclusions.

1-Methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)

In another model involving selective targeting of dopaminergic neurons in the brain, PD features are induced after administration of MPTP. The validity of the MPTP model is supported by the fact that MPTP exposure is a known, albeit rare, cause of drug-induced parkinsonism in humans (Davis et al. 1979; Langston et al. 1983). In mice, MPTP is typically administered systemically. MPTP is taken up primarily by astrocytes and converted to its metabolite MPP⁺. Then, it is taken up by dopaminergic neurons, where it exerts toxic effects. Interestingly, loss of dopaminergic neurons is observed not only in the basal ganglia but also in the enteric nervous system (Anderson et al. 2007). Varying regimens of MPTP delivery can produce varying degrees of nigrostriatal dysfunction. Acute dosing can produce lesions that result in mild to moderate cell loss, mimicking cell loss in early stages of human PD. Chronic dosing, consisting of long series of low-dose injections, results in more robust lesions representative of later stages of PD (Bezard et al. 1997). The MPTP mouse model has been invaluable in understanding the

Table 1 Pharmacological and toxicological models for Parkinson's disease

Molecule	Administration	Level of relevance	Target	I	MS	NMS	TH	BG	SP	Age	Ref
6-hydroxydopamine (6-OHDA)	Local	Derivate of dopamine	Causes oxidative stress after uptake by dopamine transporters	N	Y	NR	Y	Y	N	N	1
1-methyl-4-phenyl-1,2,3,4-tetrahydropyridine (MPTP)	Systemic	Known, albeit rare, cause of drug-induced parkinsonism in humans	Taken up by astrocytes primarily, converted to MPP ⁺ , which can be taken up by dopaminergic neurons, where it exerts toxic effects	Y	Y	Y	Y	Y	DD	Y	2
Rotenone	Systemic	Organic pesticide	Elicits mitochondrial dysfunction in a dopamine transporter independent fashion	Y	Y	NR	Y	Y	DD	NR	3
Paraquat, combined with maneb	Systemic	Paraquat is a pesticide, maneb is a fungicide	Causes redox cycling and thereby oxidative stress	Y	Y	NR	Y	Y	DD	NR	4

Age = age-related progression; BG = neurodegeneration in the basal ganglia; DD = dose-dependent; I = α -synuclein positive inclusions; MS = motor signs; N = no; NMS = nonmotor signs; NR = not reported; SP = slow progression; TH = reduced level of tyrosine hydroxylase; Y = yes

References (Ref): 1 = Ungerstedt et al. 1974; 2 = Anderson et al. 2007; Bezard et al. 1997; Davis et al. 1979; Irwin et al. 1992; Langston et al. 1983; 3 = Betarbet et al. 2000; p. 4 = Manning-Bog et al. 2002; Thiruchelvam et al. 2000

mechanisms underlying MPTP toxicity. However, despite robust replication of motor dysfunction and considerable preclinical successes in the screening for potentially therapeutic drugs, including coenzyme Q10, isradipine CR, and simvastatin (Beal et al. 1998; Ghosh et al. 2009; Meredith et al. 2008), the MPTP model has its limitations. For instance, mice may recover spontaneously after MPTP lesioning, thus hindering the assessment of therapeutic efficacy. The variability of MPTP sensitivity between distinct mouse strains and species is also problematic (Giovanni et al. 1994a, b; Sedelis et al. 2003). Studies on primates have demonstrated clear-cut parkinsonian features as well as protein aggregation after MPTP exposure. Consequently, this primate MPTP model has served as an adequate model for translational research toward the development and optimization of deep-brain stimulation therapies. In contrast, the MPTP model is less valid with other species, although as alluded above, the application of simvastatin in the MPTP mouse model produced interesting results in terms of protection against exotoxins (Ghosh et al. 2009).

Rotenone

The success of MPTP in replicating the neuropathology of human PD has led to the examination of other potential neurotoxin models for PD. Two promising models to emerge are the rotenone model and the combined paraquat/maneb model. Rotenone is an organic pesticide that inhibits complex I of the electron transport chain. It causes mitochondrial dysfunction and ultimately leads to cell loss in the nigrostriatal pathway (Betarbet et al. 2000). Rotenone can be delivered systemically via intranigral infusion or via gastric gavage to replicate human disease initiation in enteric neurons, as proposed by Braak (Braak et al. 2003). Rotenone exerts its effects in a dopamine transporter-independent fashion, thereby eliciting mitochondrial dysfunction in nondopaminergic systems as well. Nevertheless, its neurotoxic effects are predominantly dopamine neuron-selective. Data from cell culture experiments suggest that rotenone triggers intracellular dopamine release, which could in turn explain the toxic effect on dopaminergic neurons (Inden et al. 2011). However, the variability of sensitivity between individuals and species and the potential of this toxin to induce widespread systemic toxicity (Ravenstijn et al. 2008) complicates the use of the rotenone model for translational research.

Paraquat/maneb

Paraquat is a pesticide that is first converted to a cation before being reoxidized. This reaction produces superoxide radicals, which in turn cause further redox cycling and

oxidative stress. Paraquat is frequently administered in combination with maneb, a fungicide that has been linked to the development of parkinsonian symptoms in humans (Thiruchelvam et al. 2000). Exposure of rodents to a combination of paraquat and maneb results in significant nigral cell loss and nigral α -synuclein-positive inclusions. Rats prenatally exposed to maneb show increased sensitivity to paraquat toxicity in adulthood. These observations support the multihit hypothesis of PD, which suggests that early insults can sensitize the nigrostriatal system to subsequent hits in adulthood (Thiruchelvam et al. 2000).

Trichloroethylene

Trichloroethylene (TCE), a solvent widely used as a degreasing agent, is also considered an environmental risk factor for PD (Gash et al. 2008; Goldman 2010). Peritoneal administration of TCE causes dopaminergic neuronal cell death in mice (Guehl et al. 1999). In rats, oral administration of TCE induces key features of PD, i.e., loss of nigral dopaminergic neurons, reduced mitochondrial complex I activity, oxidative stress, microglial activation, and α -synuclein accumulation (Liu et al. 2010a, b). However, the relevance of this model for translational research remains to be evaluated.

Overall, pharmacological and neurotoxic models help to understand the consequences of striatal dopamine loss and to develop current dopaminergic therapies (Bove et al. 2005). A major limitation of toxin-induced models, however, is that they do not reproduce the complete spectrum of human pathology. Although they are excellent for evaluating the role of the nigrostriatal system in PD and may be adequate for testing restorative therapeutic compounds, most of these models are limited in their ability to replicate important aspects of human PD such as its progressive nature and Lewy body formation.

Genetic models for PD

In the last decade, considerable advances have been made in the identification of genes responsible for monogenic familial PD. Since 1998, 14 different genes have been identified to cause familial PD (Schapira 2006; Schapira and Tolosa 2010). In addition, recent genome-wide association studies (GWAS) led to the identification of yet another set of risk-associated genes. Besides its role in familial PD, α -synuclein is, worldwide, the most common risk factor for idiopathic PD (Pankratz et al. 2009; Satake et al. 2009). Strong association signals were also identified at the *MAPT* and *LRRK2* loci (Simon-Sánchez et al. 2009). Other important risk factors for PD are mutations in the glucocerebrosidase gene (Sidransky et al. 2009). A more

recently identified risk factor for PD, named PARK16, is associated with reduced risk for PD (Tan et al. 2010). Most recently, five additional PD risk factors have been reported: ACMSD, STK39, MCCC1/LAMP3, SYT11, and CCDC62/HIP1R (International Parkinson Disease Genomic Consortium 2011). As PD appears to be a complex genetic disease sensitive to environmental factors, the discovery of genetic factors in PD led to a new generation of PD models: the genetic PD mouse models.

α -Synuclein models

The very first identified mutations leading to familial PD were located in the gene encoding α -synuclein (Chartier-Harlin et al. 2004; Ibanez et al. 2004; Kruger et al. 1998; Polymeropoulos et al. 1997). In addition to single nucleotide polymorphisms (SNPs), rare triplications and duplications of the α -synuclein gene (*SNCA*) were reported in human PD (Ahn et al. 2008; Gasser 2009b; Singleton et al. 2003). However, α -synuclein, which typically accumulates in Lewy bodies (Spillantini et al. 1998), is not mutated in the large majority of PD patients. Nevertheless, numerous lines of mice overexpressing α -synuclein have been generated using distinct promoters and transgenes (Fernagut and Chesselet 2004). These models use transgenes with mutations causing familial PD, the full-length protein, or truncated forms of the protein. Most recapitulate some but not all aspects of PD, and effects relevant to the disease are generally correlated to the levels of α -synuclein expression (Magen and Chesselet 2010). Importantly, expression of transgenes is not limited to dopaminergic neurons but is defined by the nature of the genetic modification, i.e., promoter and genomic insertion site. However, a common advantage of these models, compared to pharmacological and toxicological models, is the slowly progressive accumulation of α -synuclein, a typical feature of idiopathic PD. The main strength of α -synuclein models is the ability to replicate α -synuclein-positive inclusions (Table 2), although the typical fibrillar halo structure of Lewy bodies, observed in human PD, is not observed in the murine α -synuclein-positive inclusions (Maries et al. 2003). A major limitation of α -synuclein models is their inability to model cell loss in the substantia nigra pars compacta. Mice overexpressing wild-type human α -synuclein exhibit reduced olfaction, autonomic dysfunction, α -synuclein accumulation, and early motor deficits in the absence of nigrostriatal neurodegeneration (Chesselet et al. 2008; Fleming et al. 2004; Fleming et al. 2008; Rockenstein et al. 2002).

α -Synuclein knockout mice are viable and show decreased striatal dopamine levels and reduced rearing. These findings may be explained by the hypothesis that α -synuclein plays a role in synaptic vesicle function. Triple

$\alpha\beta\gamma$ -synuclein knockout mice present age-dependent synaptic and neuronal dysfunction, demonstrating that synucleins contribute to the long-term operation of the nervous system (Burre et al. 2010; Greten-Harrison et al. 2010). Conversely, mice overexpressing α -synuclein develop intraneuronal inclusions and show decreased striatal dopamine levels, despite lacking nigral cell loss. However, mice overexpressing a double-mutated or a truncated form of α -synuclein under the tyrosine hydroxylase (TH) promoter present a selective loss of nigrostriatal dopaminergic neurons. The double-mutated α -synuclein variant (A30P/A53T) used in one of these studies has never been described in humans (Thiruchelvam et al. 2004). In contrast, expression of N-terminal α -synuclein fragments, comparable to those used in the second study (Wakamatsu et al. 2008a), has been detected in PD brain extracts (Li et al. 2005). In addition, the relevance of the A53T mutation in mouse models remains to be evaluated. Although the A53T mutation can cause PD in humans, threonine is the wild-type residue at this position in the murine sequence (Cabin et al. 2005; Larsen et al. 2009). Human A53T α -synuclein, however, was described to be more toxic on an α -synuclein-null background than on a murine wild-type background. This contrast between the protective effects of wild-type mouse α -synuclein and the deteriorative effects of human A53T α -synuclein on mice might be linked to the six other α -synuclein amino acid differences between these two species (Cabin et al. 2005).

Recently, data from tyrosine hydroxylase (TH) promoter-controlled α -synuclein-overexpressing mouse models led to the hypothesis that microglial activation is an early hallmark of PD (Su et al. 2008, 2009). Findings at the neurochemical and gene expression level suggest that α -synuclein also influences dopamine metabolism (Miller et al. 2007; Richfield et al. 2002; Yu et al. 2008). In addition, TH promoter-controlled overexpression of the double-mutated α -synuclein impairs the ubiquitin proteasome system in aged mice (Chen et al. 2006). The main limitation of this model is a spatially restricted pathology. This limitation can be circumvented, at least in part, by crossing α -synuclein transgenic mice with mice lacking murine α -synuclein. In this case, the absence of endogenous α -synuclein appears to drive the aggregation of transgenic α -synuclein throughout different brain regions (Prasad et al. 2011). In contrast, human Lewy body formation and α -synuclein expression in neurites are widely distributed in both the central and peripheral nervous systems (Braak et al. 2003; Halliday et al. 2005).

Platelet-derived growth factor- β (PDGF- β) promoter-controlled α -synuclein-overexpressing mouse models aim to mimic the broad distribution of α -synuclein pathology in humans (Masliah et al. 2000). Recent findings suggest that overexpression of A53T α -synuclein affects neurogenesis

Table 2 Alpha-synuclein models

Model(s)	Promoter	Background	I	MS	NMS	TH	BG	SP	Age	Ref
WT, A53T	PDGF- β	C57BL/6 × DBA2	Y	Y	NR	Y	Y	NR	Y	1
WT	KO	C57BL/6	NR	NR	NR	NR	NR	NR	NR	2
WT	PDGF- β	C57BL/6	NR	NR	NR	NR	NR	Y	NR	3
A53T	Mouse Thy-1	C57BL/6	LN	Y	NR	NR	N	NR	NR	4
WT, A30P, A53T	Mouse Thy-1	C57BL/6	Y	Y	IFC	NR	Y	NR	Y	5
<u>WT</u> , (A30P)	Mouse Thy-1	C57BL/6 × DBA2	Y	Y	Y	Y	Y	Y	N	6
Y39C	Mouse Thy-1	FVB/N	Y	Y	CD	N	N	Y	Y	7
A30P + A53T	Human Thy-1	C57BL/6 × DBA2	Y	Y	N	Y	Y	Y	N	8
(WT), (A30P), <u>A53T</u>	Mouse prion	C3H/HeJ × C57BL/6J backcrossed into C57BL/6J <i>PARKIN</i> KO	Y	Y	MD	N	Y	N	Y	9
WT, <u>A53T</u>	Mouse prion	C57BL/6 × C3H	Y	Y	RA	N	Y	Y	N	10
(WT), <u>A53T</u>	Mouse prion	FVB/N, FVB × 129, α -synuclein KO	N	Y	NR	NR	NR	Y	N	11
(WT), <u>A30P</u>	Mouse prion	C57BL/6J × DBA2 backcrossed into C57BL/6J	N	Y	ASP	N	NR	NR	Y	12
WT, <u>A30P</u> , A53T	Hamster prion	C57BL/6J × SJL	N	Y	NR	N	NR	Y	Y	13
WT, A30P, A53T	Rat THP	Swiss Webster × C57BL/DBA	N	NR	NR	N	NR	NR	NR	14
WT, <u>A30P ± A53T</u>	Rat THP	C57BL/6J	N	Y	NR	NR	NR	Y	Y	15
Truncated (1-120)	Rat THP	C57BL/6J × CBA/ca backcrossed into C57BL/6J, α -synuclein KO	Y	Y	NR	NR	N	Y	Y	16
Truncated (1-130)	Rat THP	C57BL/6J	N	N	REB	Y	Y	N	N	17
A30P + A53T	Chicken beta actin, BA	C57BL/6	NR	NR	MD	N	N	NR	Y	18
A30P + A53T	Mouse THP	C57BL/6	NR	NR	MD	N	N	NR	Y	19
A30P + A53T	Mouse prion	C57BL/6	NR	NR	NR	NR	N	NR	NR	20
WT, A30P, A53T	CaM-tTA (tet-off)	C57BL/6 (WT and A30P), C57BL/CH3 (WT and A53T)	N	Y	CD	NR	Y	Y	Y	21
A30P	KI in endogenous α -synuclein	C57BL/6	NR	Y	NR	RD	NR	Y	Y	22
Truncated (1-119)	Conditional ROSA26	C57BL/6J	N	NR	NR	RD	N	NR	N	23
A53T	Conditional ROSA26	C57BL/6J	N	NR	NR	NR	N	NR	N	23
WT, A30P, A53T	Endogenous α -synuclein (BAC)	FVB/N × 129S6/SvEvTac	N	Y	Y	NS	N	Y	N	24

Alpha-synuclein models express truncated or full-length α -synuclein with or without mutations. The expression of these constructs, in different mouse strains, is controlled by different promoters and causes distinct phenotypes. The models are grouped by project. *Lines* with a stronger phenotype are *underlined* and *lines without* phenotype are in *parentheses*

Age = age-related progression; ASP = affected synaptic plasticity; BA = beta-actin promoter; BG = neurodegeneration in the basal ganglia; CD = cognitive deficits; I = α -synuclein positive inclusions; IFC = impaired fear conditioning; LN = Lewy-like neurites; MD = mitochondrial deficits; MS = motor signs; N = no; NMS = nonmotor signs; NR = not reported; NS = not significant; RA = reduced anxiety; RD = reduced dopamine level; REB = reduced exploratory behavior; SP = slow progression; TH = reduced level of tyrosine hydroxylase; THP = tyrosine hydroxylase promoter; Y = yes

References (Ref): 1 = Hashimoto et al. 2003; Koob et al. 2010; Masliah et al. 2000; Rockenstein et al. 2002; Winner et al. 2004; Winner et al. 2008; Yacoubian et al. 2008; 2 = Sharon et al. 2003; 3 = Liu P et al. Liu et al. 2010a, b; 4 = van der Putten et al. 2000; 5 = Frasier et al. 2005; Freichel et al. 2007; Kahle et al. 2000; Neumann et al. 2002; Poon et al. 2005; Schell et al. 2009; 6 = Fernagut et al. 2007; Fleming et al. 2004; Fleming et al. 2006; Fleming et al. 2008; Koob et al. 2010; Rockenstein et al. 2002; Song et al. 2004; Wang et al. 2008; Watson et al. 2009; Wu et al. 2010; 7 = Zhou et al. 2008; 8 = Ikeda et al. 2009; Ono et al. 2009; 9 = Lee et al. 2002; Martin et al. 2006; Miller et al. 2007; Unger et al. 2006; von Coelln et al. 2006; 10 = Gao et al. 2008; George et al. 2008; Giasson et al. 2002; Graham and Sidhu 2010; Norris et al. 2007; Sotiriou et al. 2010; 11 = Cabin et al. 2005; Gispert et al. 2003; 12 = Gureviciene et al. 2007; 2009; Oksman et al. 2009; Yavich et al. 2006; Yavich et al. 2005; 13 = Gomez-Isla et al. 2003; Nieto et al. 2006; 14 = Manning-Bog et al. 2003; Matsuoka et al. 2001; Yu et al. 2008; 15 = Chen et al. 2006; Miller et al. 2007; Richfield et al. 2002; Su et al. 2009; Su et al. 2008; Thiruchelvam et al. 2004; 16 = Tofaris et al. 2006; 17 = Wakamatsu et al. 2008a; Wakamatsu et al. 2008b; 18 = Maskri et al. 2004; Stichel et al. 2007; 19 = Maskri et al. 2004; Stichel et al. 2007; 20 = Maskri et al. 2004; 21 = Lim et al. 2010; Marxreiter et al. 2009; Nuber et al. 2008; 22 = Plaas et al. 2008; 23 = Daher et al. 2009; 24 = Kuo et al. 2010

and proliferation of newly born neurons in the subventricular zone (Winner et al. 2004, 2008). Molecular analysis of mice with PDGF- β -controlled overexpression of α -synuclein suggested that interactions between α -synuclein, polyunsaturated fatty acids, and cholesterol accelerate α -synuclein aggregation (Bieschke et al. 2006; Sharon et al. 2003). Koob et al. (2010) used this model to show that treatment with the cholesterol synthesis inhibitor lovastatin alleviates these alterations. In PDGF- β/α -synuclein mice, gene expression is altered. However, neuronal death has not yet been reported in these mice (Yacoubian et al. 2008). Changes due to overexpression of α -synuclein are restricted mainly to the olfactory bulb and the hippocampus, and deficits occur relatively late. Despite changes at the gene expression level, dopamine metabolism, TH terminals, and neurogenesis, no overt pathology of the nigrostriatal system or clinical hallmarks have been observed in PDGF- β/α -synuclein mice.

In contrast to PDGF- β/α -synuclein models, prion protein promoter α -synuclein models aim to study the effects of extremely high overexpression of α -synuclein in neurons usually involved in PD, including neurons of the substantia nigra (Maskri et al. 2004). However, the translational relevance of these models, with typically severe motor deficits, is mitigated by high transgene expression in motor neurons of the spinal cord and brainstem, which confer behavioral deficits that are not characteristic for PD (Magen and Chesselet 2010). An additional drawback is the lack of cell loss in the substantia nigra and locus caeruleus.

Compared to the prion protein promoter, the Thy-1 promoter causes a more widespread expression of α -synuclein, which is not restricted to catecholaminergic neurons (Rockenstein et al. 2002), and does not cause any motor neuron loss when used in C57BL/6 mice. Nigral cell loss was observed in only a few cases, but most of the studies found typical PD phenotypes such as loss of dopamine in the striatum, motor deficits, and nonmotor deficits (Fleming et al. 2004; Ikeda et al. 2009; Ono et al. 2009; van der Putten et al. 2000). These mice further develop an age-dependent neurite terminal α -synuclein pathology, suggesting that α -synuclein is selectively phosphorylated in restricted brain regions (Schell et al. 2009). Some Thy1/ α -synuclein mouse lines present early motor and nonmotor deficits equivalent to those in PD prior to clinical diagnosis, suggesting that these models could be useful for the study of preclinical PD stages and hence for translational research (Fernagut et al. 2007; Fleming et al. 2004, 2006, 2008; Koob et al. 2010; Song et al. 2004). However, other studies using Thy1/ α -synuclein mouse lines demonstrated a much later onset, or even absence, of motor dysfunction (Freichel et al. 2007; Neumann et al. 2002; Zhou et al. 2008).

Models overexpressing α -synuclein under the calcium/calmodulin-dependent protein kinase II α (CaM) promoter were initially designed to study postnatal neurogenesis in the olfactory bulb and hippocampus. Conditional overexpression driven by the CaM promoter can be triggered by a tetracycline transactivator. The emergence of conditional transgenic models enables detailed investigation of the reversibility of α -synuclein pathology, thereby helping to focus on therapeutic strategies that aim to regulate α -synuclein expression at early stages of the disease (Lim et al. 2010; Marxreiter et al. 2009; Nuber et al. 2008). Recent models allow the control of gene expression in a region-dependent manner. One example is the Cre/loxP-based mouse model—expressing truncated N-terminal α -synuclein fragments—that shows reduced striatal dopamine levels but no neurodegeneration (Daher et al. 2009).

The most recently reported α -synuclein mouse model expresses A53T α -synuclein under a P1 artificial chromosome on an α -synuclein-null background. These mice have early motor deficits but no α -synuclein-positive inclusions, no loss of TH-positive cells in the substantia nigra, and no dopamine loss in the striatum. However, these mice have decreased colonic motility that may result from proteinase K-resistant α -synuclein aggregates found in neurons of the enteric nervous system. Olfactory function and cardiac innervation remain intact (Kuo et al. 2010). Consistently, this model could serve to investigate therapies for gastrointestinal dysfunction in early preclinical PD stages.

PARKIN, PTEN-induced putative kinase 1 (PINK1), and DJ1

In addition to the α -synuclein model, other models of familial PD have been reported. Recessive mutations in *PARKIN*, *PINK1*, and *DJ1* were mostly modeled using knockout strategies that result in the lack of the corresponding protein and, so far, to modest PD-associated deficits (Table 3).

The most frequent mutations causing early-onset recessive familial forms of PD occur in the *PARKIN* gene. *PARKIN* is an ubiquitin E3 ligase that ubiquitinates proteins to regulate a variety of cellular processes. Loss of E3 ligase activity is thought to play a pathogenic role in both inherited and sporadic PD (Dawson and Dawson 2010). More recently, *PARKIN* has been demonstrated to play a role in maintaining mitochondrial homeostasis through targeting damaged mitochondria for mitophagy (Tanaka 2010). *PARKIN* loss-of-function could thus lead to enhanced presence of dysfunctional mitochondria, a process that could contribute to PD pathogenesis. *PARKIN* knockout mice have been generated by deleting either exon 3 or 7. These lines show mild, progressive motor deficits and premotor signs typical of PD, i.e., reduced dopamine

Table 3 Summary of genetic PD perturbations and available genetic PD mouse models

Association with PD	Loci	Gene	Genotype(s) of mouse models	Promoter type	Background	I	MS	NMS	TH	BG	SP	Age	Ref
Common	PARK1 and 4	SNCA	Table 2 (T2)	T2	T2 Protamine-Cre mice (129/SvJae × (BALB/ c × C57BL/6 F1)) × C57BL/6	T2	T2	T2	T2	T2	T2	T2	2
Common	PARK2	PARKIN	Deletion of exon 7 causing a frame shift and premature termination	KO	NR N	Y	Y	Y	Y	Y	Y	N	1
Common	PARK2	PARKIN	PARKIN interrupted within exon 3	KO	129SV	N	Y	Y	N	NR	NR	NR	2
Common	PARK2	PARKIN	Deletion of exon 3	KO	C57BL/6J × 129/Sv	NR	N	Y	N	NR	NR	NR	3
Common	PARK2	PARKIN	Q311X	BAC	FvB	NR	Y	NR	Y	Y	Y	Y	4
Common	PARK2	PARKIN	Most of exon 3 was replaced with EGFP + stop	KO	C57BL/6J × 129/Sv	NR	Y	NR	N	NR	Y	Y	5
Common	PARK2	PARKIN	Exon 3 deletion (PaKO)	KO	129SvJ backcrossed into C57BL/6	NR	NR	MD	N	N	NR	Y	6
Common, combination	PARK2	PARKIN	Double mutant with PARKIN deletion (PaKO) and overexpression of A30P + A53T	KO + B A- α -synuclein	129SvJ × C57BL/6	NR	NR	MD	N	N	NR	Y	7
Common, combination	PARK2	PARKIN	Double mutant with PARKIN deletion (PaKO) α -synuclein driven by BA	KO + TH- α -synuclein	129SvJ × C57BL/6	NR	NR	MD	N	N	NR	Y	7
Common	PARK6	PINK1	Most of exon 2, and exons 3-5 were replaced by IRES-lacZ/MCIneo	KO	129SvEvBrd × C57BL/6J	NR	NR	HSN	N	NR	NR	NR	8
Common	PARK6	PINK1	Deletion of exons 4-7	PINK-1 null allele	C57BL/6 × 129/Sv	NR	NR	MD	N	N	NR	Y	9
Common	PARK6	PINK1	G309D	HR	129/SvEv	N	Y	MD	N	N	Y	Y	10
Common	PARK7	DJ1	Premature stop codon in exon 2	KO	C57BL/6	N	N	NR	N	NR	NR	N	11
Common	PARK7	DJ1	Deletion of a promoter subregion and the first 5 exons	Null allele	129 × C57BL/6J, E14Tg2A.4	N	Y	DD	N	NR	NR	Y	12
Common	PARK8	LRRK2	R1441G	BAC	FVB	NR	Y	DD	Y	NR	Y	Y	13
Common	PARK8	LRRK2	R1441C	KI	B6/129	N	MH	DD	N	NR	NR	NR	14
Common	PARK9	ATP13A2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	15
Common	Combination	PARKIN + PINK + DJ1	Triple KO	KO	C57BL/6 × 129/Sv	NR	NR	NR	N	N	NR	N	16

Table 3 continued

Association with PD	Loci	Gene	Genotype(s) of mouse models	Promoter type	Background	I	MS	NMS	TH	BG	SP	Age	Ref
Rare	PARK3	Unknown	SPR is a candidate gene for PARK3. The 2 first SPR exons are replaced with a lacZ-neomycin cassette	KO	C57BL/6J-Tyrc-Brd × 129S5/SvEvBrd	NR	Y	NR	Y	NR	NR	Y	17
Rare	PARK5	<i>UCHL1</i>	WT, I93 M. In addition, overexpression of α -synuclein was triggered via adenovirus.	PDGF- β	C57BL/6	NR	NR	NR	Y	Y	NR	NR	18
Rare	PARK11	<i>GIGYF2</i>	GIGYF2 \pm GIGYF2-/-	Gene trap	C57BL/6 × Tyr ^{c-Brd} /129Ola	Y	Y	NR	Y	NR	Y	Y	19
Rare	PARK13	<i>HTRA2</i>	Exons 2-6, and parts of exons 1 + 7 are replaced by the PGK-neo gene	KO	C57BL6/J	NR	Y	NR	Y	Y	N	N	20
Rare	PARK13	<i>HTRA2</i>	HTra2/Omi	NSE	FVB/N	NR	NR	NR	NR	NR	NR	NR	21
Rare	PARK14	<i>PLAG6</i>	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	22
Risk factor	PARK10	<i>AAOPD</i>	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	23
Risk factor	PARK12	Unknown	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	24
Risk factor	PARK15	<i>FBXO7</i>	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	25
Risk factor	PARK16	Unknown	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	26

The listed risk factors are restricted to PARK loci recently identified by means of genome-wide association studies (GWAS). Despite the lack of adequate models for some of these factors, the published mouse models cover the majority of the known genetic factors for PD

Age = age-related progression; BA = beta-actin promoter; BG = neurodegeneration in the basal ganglia; DD = dopaminergic dysfunction; HR = homologous recombination; HSN = hyperexcitable substantia nigra neurons; I = α -synuclein positive inclusions; MD = mitochondrial deficits; MH = only under multiple hits; MS = motor signs; N = no; NA = not available; NMS = nonmotor signs; NR = not reported; NSE = rat neuron-specific enolase promoter; SP = slow progression; TH = reduced level of tyrosine hydroxylase; THP = tyrosine hydroxylase promoter; Y = yes

References (Ref): 1 = O'Gorman et al. 1997; Von Coelln et al. 2004; 2 = Itier et al. 2003; Menendez et al. 2006; Periquet et al. 2005; Rodriguez-Navarro et al. 2007; 3 = Zhu et al. 2007; 4 = Lu et al. 2009; 5 = Goldberg et al. 2003; Martella et al. 2009; Palacino et al. 2009; Martella et al. 2007; Stichel et al. 2007; 6 = Stichel et al. 2004; 7 = Stichel et al. 2007; 8 = Bishop et al. 2010; Wood-Kaczmar et al. 2008; 9 = Gautier et al. 2008; Kitada et al. 2007; Martella et al. 2009; 10 = Gispert et al. 2009; 11 = Kim et al. 2005; 12 = Kim et al. 2005; 13 = Li et al. 2009; 14 = Tong et al. 2009; 15 = Ramirez et al. 2006; 16 = Kitada et al. 2009; 17 = Takazawa et al. 2008; 18 = Seisue et al. 2007; Yasuda et al. 2009; 19 = Giovannone et al. 2009; 20 = Bishop et al. 2010; Martins et al. 2004; 21 = Liu et al. 2007; 22 = Yoshino et al. 2010; 23 = Haugarvoll et al. 2009; 24 = Pankratz et al. 2003; 25 = Di Fonzo et al. 2009; 26 = Satake et al. 2009; Simon-Sanchez et al. 2009; Tan et al. 2010

release, synaptic anomalies, mitochondrial damage, and increased oxidative stress. However, none of these models shows a progressive loss of nigrostriatal dopaminergic neurons (Goldberg et al. 2003; Itier et al. 2003; Martella et al. 2009; Palacino et al. 2004; Periquet et al. 2005; Rodriguez-Navarro et al. 2007; Stichel et al. 2007; Von Coelln et al. 2004; Zhu et al. 2007). Dopaminergic cell loss can be induced by exposure to the inflammatory stressor lipopolysaccharide (LPS) (Frank-Cannon et al. 2008), illustrating that *PARKIN* mutations may make dopaminergic neurons more vulnerable to environmental insults. Lu et al. (2009) investigated whether the *PARKIN* mutation Q311X exerts dominant toxicity on nigrostriatal dopaminergic neurons. The selective expression of this mutation in dopaminergic neurons does indeed cause progressive nigrostriatal dopaminergic cell loss, α -synuclein pathology, and motor deficits (Lu et al. 2009). This implies that these mice provide a model for dopaminergic neuron death caused by *PARKIN* mutations.

Similar to *PARKIN* knockout mice, *PINK1* and *DJ1* mice have, so far, failed to reproduce canonical PD deficits. Similar observations were made in a triple transgenic model lacking *PARKIN*, *PINK1*, and *DJ1* (Kitada et al. 2009). Without additional insults, *PINK1* and *DJ1* deficiency does not cause spontaneous loss of nigrostriatal dopaminergic neurons (Aron et al. 2010; Morais et al. 2009). However, most of these mice show reduced activity, increased oxidative stress, and increased vulnerability to mitochondrial toxins (Chen et al. 2005; Gautier et al. 2008; Gispert et al. 2009; Kim et al. 2005). Overall, recessive PD models appear to have potential for the modeling of environmental factors in familial forms of PD.

Leucine-rich repeat kinase 2 (LRRK2)

Dominant *LRRK2* mutations have recently been identified as causes of familial PD. Heterozygote carriers can develop familial PD with characteristics similar to the sporadic disease forms (Cookson et al. 2008). However, penetrance is strongly age-dependent. At the age of 69 years, only 51% of carriers show signs of PD (Healy et al. 2008). Several mutations have been identified in human PD patients, the most frequent one being the G2019S mutation. Lin et al. (2009) generated *LRRK2* knockout mice and transgenic mice with inducible overexpression of the human wild-type, the G2019S mutant, or the kinase domain-deleted *LRRK2*. Neither deletion nor overexpression of these *LRRK2* constructs caused gross neurodegeneration. However, all three *LRRK2* constructs exacerbated the progression of human A53T α -synuclein-induced neuropathology. Overexpression of *LRRK2* or its mutants impaired the structure and function of the Golgi complex

and microtubule-based transport, resulting in a pathogenic somatic accumulation of α -synuclein (Lin et al. 2009).

Later, Li et al. (2010) characterized two bacterial artificial chromosome (BAC) transgenic mouse strains overexpressing either wild-type or G2019S-mutated *LRRK2*. Transgenic mice overexpressing wild-type *LRRK2* presented an elevated striatal dopamine release, were hyperactive, and showed enhanced motor performance. In contrast, mice expressing the G2019S mutant showed an age-dependent decrease in striatal dopamine content (Li et al. 2010). Melrose et al. (2010) also reported impaired dopaminergic neurotransmission in BAC *LRRK2* mice. In G2019S mice, these changes were linked to a modified localization and phosphorylation of the microtubule-binding protein tau (Melrose et al. 2010). A more recent study that also used mice that overexpress G2019S *LRRK2* by using a BAC construct, analyzed adult neurogenesis (Winner et al. 2011). The proliferation and survival of newly generated cells was significantly decreased by the overexpression of G2019S *LRRK2*, and newly generated neurons exhibited reduced dendritic arborization and fewer spines (Winner et al. 2011).

Mice harboring the R1441G mutation develop a strong behavioral phenotype and an altered dopamine release (Li et al. 2009). In contrast, mice harboring the R1441C mutation do not present prominent motor defects but only dopaminergic deficits (Tong et al. 2009). *LRRK2* seems to be involved in both familial and idiopathic PD. Therefore, *LRRK2* is a very attractive target for therapeutic intervention (Paisan-Ruiz et al. 2004; Satake et al. 2009; Simon-Sanchez et al. 2009; Zimprich et al. 2004). The value of *LRRK2* mouse models for translational research in this direction remains to be evaluated, but primary results indicate that these models might be useful for analyzing impaired cellular dynamics that cause dopaminergic dysfunction.

Models for other network perturbations in PD

Genetic models of PD are widely used. However, only a few of these models reproduce the cardinal features of the disease. Most strikingly, most genetically altered mice do not show extensive or progressive loss of nigrostriatal dopaminergic neurons (Chesselet et al. 2008). It is reasonable to assume that distinct causal factors leading to neurodegeneration in PD perturb converging molecular or cellular pathways. In fact, the concept of PD as network perturbation might be useful to guide translational research (Del Sol et al. 2010). Consequently, models for network perturbations are relevant to translational research in PD. These include the above-mentioned genetic and toxic models as well as models for genetic risk factors and

models for disease pathway perturbations such as mitochondrial dysfunction (Wellstead and Cloutier 2011).

Aphakia and weaver mice

Besides blindness, the so-called aphakia mice—lacking the transcription factor Pitx3—present complete loss of nigrostriatal dopaminergic neurons during early postnatal development. This model does not clearly reproduce progressive loss of dopaminergic neurons in PD, yet it is useful to validate behavioral mouse tests for dopaminergic deficits (Hwang et al. 2005). Pitx3 polymorphisms have been associated with an increased risk for PD (Haubnerberger et al. 2011) and may therefore be involved in human PD. In addition, Pitx3 may directly regulate expression of the vesicular monoamine transporter (*VMAT*) and the dopamine transporter (*DAT*), two genes associated with increased risk for PD (Hwang et al. 2009; Ritz et al. 2009; Takahashi et al. 1997).

In turn, weaver mutants carry a mutation in the G-protein-coupled inwardly rectifying channel (*GIRK2*) (Guatteo et al. 2004). An important characteristic of these mice for research in PD is a gradual nigrostriatal dopaminergic neurodegeneration (Cavalcanti-Kwiatkoski et al. 2010).

VMAT

Mice with reduced expression of VMAT, member 2 (*VMAT2*) have reduced monoamine storage capacity and present both a progressive loss of striatal dopamine and L-DOPA-responsive motor deficits. In addition, *VMAT2*-deficient mice present progressive nonmotor signs such as progressive deficits in olfactory discrimination, delayed gastric emptying, altered sleep latency, and age-dependent depressive behavior. This model might be valuable for translational research toward restoration of monoamine dysfunction at early nonmotor stages of PD (Taylor et al. 2009).

MitoPark mice

The so-called MitoPark mouse—lacking the mitochondrial transcription factor A (*TFAM*)—is a model of mitochondrial dysfunction in dopaminergic midbrain neurons. These mice exhibit an adult-onset progressive loss of nigrostriatal dopaminergic neurons, accompanied by the formation of intraneuronal inclusions (Ekstrand and Galter 2009; Ekstrand et al. 2007). As in human PD, impaired nigrostriatal function precedes the onset of motor deficits, and L-DOPA administration improves motor dysfunction (Galter et al. 2010; Good et al. 2011). MitoPark mice may provide a useful model for investigating neuroprotective strategies (Harvey et al. 2008). However, is it not yet fully

understood whether mitochondrial dysfunction is the cause of PD or whether it is a consequence of other perturbations within a cascade of pathogenic events.

Aging

Aging is the primary risk factor for the development of PD. In fact, there are many parallels between normal aging and early stages of PD. In normal aging, the striatal dopamine levels are reduced and there is loss of dopaminergic nigral neurons. These losses are associated with age-related increases in α -synuclein (Irwin et al. 1992; McNeill and Koek 1990). Collectively, these findings suggest that aged animals represent a useful model for the evaluation of therapies aimed at the earliest stages of PD and that it might be valuable to rethink aged animals' validity for translational research in PD.

Future challenges and needs in PD animal models and translational research

Today, we understand that the prodrome of PD can be initiated by genetic and environmental factors. However, the total number of involved genes is not yet known. In addition, different genetic backgrounds, for instance, between different consomic mouse strains, can influence the phenotypic outcome in animal models (Nishi et al. 2010). One challenge is, therefore, to better understand the complex relationships between genetic factors and PD onset and manifestation. One possibility is to identify regions of the genome that play a role in complex PD-related traits. In the case of PD, quantitative trait loci (QTLs) could be used to map the genes underlying parkinsonian phenotypes or susceptibility to PD. The Complex Trait Consortium has provided a reference panel of recombinant inbred strains designed for QTL analysis (Churchill et al. 2004; Vogel 2003). These congenic mouse strains, generated by mating distinct inbred strains and backcrossing the descendants, harbor nearly identical genomes. The resulting set of populations allows the analysis of the effects of different genetic backgrounds. The already existing genetic PD mouse models, in contrast, are designed to analyze the role of isolated genetic loci on a fixed background. Recently, GWAS aimed to better understand the genetic basis underlying PD. Both QTL analysis and whole-genome sequencing are powerful complementary tools for proceeding with these studies.

Future therapeutic interventions will likely aim at interrupting the progression of molecular and pathological prodromes. However, the causal cascade of pathogenic events is not yet fully understood. To improve the understanding of early pathogenesis, it will be necessary to

analyze the temporal and causal coupling between events at the molecular, cellular, and organism scales. Furthermore, the history of translational research in well-studied diseases (e.g., Alzheimer disease) has demonstrated that findings from mouse models may not always be extrapolated to humans (Van Dam and De Deyn 2011). The validity of animal models of human diseases will therefore need to be evaluated. Of great importance, biomarkers found in animal models might help improve clinical risk evaluation and early diagnosis. Furthermore, biomarkers will allow us to empirically choose the most adequate animal models for specific questions in translational research (Srivastava et al. 2010). Novel techniques like live cell imaging and nanosurgery in the living mouse brain (Mascaro et al. 2010), progress in the development of epigenetic analysis tools, and reduced costs for omics analysis will help in the investigation of the complexity underlying PD. The complex relationships between multiple scales and evaluation of model relevance for clinical research will require new tools for the analysis of complex biological systems, i.e., systems biology, computational biology, and interdisciplinary cooperation. These tools should be suitable for developing translational research into a predictive, preventive, personalized, and participatory medicine (Auffray et al. 2010). An important step on the way to achieve personalized medicine will be to rank the potential of therapeutic strategies per disease state. In addition to mouse models, other animal models, including other rodents, primates, *Drosophila melanogaster*, *Caenorhabditis elegans*, and *Saccharomyces cerevisiae* (Hirth 2010; Nass et al. 2008; Witt and Flower 2006) as well as cellular models, should be considered. The adequate choice of models is critical for the outcome of translational research.

Conclusion

It is broadly accepted that current PD mouse models do not adequately present typical sets of symptoms characteristic of human PD. The striking differences between brain size and complexity between rodents and humans limit the direct translation of the experimental results to the clinic. Model and species properties determine the characteristics of the syndrome. Full reproduction of neuropathological and clinical PD manifestations is not expected in the mouse. However, a useful model should at least be of mechanistic relevance, thereby allowing potential therapeutic testing. The awareness that many mouse models show premotor symptoms of PD raises the possibility of development of novel treatments. However, none of the available models accurately recapitulates the complex and progressive pathology that characterizes human PD. These important shortcomings must be taken into account when

translating information from the laboratory to the clinic. However, it is too simple to argue that the lack of reliable models for preclinical testing undermines investigations in clinical trials. An appropriate model should be chosen after considering the question to be answered. The perfect model that fits all requirements does not exist. However, some of the current models allow us to analyze precise questions within the PD disease spectrum. At present, preclinical models and models to evaluate enteric signs, nigrostriatal neurodegeneration, the multiple hits hypothesis, and acute motor dysfunction, among others, are available. In the past, translational research has advanced using appropriate models to aid the development of deep-brain stimulations and L-DOPA treatments. The current trend in translational PD research is neuroprotection or reversion of neurodegeneration. This endeavor requires models that reproduce early pathogenic events such as those observed in transgenic models. Better clinical PD stratification and identification of triggering environmental factors will further increase the demand for targeted models. The need for distinct model platforms and for the development of targeted therapeutic interventions will grow with our knowledge on PD etiology and stratification.

Acknowledgments We thank Dr. Manuel Buttini for insightful discussions and for helpful comments on the manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Ahn TB, Kim SY, Kim JY, Park SS, Lee DS, Min HJ, Kim YK, Kim SE, Kim JM, Kim HJ et al (2008) alpha-Synuclein gene duplication is present in sporadic Parkinson disease. *Neurology* 70:43–49
- Andersen JK (2009) Arvid Carlsson: an early pioneer in translational medicine. *Sci Transl Med* 1:2ps3
- Anderson G, Noorian AR, Taylor G, Anitha M, Bernhard D, Srinivasan S, Greene JG (2007) Loss of enteric dopaminergic neurons and associated changes in colon motility in an MPTP mouse model of Parkinson's disease. *Exp Neurol* 207:4–12
- Aron L, Klein P, Pham TT, Kramer ER, Wurst W, Klein R (2010) Pro-survival role for Parkinson's associated gene DJ-1 revealed in trophically impaired dopaminergic neurons. *PLoS Biol* 8:e1000349
- Auffray C, Charron D, Hood L (2010) Predictive, preventive, personalized and participatory medicine: back to the future. *Genome Med* 2:57
- Beal MF, Matthews RT, Tielemans A, Shults CW (1998) Coenzyme Q10 attenuates the 1-methyl-4-phenyl-1, 2, 3, tetrahydropyridine (MPTP) induced loss of striatal dopamine and dopaminergic axons in aged mice. *Brain Res* 783:109–114
- Betarbet R, Sherer TB, MacKenzie G, Garcia-Osuna M, Panov AV, Greenamyre JT (2000) Chronic systemic pesticide exposure

- reproduces features of Parkinson's disease. *Nat Neurosci* 3: 1301–1306
- Betz E, Dovero S, Bioulac B, Gross CE (1997) Kinetics of nigral degeneration in a chronic model of MPTP-treated mice. *Neurosci Lett* 234:47–50
- Betz E, Dovero S, Prunier C, Ravenscroft P, Chalon S, Guilloteau D, Crossman AR, Bioulac B, Brotchie JM, Gross CE (2001) Relationship between the appearance of symptoms and the level of nigrostriatal degeneration in a progressive 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine-lesioned macaque model of Parkinson's disease. *J Neurosci* 21:6853–6861
- Bieschke J, Zhang Q, Bosco DA, Lerner RA, Powers ET, Wentworth P Jr, Kelly JW (2006) Small molecule oxidation products trigger disease-associated protein misfolding. *Acc Chem Res* 39:611–619
- Bishop MW, Chakraborty S, Matthews GA, Dougalis A, Wood NW, Festenstein R, Ungless MA (2010) Hyperexcitable substantia nigra dopamine neurons in PINK1- and HtrA2/Omi-deficient mice. *J Neurophysiol* 104:3009–3020
- Bove J, Prou D, Perier C, Przedborski S (2005) Toxin-induced models of Parkinson's disease. *NeuroRx* 2:484–494
- Braak H, Del Tredici K, Rub U, de Vos RA, Jansen Steur EN, Braak E (2003) Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 24:197–211
- Burre J, Sharma M, Tsetsenis T, Buchman V, Etherton MR, Sudhof TC (2010) Alpha-synuclein promotes SNARE-complex assembly in vivo and in vitro. *Science* 329:1663–1667
- Cabin DE, Gispert-Sanchez S, Murphy D, Auburger G, Myers RR, Nussbaum RL (2005) Exacerbated synucleinopathy in mice expressing A53T SNCA on a Snca null background. *Neurobiol Aging* 26:25–35
- Carlsson A, Lindqvist M, Magnusson T (1957) 3, 4-Dihydroxyphenylalanine and 5-hydroxytryptophan as reserpine antagonists. *Nature* 180:1200
- Cavalcanti-Kwiatkoski R, Raisman-Vozari R, Ginestet L, Del Bel E (2010) Altered expression of neuronal nitric oxide synthase in weaver mutant mice. *Brain Res* 1326:40–50
- Chartier-Harlin MC, Kachergus J, Roumier C, Mouroux V, Douay X, Lincoln S, Levecque C, Larvor L, Andrieux J, Hulihan M et al (2004) Alpha-synuclein locus duplication as a cause of familial Parkinson's disease. *Lancet* 364:1167–1169
- Chaudhuri KR, Healy DG, Schapira AH (2006) Non-motor symptoms of Parkinson's disease: diagnosis and management. *Lancet Neurol* 5:235–245
- Chen L, Cagniard B, Mathews T, Jones S, Koh HC, Ding Y, Carvey PM, Ling Z, Kang UJ, Zhuang X (2005) Age-dependent motor deficits and dopaminergic dysfunction in DJ-1 null mice. *J Biol Chem* 280:21418–21426
- Chen L, Thiruchelvam MJ, Madura K, Richfield EK (2006) Proteasome dysfunction in aged human alpha-synuclein transgenic mice. *Neurobiol Dis* 23:120–126
- Chesselet MF, Fleming S, Mortazavi F, Meurers B (2008) Strengths and limitations of genetic mouse models of Parkinson's disease. *Parkinsonism Relat Disord* 14(Suppl 2):S84–S87
- Churchill GA, Airey DC, Allayee H, Angel JM, Attie AD, Beatty J, Beavis WD, Belknap JK, Bennett B, Berrettini W et al (2004) The Collaborative Cross, a community resource for the genetic analysis of complex traits. *Nat Genet* 36:1133–1137
- Cookson MR, Hardy J, Lewis PA (2008) Genetic neuropathology of Parkinson's disease. *Int J Clin Exp Pathol* 1:217–231
- D'Costa DF, Sheehan LJ, Phillips PA, Moore-Smith B (1995) The levodopa test in Parkinson's disease. *Age Ageing* 24:210–212
- Daher JP, Ying M, Banerjee R, McDonald RS, Hahn MD, Yang L, Flint Beal M, Thomas B, Dawson VL, Dawson TM et al (2009) Conditional transgenic mice expressing C-terminally truncated human alpha-synuclein (alphaSyn119) exhibit reduced striatal dopamine without loss of nigrostriatal pathway dopaminergic neurons. *Mol Neurodegener* 4:34
- Davis GC, Williams AC, Markey SP, Ebert MH, Caine ED, Reichert CM, Kopin IJ (1979) Chronic Parkinsonism secondary to intravenous injection of meperidine analogues. *Psychiatry Res* 1:249–254
- Dawson TM, Dawson VL (2010) The role of parkin in familial and sporadic Parkinson's disease. *Mov Disord* 25(Suppl 1):S32–S39
- Del Sol A, Balling R, Hood L, Galas D (2010) Diseases as network perturbations. *Curr Opin Biotechnol* 21:566–571
- Di Fonzo A, Dekker MC, Montagna P, Baruzzi A, Yonova EH, Correia Guedes L, Szczerbinska A, Zhao T, Dubbel-Hulsman LO, Wouters CH et al (2009) FBXO7 mutations cause autosomal recessive, early-onset parkinsonian-pyramidal syndrome. *Neurology* 72:240–245
- Ekstrand MI, Galter D (2009) The MitoPark Mouse - an animal model of Parkinson's disease with impaired respiratory chain function in dopamine neurons. *Parkinsonism Relat Disord* 15(Suppl 3): S185–S188
- Ekstrand MI, Terzioglu M, Galter D, Zhu S, Hofstetter C, Lindqvist E, Thams S, Bergstrand A, Hansson FS, Trifunovic A et al (2007) Progressive parkinsonism in mice with respiratory-chain-deficient dopamine neurons. *Proc Natl Acad Sci USA* 104:1325–1330
- Emborg ME (2004) Evaluation of animal models of Parkinson's disease for neuroprotective strategies. *J Neurosci Methods* 139: 121–143
- Fearnley JM, Lees AJ (1991) Ageing and Parkinson's disease: substantia nigra regional selectivity. *Brain* 114(Pt 5):2283–2301
- Fernagut PO, Chesselet MF (2004) Alpha-synuclein and transgenic mouse models. *Neurobiol Dis* 17:123–130
- Fernagut PO, Hutson CB, Fleming SM, Tetreault NA, Salcedo J, Masliah E, Chesselet MF (2007) Behavioral and histopathological consequences of paraquat intoxication in mice: effects of alpha-synuclein over-expression. *Synapse* 61:991–1001
- Fleming SM, Salcedo J, Fernagut PO, Rockenstein E, Masliah E, Levine MS, Chesselet MF (2004) Early and progressive sensorimotor anomalies in mice overexpressing wild-type human alpha-synuclein. *J Neurosci* 24:9434–9440
- Fleming SM, Salcedo J, Hutson CB, Rockenstein E, Masliah E, Levine MS, Chesselet MF (2006) Behavioral effects of dopaminergic agonists in transgenic mice overexpressing human wildtype alpha-synuclein. *Neuroscience* 142:1245–1253
- Fleming SM, Tetreault NA, Mulligan CK, Hutson CB, Masliah E, Chesselet MF (2008) Olfactory deficits in mice overexpressing human wildtype alpha-synuclein. *Eur J Neurosci* 28:247–256
- Forno LS (1996) Neuropathology of Parkinson's disease. *J Neuropathol Exp Neurol* 55:259–272
- Frank-Cannon TC, Tran T, Ruhn KA, Martinez TN, Hong J, Marvin M, Hartley M, Trevino I, O'Brien DE, Casey B et al (2008) Parkin deficiency increases vulnerability to inflammation-related nigral degeneration. *J Neurosci* 28:10825–10834
- Frasier M, Walzer M, McCarthy L, Magnuson D, Lee JM, Haas C, Kahle P, Wolozin B (2005) Tau phosphorylation increases in symptomatic mice overexpressing A30P alpha-synuclein. *Exp Neurol* 192:274–287
- Freichel C, Neumann M, Ballard T, Muller V, Woolley M, Ozmen L, Borroni E, Kretschmar HA, Haass C, Spooren W et al (2007) Age-dependent cognitive decline and amygdala pathology in alpha-synuclein transgenic mice. *Neurobiol Aging* 28:1421–1435
- Fujishiro H, Frigerio R, Burnett M, Klos KJ, Josephs KA, Delledonne A, Parisi JE, Ahlskog JE, Dickson DW (2008) Cardiac sympathetic denervation correlates with clinical and pathologic stages of Parkinson's disease. *Mov Disord* 23:1085–1092
- Galter D, Pernold K, Yoshitake T, Lindqvist E, Hoffer B, Kehr J, Larsson NG, Olson L (2010) MitoPark mice mirror the slow

- progression of key symptoms and L-DOPA response in Parkinson's disease. *Genes Brain Behav* 9:173–181
- Gao HM, Kotzbauer PT, Uryu K, Leight S, Trojanowski JQ, Lee VM (2008) Neuroinflammation and oxidation/nitration of alpha-synuclein linked to dopaminergic neurodegeneration. *J Neurosci* 28:7687–7698
- Gash DM, Rutland K, Hudson NL, Sullivan PG, Bing G, Cass WA, Pandya JD, Liu M, Choi DY, Hunter RL et al (2008) Trichloroethylene: Parkinsonism and complex 1 mitochondrial neurotoxicity. *Ann Neurol* 63:184–192
- Gasser T (2009a) Genomic and proteomic biomarkers for Parkinson disease. *Neurology* 72:S27–S31
- Gasser T (2009b) Molecular pathogenesis of Parkinson disease: insights from genetic studies. *Expert Rev Mol Med* 11:e22
- Gautier CA, Kitada T, Shen J (2008) Loss of PINK1 causes mitochondrial functional defects and increased sensitivity to oxidative stress. *Proc Natl Acad Sci USA* 105:11364–11369
- George S, van den Buuse M, San Mok S, Masters CL, Li QX, Culvenor JG (2008) Alpha-synuclein transgenic mice exhibit reduced anxiety-like behaviour. *Exp Neurol* 210:788–792
- Ghosh A, Roy A, Matras J, Brahmachari S, Gendelman HE, Pahan K (2009) Simvastatin inhibits the activation of p21ras and prevents the loss of dopaminergic neurons in a mouse model of Parkinson's disease. *J Neurosci* 29:13543–13556
- Giasson BI, Duda JE, Quinn SM, Zhang B, Trojanowski JQ, Lee VM (2002) Neuronal alpha-synucleinopathy with severe movement disorder in mice expressing A53T human alpha-synuclein. *Neuron* 34:521–533
- Giovanni A, Sieber BA, Heikkila RE, Sonsalla PK (1994a) Studies on species sensitivity to the dopaminergic neurotoxin 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine. Part 1: Systemic administration. *J Pharmacol Exp Ther* 270:1000–1007
- Giovanni A, Sonsalla PK, Heikkila RE (1994b) Studies on species sensitivity to the dopaminergic neurotoxin 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine. Part 2: Central administration of 1-methyl-4-phenylpyridinium. *J Pharmacol Exp Ther* 270:1008–1014
- Giovannone B, Tsiaras WG, de la Monte S, Klysik J, Lautier C, Karashchuk G, Goldwurm S, Smith RJ (2009) *GIGYF2* gene disruption in mice results in neurodegeneration and altered insulin-like growth factor signaling. *Hum Mol Genet* 18:4629–4639
- Gispert S, Del Turco D, Garrett L, Chen A, Bernard DJ, Hamm-Clement J, Korf HW, Deller T, Braak H, Auburger G et al (2003) Transgenic mice expressing mutant A53T human alpha-synuclein show neuronal dysfunction in the absence of aggregate formation. *Mol Cell Neurosci* 24:419–429
- Gispert S, Ricciardi F, Kurz A, Azizov M, Hoepken HH, Becker D, Voos W, Leuner K, Muller WE, Kudin AP et al (2009) Parkinson phenotype in aged PINK1-deficient mice is accompanied by progressive mitochondrial dysfunction in absence of neurodegeneration. *PLoS One* 4:e5777
- Goldberg MS, Fleming SM, Palacino JJ, Cepeda C, Lam HA, Bhatnagar A, Meloni EG, Wu N, Ackerson LC, Klapstein GJ et al (2003) Parkin-deficient mice exhibit nigrostriatal deficits but not loss of dopaminergic neurons. *J Biol Chem* 278:43628–43635
- Goldman SM (2010) Trichloroethylene and Parkinson's disease: dissolving the puzzle. *Expert Rev Neurother* 10:835–837
- Gomez-Isla T, Irizarry MC, Mariash A, Cheung B, Soto O, Schrump S, Sondel J, Kotilinek L, Day J, Schwarzschild MA et al (2003) Motor dysfunction and gliosis with preserved dopaminergic markers in human alpha-synuclein A30P transgenic mice. *Neurobiol Aging* 24:245–258
- Good CH, Hoffman AF, Hoffer BJ, Chefer VI, Shippenberg TS, Backman CM, Larsson NG, Olson L, Gellhaar S, Galter D et al (2011) Impaired nigrostriatal function precedes behavioral deficits in a genetic mitochondrial model of Parkinson's disease. *FASEB J* 25(4):1333–1344
- Graham DR, Sidhu A (2010) Mice expressing the A53T mutant form of human alpha-synuclein exhibit hyperactivity and reduced anxiety-like behavior. *J Neurosci Res* 88:1777–1783
- Greffard S, Verny M, Bonnet AM, Beinis JY, Gallinari C, Meaume S, Piette F, Hauw JJ, Duyckaerts C (2006) Motor score of the Unified Parkinson Disease Rating Scale as a good predictor of Lewy body-associated neuronal loss in the substantia nigra. *Arch Neurol* 63:584–588
- Greten-Harrison B, Polydoro M, Morimoto-Tomita M, Diao L, Williams AM, Nie EH, Makani S, Tian N, Castillo PE, Buchman VL et al (2010) alphabetagamma-Synuclein triple knockout mice reveal age-dependent neuronal dysfunction. *Proc Natl Acad Sci USA* 107:19573–19578
- Guateeo E, Bengtson CP, Bernardi G, Mercuri NB (2004) Voltage-gated calcium channels mediate intracellular calcium increase in weaver dopaminergic neurons during stimulation of D2 and GABAB receptors. *J Neurophysiol* 92:3368–3374
- Guehl D, Bezard E, Dovero S, Boraud T, Bioulac B, Gross C (1999) Trichloroethylene and parkinsonism: a human and experimental observation. *Eur J Neurol* 6:609–611
- Gureviciene I, Gurevicius K, Tanila H (2007) Role of alpha-synuclein in synaptic glutamate release. *Neurobiol Dis* 28:83–89
- Gureviciene I, Gurevicius K, Tanila H (2009) Aging and alpha-synuclein affect synaptic plasticity in the dentate gyrus. *J Neural Transm* 116:13–22
- Halliday GM, Ophof A, Broe M, Jensen PH, Kettle E, Fedorow H, Cartwright MI, Griffiths FM, Shepherd CE, Double KL (2005) Alpha-synuclein redistributes to neuromelanin lipid in the substantia nigra early in Parkinson's disease. *Brain* 128:2654–2664
- Hamani C, Nobrega JN, Lozano AM (2010) Deep brain stimulation in clinical practice and in animal models. *Clin Pharmacol Ther* 88:559–562
- Harvey BK, Wang Y, Hoffer BJ (2008) Transgenic rodent models of Parkinson's disease. *Acta Neurochir Suppl* 101:89–92
- Hashimoto M, Rockenstein E, Masliah E (2003) Transgenic models of alpha-synuclein pathology: past, present, and future. *Ann N Y Acad Sci* 991:171–188
- Haubenberger D, Reinthaler E, Mueller JC, Pirker W, Katzenschlager R, Froehlich R, Bruecke T, Daniel G, Auff E, Zimprich A (2011) Association of transcription factor polymorphisms PITX3 and EN1 with Parkinson's disease. *Neurobiol Aging* 32:302–307
- Haugarvoll K, Toft M, Skipper L, Heckman MG, Crook JE, Soto A, Ross OA, Hulihan MM, Kachergus JM, Sando SB et al (2009) Fine-mapping and candidate gene investigation within the PARK10 locus. *Eur J Hum Genet* 17:336–343
- Hawkes CH (2008) The prodromal phase of sporadic Parkinson's disease: does it exist and if so how long is it? *Mov Disord* 23:1799–1807
- Hawkes CH, Del Tredici K, Braak H (2010) A timeline for Parkinson's disease. *Parkinsonism Relat Disord* 16:79–84
- Healy DG, Falchi M, O'Sullivan SS, Bonifati V, Durr A, Bressman S, Brice A, Aasly J, Zabetian CP, Goldwurm S et al (2008) Phenotype, genotype, and worldwide genetic penetrance of LRRK2-associated Parkinson's disease: a case-control study. *Lancet Neurol* 7:583–590
- Hirth F (2010) *Drosophila melanogaster* in the study of human neurodegeneration. *CNS Neurol Disord Drug Targets* 9:504–523
- Hwang DY, Fleming SM, Ardayfio P, Moran-Gates T, Kim H, Tarazi FI, Chesselet MF, Kim KS (2005) 3, 4-Dihydroxyphenylalanine reverses the motor deficits in Pitx3-deficient aphakia mice: behavioral characterization of a novel genetic model of Parkinson's disease. *J Neurosci* 25:2132–2137

- Hwang DY, Hong S, Jeong JW, Choi S, Kim H, Kim J, Kim KS (2009) Vesicular monoamine transporter 2 and dopamine transporter are molecular targets of Pitx3 in the ventral midbrain dopamine neurons. *J Neurochem* 111:1202–1212
- Ibanez P, Bonnet AM, Debarges B, Lohmann E, Tison F, Pollak P, Agid Y, Durr A, Brice A (2004) Causal relation between alpha-synuclein gene duplication and familial Parkinson's disease. *Lancet* 364:1169–1171
- Ikeda M, Kawarabayashi T, Harigaya Y, Sasaki A, Yamada S, Matsubara E, Murakami T, Tanaka Y, Kurata T, Wuhua X et al (2009) Motor impairment and aberrant production of neurochemicals in human alpha-synuclein A30P + A53T transgenic mice with alpha-synuclein pathology. *Brain Res* 1250:232–241
- Inden M, Kitamura Y, Abe M, Tamaki A, Takata K, Taniguchi T (2011) Parkinsonian rotenone mouse model: reevaluation of long-term administration of rotenone in C57BL/6 mice. *Biol Pharm Bull* 34:92–96
- International Parkinson Disease Genomics Consortium, Nalls MA, Plagnol V, Hernandez DG, Sharma M, Sheerin UM, Saad M, Simón-Sánchez J, Schulte C, Lesage S, Sveinbjörnsdóttir S, Stefánsson K, Martinez M, Hardy J, Heutink P, Brice A, Gasser T, Singleton AB, Wood NW (2011) Imputation of sequence variants for identification of genetic risks for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet* 377(9766):641–649
- Irwin I, Finnegan KT, Delaney LE, Di Monte D, Langston JW (1992) The relationships between aging, monoamine oxidase, striatal dopamine and the effects of MPTP in C57BL/6 mice: a critical reassessment. *Brain Res* 572:224–231
- Isaias IU, Antonini A (2010) Single-photon emission computed tomography in diagnosis and differential diagnosis of Parkinson's disease. *Neurodegener Dis* 7:319–329
- Itier JM, Ibanez P, Mena MA, Abbas N, Cohen-Salmon C, Bohme GA, Laville M, Pratt J, Corti O, Pradier L et al (2003) Parkin gene inactivation alters behaviour and dopamine neurotransmission in the mouse. *Hum Mol Genet* 12:2277–2291
- Jankovic J (2008) Parkinson's disease: clinical features and diagnosis. *J Neurol Neurosurg Psychiatry* 79:368–376
- Kahle PJ, Neumann M, Ozmen L, Muller V, Jacobsen H, Schindzielorz A, Okochi M, Leimer U, van Der Putten H, Probst A et al (2000) Subcellular localization of wild-type and Parkinson's disease-associated mutant alpha-synuclein in human and transgenic mouse brain. *J Neurosci* 20:6365–6373
- Katzenbach R, Lees AJ (2004) Olfaction and Parkinson's syndromes: its role in differential diagnosis. *Curr Opin Neurol* 17:417–423
- Kim RH, Smith PD, Aleyasin H, Hayley S, Mount MP, Pownall S, Wakeham A, You-Ten AJ, Kalia SK, Horne P et al (2005) Hypersensitivity of DJ-1-deficient mice to 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine (MPTP) and oxidative stress. *Proc Natl Acad Sci USA* 102:5215–5220
- Kitada T, Pisani A, Porter DR, Yamaguchi H, Tscherter A, Martella G, Bonsi P, Zhang C, Pothos EN, Shen J (2007) Impaired dopamine release and synaptic plasticity in the striatum of PINK1-deficient mice. *Proc Natl Acad Sci USA* 104:11441–11446
- Kitada T, Tong Y, Gautier CA, Shen J (2009) Absence of nigral degeneration in aged parkin/DJ-1/PINK1 triple knockout mice. *J Neurochem* 111:696–702
- Koob AO, Ubhi K, Paulsson JF, Kelly J, Rockenstein E, Mante M, Adame A, Masliah E (2010) Lovastatin ameliorates alpha-synuclein accumulation and oxidation in transgenic mouse models of alpha-synucleinopathies. *Exp Neurol* 221:267–274
- Kruger R, Kuhn W, Muller T, Woitalla D, Graeber M, Kosel S, Przuntek H, Epplen JT, Schols L, Riess O (1998) Ala30Pro mutation in the gene encoding alpha-synuclein in Parkinson's disease. *Nat Genet* 18:106–108
- Kuo YM, Li Z, Jiao Y, Gaborit N, Pani AK, Orrison BM, Bruneau BG, Giasson BI, Smeyne RJ, Gershon MD et al (2010) Extensive enteric nervous system abnormalities in mice transgenic for artificial chromosomes containing Parkinson disease-associated alpha-synuclein gene mutations precede central nervous system changes. *Hum Mol Genet* 19:1633–1650
- Lang AE, Obeso JA (2004) Challenges in Parkinson's disease: restoration of the nigrostriatal dopamine system is not enough. *Lancet Neurol* 3:309–316
- Langston JW (2006) The Parkinson's complex: parkinsonism is just the tip of the iceberg. *Ann Neurol* 59:591–596
- Langston JW, Ballard P, Tetrud JW, Irwin I (1983) Chronic Parkinsonism in humans due to a product of meperidine-analog synthesis. *Science* 219:979–980
- Larsen K, Hedegaard C, Bertelsen MF, Bendixen C (2009) Threonine 53 in alpha-synuclein is conserved in long-living non-primate animals. *Biochem Biophys Res Commun* 387:602–605
- Lee MK, Stirling W, Xu Y, Xu X, Qui D, Mandir AS, Dawson TM, Copeland NG, Jenkins NA, Price DL (2002) Human alpha-synuclein-harboring familial Parkinson's disease-linked Ala-53 → Thr mutation causes neurodegenerative disease with alpha-synuclein aggregation in transgenic mice. *Proc Natl Acad Sci USA* 99:8968–8973
- Lees AJ (2009) The Parkinson chimera. *Neurology* 72:S2–S11
- Li W, West N, Colla E, Pletnikova O, Troncoso JC, Marsh L, Dawson TM, Jakala P, Hartmann T, Price DL et al (2005) Aggregation promoting C-terminal truncation of alpha-synuclein is a normal cellular process and is enhanced by the familial Parkinson's disease-linked mutations. *Proc Natl Acad Sci USA* 102:2162–2167
- Li YJ, Deng J, Mayhew GM, Grimsley JW, Huo X, Vance JM (2007) Investigation of the *PARK10* gene in Parkinson disease. *Ann Hum Genet* 71:639–647
- Li Y, Liu W, Oo TF, Wang L, Tang Y, Jackson-Lewis V, Zhou C, Geghman K, Bogdanov M, Przedborski S et al (2009) Mutant LRRK2(R1441G) BAC transgenic mice recapitulate cardinal features of Parkinson's disease. *Nat Neurosci* 12:826–828
- Li X, Patel JC, Wang J, Avshalumov MV, Nicholson C, Buxbaum JD, Elder GA, Rice ME, Yue Z (2010) Enhanced striatal dopamine transmission and motor performance with LRRK2 overexpression in mice is eliminated by familial Parkinson's disease mutation G2019S. *J Neurosci* 30:1788–1797
- Lim Y, Kehm VM, Li C, Trojanowski JQ, Lee VM (2010) Forebrain overexpression of alpha-synuclein leads to early postnatal hippocampal neuron loss and synaptic disruption. *Exp Neurol* 221:86–97
- Lin X, Parisiadou L, Gu XL, Wang L, Shim H, Sun L, Xie C, Long CX, Yang WJ, Ding J et al (2009) Leucine-rich repeat kinase 2 regulates the progression of neuropathology induced by Parkinson's-disease-related mutant alpha-synuclein. *Neuron* 64:807–827
- Liu MJ, Liu ML, Shen YF, Kim JM, Lee BH, Lee YS, Hong ST (2007) Transgenic mice with neuron-specific overexpression of HtrA2/Omi suggest a neuroprotective role for HtrA2/Omi. *Biochem Biophys Res Commun* 362:295–300
- Liu M, Choi DY, Hunter RL, Pandya JD, Cass WA, Sullivan PG, Kim HC, Gash DM, Bing G (2010a) Trichloroethylene induces dopaminergic neurodegeneration in Fisher 344 rats. *J Neurochem* 112:773–783
- Liu P, Wang X, Gao N, Zhu H, Dai X, Xu Y, Ma C, Huang L, Liu Y, Qin C (2010b) G protein-coupled receptor kinase 5, overexpressed in the alpha-synuclein up-regulation model of Parkinson's disease, regulates bcl-2 expression. *Brain Res* 1307:134–141

- Lu XH, Fleming SM, Meurers B, Ackerson LC, Mortazavi F, Lo V, Hernandez D, Sulzer D, Jackson GR, Maidment NT et al (2009) Bacterial artificial chromosome transgenic mice expressing a truncated mutant parkin exhibit age-dependent hypokinetic motor deficits, dopaminergic neuron degeneration, and accumulation of proteinase K-resistant alpha-synuclein. *J Neurosci* 29: 1962–1976
- Magen I, Chesselet MF (2010) Genetic mouse models of Parkinson's disease: The state of the art. *Prog Brain Res* 184:53–87
- Manning-Bog AB, McCormack AL, Li J, Uversky VN, Fink AL, Di Monte DA (2002) The herbicide paraquat causes up-regulation and aggregation of alpha-synuclein in mice: paraquat and alpha-synuclein. *J Biol Chem* 277:1641–1644
- Manning-Bog AB, McCormack AL, Purisai MG, Bolin LM, Di Monte DA (2003) Alpha-synuclein overexpression protects against paraquat-induced neurodegeneration. *J Neurosci* 23:3095–3099
- Maries E, Dass B, Collier TJ, Kordower JH, Steele-Collier K (2003) The role of alpha-synuclein in Parkinson's disease: insights from animal models. *Nat Rev Neurosci* 4:727–738
- Markesberry WR, Jicha GA, Liu H, Schmitt FA (2009) Lewy body pathology in normal elderly subjects. *J Neuropathol Exp Neurol* 68:816–822
- Martella G, Platania P, Vita D, Sciamanna G, Cuomo D, Tassone A, Tscherter A, Kitada T, Bonsi P, Shen J et al (2009) Enhanced sensitivity to group II mGlu receptor activation at corticostriatal synapses in mice lacking the familial parkinsonism-linked genes PINK1 or Parkin. *Exp Neurol* 215:388–396
- Martin LJ, Pan Y, Price AC, Sterling W, Copeland NG, Jenkins NA, Price DL, Lee MK (2006) Parkinson's disease alpha-synuclein transgenic mice develop neuronal mitochondrial degeneration and cell death. *J Neurosci* 26:41–50
- Martins LM, Morrison A, Klupsch K, Fedele V, Moisoi N, Teismann P, Abuin A, Grau E, Geppert M, Livi GP et al (2004) Neuroprotective role of the Reaper-related serine protease HtrA2/Omi revealed by targeted deletion in mice. *Mol Cell Biol* 24:9848–9862
- Marxreiter F, Nuber S, Kandasamy M, Klucken J, Aigner R, Burgmayer R, Couillard-Despres S, Riess O, Winkler J, Winner B (2009) Changes in adult olfactory bulb neurogenesis in mice expressing the A30P mutant form of alpha-synuclein. *Eur J Neurosci* 29:879–890
- Mascaro AL, Sacconi L, Pavone FS (2010) Multi-photon nanosurgery in live brain. *Front neuroenergetics* 2, pii 21
- Maskri L, Zhu X, Fritzen S, Kuhn K, Ullmer C, Engels P, Andriske M, Stichel CC, Lubbert H (2004) Influence of different promoters on the expression pattern of mutated human alpha-synuclein in transgenic mice. *Neurodegener Dis* 1:255–265
- Masliah E, Rockenstein E, Veinbergs I, Mallory M, Hashimoto M, Takeda A, Sagara Y, Sisk A, Mucke L (2000) Dopaminergic loss and inclusion body formation in alpha-synuclein mice: implications for neurodegenerative disorders. *Science* 287:1265–1269
- Matsuoka Y, Vila M, Lincoln S, McCormack A, Picciano M, LaFrancois J, Yu X, Dickson D, Langston WJ, McGowan E et al (2001) Lack of nigral pathology in transgenic mice expressing human alpha-synuclein driven by the tyrosine hydroxylase promoter. *Neurobiol Dis* 8:535–539
- McNeill TH, Koek LL (1990) Differential effects of advancing age on neurotransmitter cell loss in the substantia nigra and striatum of C57BL/6 N mice. *Brain Res* 521:107–117
- Melrose HL, Dachsel JC, Behrouz B, Lincoln SJ, Yue M, Hinkle KM, Kent CB, Korvatska E, Taylor JP, Witten L et al (2010) Impaired dopaminergic neurotransmission and microtubule-associated protein tau alterations in human LRRK2 transgenic mice. *Neurobiol Dis* 40:503–517
- Menendez J, Rodriguez-Navarro JA, Solano RM, Casarejos MJ, Rodal I, Guerrero R, Sanchez MP, Avila J, Mena MA, de Yebenes JG (2006) Suppression of Parkin enhances nigrostriatal and motor neuron lesion in mice over-expressing human-mutated tau protein. *Hum Mol Genet* 15:2045–2058
- Meredith GE, Totterdell S, Potashkin JA, Surmeier DJ (2008) Modeling PD pathogenesis in mice: advantages of a chronic MPTP protocol. *Parkinsonism Relat Disord* 14(Suppl 2):S112–S115
- Miller RM, Kiser GL, Kaysser-Kranich T, Casaceli C, Colla E, Lee MK, Palaniappan C, Federoff HJ (2007) Wild-type and mutant alpha-synuclein induce a multi-component gene expression profile consistent with shared pathophysiology in different transgenic mouse models of PD. *Exp Neurol* 204:421–432
- Morais VA, Verstreken P, Roethig A, Smet J, Snellinx A, Vanbrabant M, Haddad D, Frezza C, Mandemakers W, Vogt-Weisenhorn D et al (2009) Parkinson's disease mutations in PINK1 result in decreased Complex I activity and deficient synaptic function. *EMBO Mol Med* 1:99–111
- Nass R, Merchant KM, Ryan T (2008) *Caenorhabditis elegans* in Parkinson's disease drug discovery: addressing an unmet medical need. *Mol Interv* 8:284–293
- Neumann M, Kahle PJ, Giasson BI, Ozmen L, Borroni E, Spooren W, Muller V, Odoy S, Fujiwara H, Hasegawa M et al (2002) Misfolded proteinase K-resistant hyperphosphorylated alpha-synuclein in aged transgenic mice with locomotor deterioration and in human alpha-synucleinopathies. *J Clin Invest* 110:1429–1439
- Nieto M, Gil-Bea FJ, Dalfo E, Cuadrado M, Cabodevilla F, Sanchez B, Catena S, Sesma T, Ribe E, Ferrer I et al (2006) Increased sensitivity to MPTP in human alpha-synuclein A30P transgenic mice. *Neurobiol Aging* 27:848–856
- Nishi A, Ishii A, Takahashi A, Shiroishi T, Koide T (2010) QTL analysis of measures of mouse home-cage activity using B6/MSM consomic strains. *Mamm Genome* 21:477–485
- Norris EH, Uryu K, Leight S, Giasson BI, Trojanowski JQ, Lee VM (2007) Pesticide exposure exacerbates alpha-synucleinopathy in an A53T transgenic mouse model. *Am J Pathol* 170:658–666
- Nuber S, Petrasch-Parwez E, Winner B, Winkler J, von Horsten S, Schmidt T, Boy J, Kuhn M, Nguyen HP, Teismann P et al (2008) Neurodegeneration and motor dysfunction in a conditional model of Parkinson's disease. *J Neurosci* 28:2471–2484
- O'Gorman S, Dagenais NA, Qian M, Marchuk Y (1997) Protamine-Cre recombinase transgenes efficiently recombine target sequences in the male germ line of mice, but not in embryonic stem cells. *Proc Natl Acad Sci USA* 94:14602–14607
- Oksman M, Tanila H, Yavich L (2009) Behavioural and neurochemical response of alpha-synuclein A30P transgenic mice to the effects of L-DOPA. *Neuropharmacology* 56:647–652
- Ono K, Ikemoto M, Kawarabayashi T, Ikeda M, Nishinakagawa T, Hosokawa M, Shoji M, Takahashi M, Nakashima M (2009) A chemical chaperone, sodium 4-phenylbutyric acid, attenuates the pathogenic potency in human alpha-synuclein A30P + A53T transgenic mice. *Parkinsonism Relat Disord* 15:649–654
- Paisan-Ruiz C, Jain S, Evans EW, Gilks WP, Simon J, van der Brug M, Lopez de Munain A, Aparicio S, Gil AM, Khan N et al (2004) Cloning of the gene containing mutations that cause PARK8-linked Parkinson's disease. *Neuron* 44:595–600
- Paisan-Ruiz C, Guevara R, Federoff M, Hanagasi H, Sina F, Elahi E, Schneider SA, Schwingenschuh P, Bajaj N, Emre M et al (2010) Early-onset L-dopa-responsive parkinsonism with pyramidal signs due to ATP13A2, PLA2G6, FBXO7 and spatacsin mutations. *Mov Disord* 25:1791–1800
- Palacino JJ, Sagi D, Goldberg MS, Krauss S, Motz C, Wacker M, Klose J, Shen J (2004) Mitochondrial dysfunction and oxidative damage in parkin-deficient mice. *J Biol Chem* 279:18614–18622

- Pankratz N, Nichols WC, Uniacke SK, Halter C, Murrell J, Rudolph A, Shults CW, Conneally PM, Foroud T (2003) Genome-wide linkage analysis and evidence of gene-by-gene interactions in a sample of 362 multiplex Parkinson disease families. *Hum Mol Genet* 12:2599–2608
- Pankratz N, Wilk JB, Latourelle JC, DeStefano AL, Halter C, Pugh EW, Doheny KF, Gusella JF, Nichols WC, Foroud T et al (2009) Genomewide association study for susceptibility genes contributing to familial Parkinson disease. *Hum Genet* 124:593–605
- Periquet M, Corti O, Jacquier S, Brice A (2005) Proteomic analysis of parkin knockout mice: alterations in energy metabolism, protein handling and synaptic function. *J Neurochem* 95:1259–1276
- Pienaar IS, Daniels WM, Gotz J (2008) Neoproteomics as a promising tool in Parkinson's disease research. *J Neural Transm* 115:1413–1430
- Plaas M, Karis A, Innos J, Rebane E, Baekelandt V, Vaermann A, Luuk H, Vasar E, Koks S (2008) Alpha-synuclein A30P point-mutation generates age-dependent nigrostriatal deficiency in mice. *J Physiol Pharmacol* 59:205–216
- Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, Pike B, Root H, Rubenstein J, Boyer R et al (1997) Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science* 276:2045–2047
- Poon HF, Frasier M, Shreve N, Calabrese V, Wolozin B, Butterfield DA (2005) Mitochondrial associated metabolic proteins are selectively oxidized in A30P alpha-synuclein transgenic mice—a model of familial Parkinson's disease. *Neurobiol Dis* 18:492–498
- Prasad K, Tarasewicz E, Ohman Strickland PA, O'Neill M, Mitchell SN, Merchant K, Tep S, Hilton K, Datwani A, Buttini M et al (2011) Biochemical and morphological consequences of human alpha-synuclein expression in a mouse alpha-synuclein null background. *Eur J Neurosci* 33:642–652
- Rajput AH, Voll A, Rajput ML, Robinson CA, Rajput A (2009) Course in Parkinson disease subtypes: A 39-year clinicopathologic study. *Neurology* 73:206–212
- Ramirez A, Heimbach A, Grundemann J, Stiller B, Hampshire D, Cid LP, Goebel I, Mubaidin AF, Wriekat AL, Roeper J et al (2006) Hereditary parkinsonism with dementia is caused by mutations in ATP13A2, encoding a lysosomal type 5 P-type ATPase. *Nat Genet* 38:1184–1191
- Rascol O (2009) "Disease-modification" trials in Parkinson disease: target populations, endpoints and study design. *Neurology* 72:S51–S58
- Ravenstijn PG, Merlini M, Hameetman M, Murray TK, Ward MA, Lewis H, Ball G, Mottart C, de Goyet C, Lemarchand T et al (2008) The exploration of rotenone as a toxin for inducing Parkinson's disease in rats, for application in BBB transport and PK-PD experiments. *J Pharmacol Toxicol Methods* 57:114–130
- Richfield EK, Thiruchelvam MJ, Cory-Slechta DA, Wuertzer C, Gainetdinov RR, Caron MG, Di Monte DA, Federoff HJ (2002) Behavioral and neurochemical effects of wild-type and mutated human alpha-synuclein in transgenic mice. *Exp Neurol* 175:35–48
- Ritz BR, Mantripragada AD, Costello S, Lincoln SJ, Farrer MJ, Cockburn M, Bronstein J (2009) Dopamine transporter genetic variants and pesticides in Parkinson's disease. *Environ Health Perspect* 117:964–969
- Robinson PA (2010) Understanding the molecular basis of Parkinson's disease, identification of biomarkers and routes to therapy. *Expert Rev Proteomics* 7:565–578
- Rockenstein E, Mallory M, Hashimoto M, Song D, Shults CW, Lang I, Masliah E (2002) Differential neuropathological alterations in transgenic mice expressing alpha-synuclein from the platelet-derived growth factor and Thy-1 promoters. *J Neurosci Res* 68:568–578
- Rodriguez-Navarro JA, Casarejos MJ, Menendez J, Solano RM, Rodal I, Gomez A, Yebenes JG, Mena MA (2007) Mortality, oxidative stress and tau accumulation during ageing in parkin null mice. *J Neurochem* 103:98–114
- Satake W, Nakabayashi Y, Mizuta I, Hirota Y, Ito C, Kubo M, Kawaguchi T, Tsunoda T, Watanabe M, Takeda A et al (2009) Genome-wide association study identifies common variants at four loci as genetic risk factors for Parkinson's disease. *Nat Genet* 41:1303–1307
- Savica R, Rocca WA, Ahlskog JE (2010) When does Parkinson disease start? *Arch Neurol* 67:798–801
- Schapira AH (2006) Etiology of Parkinson's disease. *Neurology* 66: S10–S23
- Schapira AH, Tolosa E (2010) Molecular and clinical prodrome of Parkinson disease: implications for treatment. *Nat Rev Neurol* 6:309–317
- Schapira AH, Agid Y, Barone P, Jenner P, Lemke MR, Poewe W, Rascol O, Reichmann H, Tolosa E (2009) Perspectives on recent advances in the understanding and treatment of Parkinson's disease. *Eur J Neurol* 16:1090–1099
- Schell H, Hasegawa T, Neumann M, Kahle PJ (2009) Nuclear and neuritic distribution of serine-129 phosphorylated alpha-synuclein in transgenic mice. *Neuroscience* 160:796–804
- Sedelis M, Hofele K, Schwarting RK, Huston JP, Belknap JK (2003) Chromosomal loci influencing the susceptibility to the parkinsonian neurotoxin 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine. *J Neurosci* 23:8247–8253
- Setsuie R, Wang YL, Mochizuki H, Osaka H, Hayakawa H, Ichihara N, Li H, Furuta A, Sano Y, Sun YJ et al (2007) Dopaminergic neuronal loss in transgenic mice expressing the Parkinson's disease-associated UCH-L1 I93 M mutant. *Neurochem Int* 50: 119–129
- Sharon R, Bar-Joseph I, Mirick GE, Serhan CN, Selkoe DJ (2003) Altered fatty acid composition of dopaminergic neurons expressing alpha-synuclein and human brains with alpha-synucleinopathies. *J Biol Chem* 278:49874–49881
- Sidransky E, Nalls MA, Aasly JO, Aharon-Peretz J, Annesi G, Barbosa ER, Bar-Shira A, Berg D, Bras J, Brice A et al (2009) Multicenter analysis of glucocerebrosidase mutations in Parkinson's disease. *N Engl J Med* 361:1651–1661
- Simon-Sanchez J, Schulte C, Bras JM, Sharma M, Gibbs JR, Berg D, Paisan-Ruiz C, Lichtner P, Scholz SW, Hernandez DG et al (2009) Genome-wide association study reveals genetic risk underlying Parkinson's disease. *Nat Genet* 41:1308–1312
- Singleton AB, Farrer M, Johnson J, Singleton A, Hague S, Kachergus J, Hulihan M, Peuralinna T, Dutra A, Nussbaum R et al (2003) alpha-Synuclein locus triplication causes Parkinson's disease. *Science* 302:841
- Song DD, Shults CW, Sisk A, Rockenstein E, Masliah E (2004) Enhanced substantia nigra mitochondrial pathology in human alpha-synuclein transgenic mice after treatment with MPTP. *Exp Neurol* 186:158–172
- Sotiriou E, Vassilatis DK, Vila M, Stefanis L (2010) Selective noradrenergic vulnerability in alpha-synuclein transgenic mice. *Neurobiol Aging* 31:2103–2114
- Spillantini MG, Crowther RA, Jakes R, Cairns NJ, Lantos PL, Goedert M (1998) Filamentous alpha-synuclein inclusions link multiple system atrophy with Parkinson's disease and dementia with Lewy bodies. *Neurosci Lett* 251:205–208
- Srivastava G, Singh K, Tiwari MN, Singh MP (2010) Proteomics in Parkinson's disease: current trends, translational snags and future possibilities. *Expert Rev Proteomics* 7:127–139
- Steg G (1964) Alpha-rigidity in reserpinated rats. *Experientia* 20:79–80
- Stichel CC, Zhu XR, Bader V, Linnartz B, Schmidt S, Lubbert H (2007) Mono- and double-mutant mouse models of Parkinson's

- disease display severe mitochondrial damage. *Hum Mol Genet* 16:2377–2393
- Su X, Maguire-Zeiss KA, Giuliano R, Prifti L, Venkatesh K, Federoff HJ (2008) Synuclein activates microglia in a model of Parkinson's disease. *Neurobiol Aging* 29:1690–1701
- Su X, Federoff HJ, Maguire-Zeiss KA (2009) Mutant alpha-synuclein overexpression mediates early proinflammatory activity. *Neurotox Res* 16:238–254
- Takahashi N, Miner LL, Sora I, Ujike H, Revay RS, Kostic V, Jackson-Lewis V, Przedborski S, Uhl GR (1997) VMAT2 knockout mice: heterozygotes display reduced amphetamine-conditioned reward, enhanced amphetamine locomotion, and enhanced MPTP toxicity. *Proc Natl Acad Sci USA* 94:9938–9943
- Takazawa C, Fujimoto K, Homma D, Sumi-Ichinose C, Nomura T, Ichinose H, Katoh S (2008) A brain-specific decrease of the tyrosine hydroxylase protein in sepiapterin reductase-null mice as a mouse model for Parkinson's disease. *Biochem Biophys Res Commun* 367:787–792
- Tan EK, Kwok HH, Tan LC, Zhao WT, Prakash KM, Au WL, Pavanni R, Ng YY, Satake W, Zhao Y et al (2010) Analysis of GWAS-linked loci in Parkinson disease reaffirms PARK16 as a susceptibility locus. *Neurology* 75:508–512
- Tanaka A (2010) Parkin-mediated selective mitochondrial autophagy, mitophagy: Parkin purges damaged organelles from the vital mitochondrial network. *FEBS Lett* 584:1386–1392
- Taylor TN, Caudle WM, Shepherd KR, Noorian A, Jackson CR, Iuvone PM, Weinshenker D, Greene JG, Miller GW (2009) Nonmotor symptoms of Parkinson's disease revealed in an animal model with reduced monoamine storage capacity. *J Neurosci* 29:8103–8113
- Thiruchelvam M, Richfield EK, Baggs RB, Tank AW, Cory-Slechta DA (2000) The nigrostriatal dopaminergic system as a preferential target of repeated exposures to combined paraquat and maneb: implications for Parkinson's disease. *J Neurosci* 20: 9207–9214
- Thiruchelvam MJ, Powers JM, Cory-Slechta DA, Richfield EK (2004) Risk factors for dopaminergic neuron loss in human alpha-synuclein transgenic mice. *Eur J Neurosci* 19:845–854
- Tofaris GK, Garcia Reitbock P, Humby T, Lambourne SL, O'Connell M, Ghetti B, Gossage H, Emson PC, Wilkinson LS, Goedert M et al (2006) Pathological changes in dopaminergic nerve cells of the substantia nigra and olfactory bulb in mice transgenic for truncated human alpha-synuclein(1–120): implications for Lewy body disorders. *J Neurosci* 26:3942–3950
- Tolosa E, Gaig C, Santamaria J, Compta Y (2009) Diagnosis and the premotor phase of Parkinson disease. *Neurology* 72:S12–S20
- Tong Y, Pisani A, Martella G, Karouani M, Yamaguchi H, Pothos EN, Shen J (2009) R1441C mutation in LRRK2 impairs dopaminergic neurotransmission in mice. *Proc Natl Acad Sci USA* 106:14622–14627
- Unger EL, Eve DJ, Perez XA, Reichenbach DK, Xu Y, Lee MK, Andrews AM (2006) Locomotor hyperactivity and alterations in dopamine neurotransmission are associated with overexpression of A53T mutant human alpha-synuclein in mice. *Neurobiol Dis* 21:431–443
- Ungerstedt U (1968) 6-Hydroxy-dopamine induced degeneration of central monoamine neurons. *Eur J Pharmacol* 5:107–110
- Ungerstedt U, Ljungberg T, Steg G (1974) Behavioral, physiological, and neurochemical changes after 6-hydroxydopamine-induced degeneration of the nigro-striatal dopamine neurons. *Adv Neurol* 5:421–426
- Van Dam D, De Deyn PP (2011) Animal models in the drug discovery pipeline for Alzheimer's disease. *Br J Pharmacol.* doi: 10.1111/j.1476-5381.2011.01299.x
- van der Putten H, Wiederhold KH, Probst A, Barbieri S, Mistl C, Danner S, Kauffmann S, Hofele K, Spooren WP, Ruegg MA et al (2000) Neuropathology in mice expressing human alpha-synuclein. *J Neurosci* 20:6021–6029
- Vogel G (2003) Genetics. Scientists dream of 1001 complex mice. *Science* 301:456–457
- Von Coelln R, Thomas B, Savitt JM, Lim KL, Sasaki M, Hess EJ, Dawson VL, Dawson TM (2004) Loss of locus coeruleus neurons and reduced startle in parkin null mice. *Proc Natl Acad Sci USA* 101:10744–10749
- Von Coelln R, Thomas B, Andraibi SA, Lim KL, Savitt JM, Saffary R, Stirling W, Bruno K, Hess EJ, Lee MK et al (2006) Inclusion body formation and neurodegeneration are parkin independent in a mouse model of alpha-synucleinopathy. *J Neurosci* 26:3685–3696
- Wakamatsu M, Ishii A, Iwata S, Sakagami J, Ukai Y, Ono M, Kanbe D, Muramatsu S, Kobayashi K, Iwatsubo T et al (2008a) Selective loss of nigral dopamine neurons induced by overexpression of truncated human alpha-synuclein in mice. *Neurobiol Aging* 29:574–585
- Wakamatsu M, Iwata S, Funakoshi T, Yoshimoto M (2008b) Dopamine receptor agonists reverse behavioral abnormalities of alpha-synuclein transgenic mouse, a new model of Parkinson's disease. *J Neurosci Res* 86:640–646
- Wang L, Fleming SM, Chesselet MF, Tache Y (2008) Abnormal colonic motility in mice overexpressing human wild-type alpha-synuclein. *Neuroreport* 19:873–876
- Watson JB, Hatami A, David H, Masliah E, Roberts K, Evans CE, Levine MS (2009) Alterations in corticostratial synaptic plasticity in mice overexpressing human alpha-synuclein. *Neuroscience* 159:501–513
- Wellstead P, Cloutier M (2011) An energy systems approach to Parkinson's disease. *Wiley Interdiscip Rev Syst Biol Med* 3:1–6
- Winner B, Lie DC, Rockenstein E, Aigner R, Aigner L, Masliah E, Kuhn HG, Winkler J (2004) Human wild-type alpha-synuclein impairs neurogenesis. *J Neuropathol Exp Neurol* 63:1155–1166
- Winner B, Rockenstein E, Lie DC, Aigner R, Mante M, Bogdahn U, Couillard-Despres S, Masliah E, Winkler J (2008) Mutant alpha-synuclein exacerbates age-related decrease of neurogenesis. *Neurobiol Aging* 29:913–925
- Winner B, Melrose HL, Zhao C, Hinkle KM, Yue M, Kent C, Braithwaite AT, Ogholikhan S, Aigner R, Winkler J et al (2011) Adult neurogenesis and neurite outgrowth are impaired in LRRK2 G2019S mice. *Neurobiol Dis* 41:706–716
- Witt SN, Flower TR (2006) alpha-Synuclein, oxidative stress and apoptosis from the perspective of a yeast model of Parkinson's disease. *FEMS Yeast Res* 6:1107–1116
- Wood-Kaczmar A, Gandhi S, Yao Z, Abramov AY, Miljan EA, Keen G, Stanyer L, Hargreaves I, Klupsch K, Deas E et al (2008) PINK1 is necessary for long term survival and mitochondrial function in human dopaminergic neurons. *PLoS One* 3:e2455
- Wu N, Joshi PR, Cepeda C, Masliah E, Levine MS (2010) Alpha-synuclein overexpression in mice alters synaptic communication in the corticostratial pathway. *J Neurosci Res* 88:1764–1776
- Yacoubian TA, Cantuti-Castelvetri I, Bouzou B, Asteris G, McLean PJ, Hyman BT, Standaert DG (2008) Transcriptional dysregulation in a transgenic model of Parkinson disease. *Neurobiol Dis* 29:515–528
- Yasuda T, Nihira T, Ren YR, Cao XQ, Wada K, Setsuie R, Kabuta T, Hattori N, Mizuno Y, Mochizuki H (2009) Effects of UCH-L1 on alpha-synuclein over-expression mouse model of Parkinson's disease. *J Neurochem* 108:932–944
- Yavich L, Oksman M, Tanila H, Kerokoski P, Hiltunen M, van Groen T, Puolivali J, Mannisto PT, Garcia-Horsman A, MacDonald E et al (2005) Locomotor activity and evoked dopamine release are

- reduced in mice overexpressing A30P-mutated human alpha-synuclein. *Neurobiol Dis* 20:303–313
- Yavich L, Jakala P, Tanila H (2006) Abnormal compartmentalization of norepinephrine in mouse dentate gyrus in alpha-synuclein knockout and A30P transgenic mice. *J Neurochem* 99:724–732
- Yoshino H, Tomiyama H, Tachibana N, Ogaki K, Li Y, Funayama M, Hashimoto T, Takashima S, Hattori N (2010) Phenotypic spectrum of patients with PLA2G6 mutation and PARK14-linked parkinsonism. *Neurology* 75:1356–1361
- Yu WH, Matsuoka Y, Sziraki I, Hashim A, Lafrancois J, Sershen H, Duff KE (2008) Increased dopaminergic neuron sensitivity to 1-methyl-4-phenyl-1, 2, 3, 6-tetrahydropyridine (MPTP) in transgenic mice expressing mutant A53T alpha-synuclein. *Neurochem Res* 33:902–911
- Zhou W, Milder JB, Freed CR (2008) Transgenic mice overexpressing tyrosine-to-cysteine mutant human alpha-synuclein: a progressive neurodegenerative model of diffuse Lewy body disease. *J Biol Chem* 283:9863–9870
- Zhu XR, Maskri L, Herold C, Bader V, Stichel CC, Gunturkun O, Lubbert H (2007) Non-motor behavioural impairments in parkin-deficient mice. *Eur J Neurosci* 26:1902–1911
- Zimprich A, Biskup S, Leitner P, Lichtner P, Farrer M, Lincoln S, Kachergus J, Hulihan M, Uitti RJ, Calne DB et al (2004) Mutations in LRRK2 cause autosomal-dominant parkinsonism with pleomorphic pathology. *Neuron* 44:601–607