Unique coral community in the Persian Gulf

Received: 6 September 2008 / Accepted: 18 October 2008 / Published online: 9 November 2008
© Springer-Verlag 2008

Fig. 1 **a** Shallow zone was exposed during low tide, **b** *Montipora* cf. *danai*, **c** *Platygyra daedalea*, **d** *Stylophora pistillata*, **e** *Sarcophyton* sp., **f** *Sinularia erecta*, **g** Schematic profile of the described area

A fringing coral community at Larak Island (26°53'N, 56°23'E), situated at the boundary between the Persian Gulf and Gulf of Oman in the Strait of Hormuz, is described. Little published information exists about Iranian reefs and their fauna and such a coral community has not yet been reported in the Persian Gulf (Shinn 1976; Sheppard and Sheppard 1991; Riegl 1999; Sheppard et al. 1992; Rezai et al. 2004).

Larak Island has mainly sandy shores with patchy reef development mostly in its northern and western parts. The described area extends about 1 km northeast of the Island. The tidal regime in the area is mixed and tidal range may reach 4 m during the year. Exposed to air for a few hours a day during spring low tide (Fig. 1a), the whole shallow zone (<5 m) coral community was dominated by soft corals, such as *Sinularia compressa*, *S. erecta* (Fig. 1f), other soft coral species, and hard corals, e.g., *Montipora* cf. *danai*, *Platygyra daedalea*, *Stylophora pistillata* (Fig. 1b–d), and *Favia favius*. Total live hard and soft coral cover in this zone was 18 and 15%, respectively. A deeper coral zone (5–8 m), dominated by *Acropora downingi* and *A. khayranensis*, occurred down slope to 8 m in depth (Fig. 1g) with 78.6% cover. Below that, *Sarcophyton* sp. (Fig. 1e), *Junceella juncea*, and *Goniopora* sp. become dominant and continue to 11 m depth (Fig. 1g).

The described area is of particular ecological importance and unique in the Persian Gulf, as soft corals were dominant in the coral community and that they can tolerate such harsh environmental fluctuations, which is likely due to some special mechanisms and changes in their physiology. It is not known whether this structure represents a true framework reef or whether the corals simply follow existing morphology.

References

- Rezai H, Wilson S, Claereboudt M, Riegl B (2004) Coral reef status in ROPME Sea area. In: Wilkinson CR (ed) Status of coral reefs of the world. Australian Institute of Marine Science, Townsville, pp 155–170
- Riegl B (1999) Corals in a non-reef setting in the southern Arabian Gulf (Dubai, UAE): fauna and community structure in response to recurring mass mortality. *Coral Reefs* 18:63–73
- Sheppard CRC, Sheppard ALS (1991) Corals and coral communities of Arabia. *Fauna of Saudi Arabia* 12:3–170
- Sheppard CRC, Price A, Roberts C (1992) Marine ecology of the Arabian region. Academic Press, Cambridge
- Shinn EA (1976) Coral reef recovery in Florida and Persian Gulf. *Environ Geol* 1:241–254

K. Samimi Namin (✉) · H. Rezai · K. Kabiri

Iranian National Center for Oceanography, Etemadzadeh Street, Fatemi Ave, P.O. Box: 14155-4181, Tehran, Iran
e-mail: kaveh_s_n@yahoo.com

Z. Zohari

Faculty of Marine Sciences, Islamic Azad University, Tehran North Branch, Tehran, Iran

Reef sites

Coral Reefs (2009) 28:27
DOI 10.1007/s00338-008-0442-z