Mushroom coral regeneration from a detached stalk

Received: 26 January 2009 / Accepted: 25 March 2009 / Published online: 18 April 2009
© Springer-Verlag 2009

Fig. 1 Stalk regeneration from a free living fungiid coral, *Fungia granulosa*. **a** A 1-year-old polyp cultured in the lab. **b** An aboral view of 'a', notice the stalk "stump" in the center of the polyp. **c** Detached stalk, an aboral view. **d** Detached stalk, an oral view. Notice the appearance of mouth, septae, and tentacles. **e** Fully extended tissue of a regenerated stalk. Note the 'daughter' polyp appears more bleached than the 'parent' due to fewer zooxanthellae (shown in **a**) (scale = 2.5 mm)

Following settlement, attachment, and metamorphosis of the planula larva, fungiid corals develop by forming an elongated calcareous stalk (covered by tissue) on which the polyp "cap" grows. This "cap" then grows radially producing a polyp with a mushroom-like appearance (Hoeksema 1989). Radial growth of the polyp "cap" continues until it detaches from the stalk and becomes a free living solitary polyp (Boschma 1922). Previous studies show that this detachment occurs through the active dissolution of the skeleton across a distinct plane by the calicoblastic cells of the polyp (Yamashiro and Yamazato 1987, 1996). Following detachment in the field, the calcareous stalk remains attached to the substrate, and regenerates one or more polyp "caps". This process can repeat itself one or more times (Hoeksema 1989).

We present evidence of a spontaneous release from the substrate of the whole polyp (stalk and cap) of a *Fungia granulosa* coral cultured in the laboratory. Following this release (Fig. 1a, b), the "cap" detaches from its stalk (Fig. 1c). A full regeneration of the "disc-shaped" free stalk into another free living mushroom-shaped polyp occurs (Fig. 1d, e). Due to its short height (approximately 1 mm), the regenerated stalk "stump" looks like a normal free living young polyp (Fig. 1e). This phenomenon is especially useful for studying the spontaneous decalcification mecha-

nisms that occur during the detachment process as a model for mechanisms that occur during polyp and colony growth, and processes of asexual reproduction, such as polyp expulsion (Kramarsky-Winter et al. 1997).

Acknowledgments The authors would like to thank Varda Wexler (Tel Aviv University) for her graphic assistance and Amikam Shoob (Tel Aviv University) for the photography. This research was supported by the Israel Science Foundation (ISF) to YL.

References

- Boschma H (1922) On budding and coalescence of buds in *Fungia fungites* and *Fungia actiniformis*. Proceedings Koninklijke Nederlandse Academie van Wetenschappen, Amsterdam 24: 257–268
- Hoeksema BW (1989) Taxonomy, phylogeny and biogeography of mushroom corals (Scleractinia: Fungiidae). Zoologische Verhandelingen (Leiden) 254:1–295
- Kramarsky-Winter E, Fine M, Loya Y (1997) Coral polyp expulsion. Nature 387:137
- Yamashiro H, Yamazato K (1987) Studies on the detachment of the discs of the mushroom coral *Fungia fungites* with special reference to hard structural changes. Galaxea 6:283–294
- Yamashiro H, Yamazato K (1996) Morphological studies of the soft tissues involved in skeletal dissolution in the coral *Fungia fungites*. Coral Reefs 15:177–180

M. Vizel (✉) · E. Kramarsky-Winter · Y. Loya
Department of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel
e-mail: vizelmay@post.tau.ac.il

Reef sites

Coral Reefs (2009) 28:939
DOI 10.1007/s00338-009-0496-6