Ying Cao · Yutaka Fujito · Kiyoji Matsuyama
Mamoru Aoki

Effects of electrical stimulation of the medullary raphe nuclei on respiratory movement in rats

Published online: 28 February 2006
© Springer-Verlag 2006

J Comp Physiol A (2006) DOI 10.1007/s00359-005-0087-0

In the **Materials and methods** section—Stimulation and recording—of the original publication, an incorrect value was given in the following sentence:

“Train pulse stimuli (100 Hz, 100–200 pulses, 20 μ s pulse width, 10–30 μ A) were applied four to six times at intervals of 10–15 s in the raphe nuclei and nearby regions.”

The correct sentence must read:

“Train pulse stimuli (100 Hz, 100–200 pulses, **200** μ s pulse width, 10–30 μ A) were applied four to six times at intervals of 10–15 s in the raphe nuclei and nearby regions.”

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00359-005-0087-0>

Y. Cao · Y. Fujito (✉) · K. Matsuyama · M. Aoki
Department of Physiology, Sapporo Medical University
School of Medicine, South 1, West 17, Chuo-ku,
060-8556, Sapporo, Japan
E-mail: fujito@sapmed.ac.jp
Tel.: +81-11-6112111
Fax: +81-11-6441020

Present address: Y. Cao
Institute of Physiology, Shandong University School of Medicine,
Jinan, P. R. China