

Temperature adaptation of lipids in diapausing *Ostrinia nubilalis*: an experimental study to distinguish environmental versus endogenous controls

Elvira L. Vukašinić¹ · David W. Pond² · Gordana Grubor-Lajšić¹ · M. Roger Worland³ · Danijela Kojić¹ · Jelena Purać¹ · Željko D. Popović¹ · Duško P. Blagojević⁴

Received: 11 October 2016 / Revised: 16 May 2017 / Accepted: 17 May 2017 / Published online: 1 June 2017
© Springer-Verlag Berlin Heidelberg 2017

Abstract Larvae of the European corn borer (*Ostrinia nubilalis* Hubn.) were cold acclimated during different phases of diapause to determine if changes in the fatty acid composition lipids occur as part of a programmed diapause strategy, or as a response to low temperatures during winter. Cold acclimation of fifth instar larvae of *O. nubilalis* during diapause had modest effects further on the readjustments in fatty acid composition of triacylglycerols and phospholipids. Overall, FA unsaturation (UFAs/SFAs ratio) was stable, with the exception of the triacylglycerols fraction after exposure to -3 and -10 °C in mid-diapause (MD) when it significantly increased. Differential scanning calorimetry (DSC) was used to examine phase transitions of total body lipid of cold-acclimated larvae in diapause. Thermal analysis indicated that changes in the melt transition temperatures of whole body total lipids were subtle, but consistent with the modest changes in the level of FA unsaturation observed. We conclude that lipid rearrangements are a function of the endogenous “diapause

program” rather than a direct effect of low temperatures, which proved to have limited impact on lipid changes in diapausing larvae of *O. nubilalis*.

Keywords Cold acclimation · Fatty acid composition · *Ostrinia nubilalis* · Lipid melt transition temperatures · Phospholipid · Triacylglycerol

Abbreviations

DSC	Differential scanning calorimetry
FA	Fatty acid
SFA	Saturated fatty acid
UFA	Unsaturated fatty acid
MUFA	Monounsaturated fatty acid
PUFA	Polyunsaturated fatty acid
UFAs/SFAs ratio	Ratio of unsaturated/saturated FAs
FAME	Fatty acid methyl ester
GPEtn/GPChol ratio	Ratio of glycerophosphoethanolamine/glycerophosphocholine

Communicated by G. Heldmaier.

Gordana Grubor-Lajšić: deceased

✉ Elvira L. Vukašinić
elvira.vukasiniovic@dbe.uns.ac.rs

- ¹ Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 3, 21000 Novi Sad, Serbia
- ² Scottish Association for Marine Science, Oban, Argyll PA37 1QA, Scotland, UK
- ³ British Antarctic Survey, Cambridge CB3 0ET, UK
- ⁴ Department of Physiology, Institute for Biological Research, University of Belgrade, Bulevar despota Stefana 142, 11000 Belgrade, Serbia

Introduction

Some insects of temperate and colder climate zones survive harsh environmental conditions by overwintering in a state of diapause. In many of these insect species, diapause is closely connected to cold hardiness and enhances cold stress tolerance (Denlinger 1991, 2002; Bale and Hayward 2010). Even though the level of metabolism is suppressed during dormancy, diapause is not a static state but rather a genetically controlled dynamic physiological process with different successive phases (Bennet et al. 1997; Košťál 2006). Many biochemical and physiological changes that occur during the onset of diapause prepare the organism

for forthcoming overwintering challenges. It is important to emphasize that these physiological mechanisms are initiated exclusively as a result of entering diapause and not as a direct response to low temperature. However, these physiological mechanisms may be further enhanced by subsequent cold exposure (Bale and Hayward 2010). Specifically, diapause induces the synthesis and accumulation of cold-protective metabolites (e.g., sugars and sugar alcohols) and associated chaperones, a cessation of the cell cycle and changes in the storage and membrane lipid compositions (Tamariello and Denlinger 1998; Hodková et al. 1999; Košťál and Šimek 2000; Michaud and Denlinger 2006; Tomčala et al. 2006; Rinehart et al. 2007; MacRae 2010; Vukašinić et al. 2013, 2015).

In some poikilotherms, including insects, restructuring of the lipid composition of cellular membranes is closely related to changing ambient temperature (Cossins and Sinensky 1984; Hazel 1989, 1995; Košťál 2010; Purać et al. 2011). The adaptive restructuring of membrane phospholipids involves an increase of the overall FA unsaturation (UFAs/SFAs ratio) in conjunction with the modifications of phospholipid headgroups, as increase of the GPEth/GPchol ratio. Additional changes considering membrane cold adaptation involve an increase of the cholesterol content, the placement of unsaturated fatty acids on the *sn2* instead *sn1* position of the glycerol backbone of the molecule since unsaturated FA at *sn2* position has a greater effect on membrane fluidity than at the *sn1* position and an increase in the proportion of 16-carbon rather than 18-carbon fatty acids in the plasma membrane, to compensate for the rigidifying effect of sub-zero temperature during winter (Sinensky 1974; Cossins 1994; Hazel 1995; Hochachka and Somero 2002; Košťál et al. 2003; Michaud and Denlinger 2006). At low body temperatures, optimal maintenance of the phase and fluidity of lipid deposits and membrane lipids ensures the availability of energy stores during winter for enzymatic breakdown, while maintaining the functionality and/or structural integrity of membranes (Cossins 1994; Marshall et al. 2014).

The European corn borer (*Ostrinia nubilalis*, Hubn.) is a widespread agricultural pest in the temperate regions of Euroasia, North America and North Africa. This moth normally survives harsh winter conditions inside corn stalk debris, remaining on the soil surface after the crop has been harvested. *O. nubilalis* overwinters as a fifth instar diapausing larvae, which is freeze-tolerant with a considerable supercooling capacity between -20.5 and -25 °C (Grubor-Lajšić et al. 1991). Metabolism of diapausing fifth instar larvae is suppressed and diverted towards: (1) the synthesis and accumulation of cold-protective metabolites, e.g., glycerol and trehalose which are the main cryo and anhydroprotectors (Grubor-Lajšić et al. 1991, 1992; Kojić et al. 2010) and closely connected with

regulation of water content during overwintering (Kojić et al. 2010), (2) readjustment of lipid (storage and membrane) FA composition (Vukašinić et al. 2013, 2015), (3) the expression of stress-related genes (Popović et al. 2015) and (4) rearrangements of the antioxidative defense system (Stanić et al. 2004; Jovanović-Galović et al. 2004, 2007; Kojić et al. 2009). Since some of the overwintering cold stress tolerance mechanisms in diapausing larvae of *O. nubilalis* reach their full potential only after exposure to low temperatures (Grubor-Lajšić et al. 1992; Stanić et al. 2004; Kojić et al. 2010), it is considered that exposure to low temperatures and cold-acclimation are essential for the development of cold hardiness during photoperiod-induced diapause.

Previous studies on larvae of *O. nubilalis* have shown that major compositional remodeling of lipids and changes of their biophysical properties were initiated by the diapause program and preceded the occurrence of low temperatures in fields (Vukašinić et al. 2013, 2015). Higher overall FA unsaturation (UFAs/SFAs ratio) in diapause was the result of concomitant increase in content of MUFAs and decrease in contents of PUFAs and SFAs. Such changes in the FA composition significantly contributed to the depression of melt transition temperatures during the time course of diapause, which was recorded using DSC (Vukašinić et al. 2015).

In this study, the FA composition of triacylglycerols and phospholipids of cold-acclimated fifth instar larvae of *O. nubilalis* during different phases of diapause were analyzed to clarify for the first time in this species, whether the changes in the FA compositions of lipids in overwintering larvae occur as a part of the programmed diapause (endogenous cues) and/or as a direct effect of exposure to low temperatures.

Materials and methods

Insect collection

Fifth instar diapausing larvae of *O. nubilalis* were collected from corn stalk residues left on the soil surface of a harvested field in the vicinity of Novi Sad (AP Vojvodina, Serbia) throughout the diapausing period: autumn (October 2010, early diapause, ED), winter (January 2011, mid-diapause, MD) and early spring (March 2011, termination of diapause, TD). This sampling strategy included all phases of diapause and was consistent with earlier studies (Vukašinić et al. 2013, 2015). The range of ambient air temperatures in the weeks prior to larvae collection were: October 10 to 5 °C, January -5 to -0.5 °C, and March -14 to 5.5 °C.

Laboratory cold acclimation

Immediately after collection larvae were stored under controlled conditions (5 °C) for 10 days. A sample was then taken for analysis and the remainder gradually cooled (1 °C day⁻¹) to the first experimental acclimation temperature (−3 °C). At this point, half were held at −3 °C for a further 10 days while the remaining larvae were cooled at the same rate to −10 °C. The latter sample was then held at −10 °C for 10 days. All samples were prepared for analysis after the 10 day acclimation period. This treatment was repeated for each field sample (Fig. 1).

Extraction of total lipids

Three pools, three larvae per pool, were prepared for each experimental group. Total lipids were extracted from whole bodies (three larvae per pool) by homogenizing (T10 Basic Ultra-Turrax Homogenizer, IKA, Germany) them in ice-cold chloroform:methanol (2:1 v/v) according to a modified method of Folch et al. (1957). The homogenates of the whole body pool were filtered through Whatman No.1 filter paper, pre-washed prior to filtration with chloroform. After the addition of 1/4 volume of KCl (0.88% w/v) to the filtrate, samples were thoroughly vortexed and centrifuged for 5 min at 0.2G to achieve phase separation. The upper methanol/water layer was discarded, while the lower chloroform phase containing the total lipid extract was dried under a stream of nitrogen using an N-EVAP system (Organomation). Total lipid extracts from larval whole bodies were resuspended in chloroform (10 mg ml⁻¹ final concentration) and stored at −20 °C until analysis.

Larval whole body triacylglycerol and phospholipid FA composition analysis

Triacylglycerol and phospholipid fractions prepared from total lipids extracted from whole bodies of fifth instar diapausing larvae of *O. nubilalis* were separated by thin-layer chromatography (TLC) using 20×20 cm silica gel plates with hexane:diethyl ether:glacial acetic acid (80:20:2 v/v/v). The analyzed lipid fraction was visualized under an ultraviolet light (254 nm), after spraying plates with 2', 7'-dichlorofluorescein dissolved in methanol (0.1% w/v), and identified by comparison with commercial standards. Triacylglycerol and phospholipid fractions were scraped from the developed plates into reaction vials and trans-methylated with 1% sulfuric acid in methanol in sealed vials at 50 °C for 16 h (Christie 1982). After trans-methylation, milli-Q water and hexane:diethyl ether (1:1 v/v), in the ratio of 1:1 v/v, was added to each vial, which was vortexed and centrifuged at 0.2G for 2 min. From the resultant two-phase mixture, the upper organic phase containing the FAMES was removed and the extraction was repeated by adding hexane:diethyl ether (1:1 v/v). After the re-extraction, the upper organic phase containing the FAMES was removed again. Two milliliter of NaHCO₃ (2% w/v) was added to the combined upper organic phase which was then vortexed and centrifuged at 0.2G for 2 min. The lower layer was discarded while the upper phase containing the FAMES was evaporated under nitrogen. The unpurified FAMES from triacylglycerol and phospholipid fractions were re-dissolved in 50–100 μL of hexane and applied to the base of a 20×20 cm silica gel TLC plate using a Hamilton micro syringe. The TLC plate was developed in a hexane:diethyl ether:glacial acetic acid (90:10:1 v/v/v) solvent system. The FAMES were visualized under an ultraviolet light (254 nm) after spraying plates with 2', 7'-dichlorofluorescein dissolved in methanol (0.1% w/v). Their position was then marked using a pencil and scraped from the plates into reaction vials. Hexane:diethyl ether (1:1 v/v) and NaHCO₃ (2%, w/v), in the ratio of 2:1 v/v, was added to each vial, which was then vortexed and centrifuged at 0.2G for 2 min. The upper layer was then removed to a second clean vial. An additional quantity of hexane:diethyl ether (1:1 v/v) was added to the lower phase in the first vial, which was vortexed and centrifuged at 0.2G for 2 min. The collected upper layer containing the purified FAMES was finally evaporated under a stream of nitrogen. The purified FAMES from triacylglycerol and phospholipid fractions were re-dissolved and analyzed using TRACE 2000, Thermo Electron, Gas Chromatograph (Thermo Scientific, UK) equipped with a Restek Stabilwax column (0.32 mm i.d. x 30 m). Hydrogen was used as the carrier gas (Pond et al. 2008).




Fig. 1 Schematic illustration of experiment (explanation given in the text). ED early diapause, MD mid-diapause, TD termination of diapause

Differential scanning calorimetry analysis

A differential scanning calorimeter (DSC 820, Mettler Toledo) was used to examine thermal changes in total lipids. Aliquots of dissolved total lipids extracted from whole bodies of fifth instar larvae of *O. nubilalis* were added to a 40 μl aluminium DSC sample pan and dried under a stream of nitrogen until a dry sample of about 1.5 mg had been accumulated. A lid was then sealed onto the pan and the sample analysed in the DSC using the following temperature program. Samples were initially cooled to -60°C at 4°C min^{-1} , held isothermally for 1 min at this temperature then warmed to 40°C at 4°C min^{-1} . The thermograms were evaluated using STARe software (version 6).

Analysis was only conducted using whole body total lipids (TLs) and not the triacylglycerols fraction (TAGs) as in a previous study we determined that they resulted in very similar thermograms. We determined that melt changes were predominately attributable to TAGs with other lipid classes contributing little to the melt of the complex mixture of total lipids (Vukašinovic et al. 2015).

For each endothermic event (melt) the peak, onset and end set during the melt were measured. The peak of the melt endotherm was taken as the temperature at which the difference between the DSC curve and the interpolated baseline is at a maximum; the onset temperature, the temperature at which melting starts and the end set, the temperature at which melting finishes. The area under the melt endotherm represents the amount of heat energy [mJ] consumed during phase change, melting.

Statistical analysis

All results are expressed as mean \pm standard error (SE). The peak areas were converted to percentages of total content of FAs within the same chromatogram (relative proportions) for each FA detected and the logarithmic transformations of these values were analyzed using one-way analyses of variance (one-way ANOVA) followed by Tukey's HSD post hoc test with a level of significance at least $p < 0.05$ between groups (Hinkle et al. 1994) using STATISTICA software, version 12 (Stat Soft Inc., Tulsa, OK, USA).

To summarize data, the overall change in FA unsaturation in the triacylglycerols and phospholipids, the ratio of unsaturated to saturated fatty acids (UFAs/SFAs ratio) was calculated as the ratio of the total proportion of all unsaturated fatty acids [UFAs = (MUFAs + PUFAs)] over the total proportion of all saturated fatty acids (SFAs).

Results

FA composition of triacylglycerol and phospholipid fractions of cold-acclimated diapausing larvae of *O. nubilalis*

First of all, the overall FA unsaturation in triacylglycerols and phospholipids of *O. nubilalis* during diapause (ED, MD, TD), regarding larvae stored under controlled conditions (5°C), were determined (Fig. 2). The overall FA unsaturation (UFAs/SFAs ratio) did not change during time course of diapause until the termination period (TD), when it decreased (Fig. 2b, d), significantly in phospholipids (Fig. 2d), primarily as a consequence of a significant decrease in the relative proportion of MUFAs and a concomitant increase in the proportion of SFAs and PUFAs (Fig. 2c).

Seven FAs with 16 and 18 carbons were determined in triacylglycerol and phospholipid fractions: palmitic (16:0), palmitoleic (16:1n-7), stearic (18:0), oleic (18:1n-9), vaccenic (18:1n-7), linoleic (18:2n-6) and α -linolenic acid (18:3n-3) (Figs. 3a, 4a). In triacylglycerol, palmitic (16:0), palmitoleic (16:1n-7) and oleic acid (18:1n-9) dominated and comprise more than 90% of the total FA content (Fig. 3a). FAs in phospholipids were more evenly represented with in general a higher abundance of PUFAs, especially linoleic acid (18:2n-6) (Fig. 4a).

In the case of triacylglycerols, the major structural class of FAs was MUFAs (Fig. 3b), with palmitoleic (16:1n-7) and oleic acid (18:1n-9) being the dominant FAs (Fig. 3a). The proportion of MUFAs did not change significantly in response to cold treatment during the time course of diapause (Fig. 3b). The relative abundance of SFAs changed only in mid-diapause (MD), when significantly decreased after exposure to -3 and -10°C (Fig. 3b), as a result of a significant decrease in the relative proportion of palmitic acid (16:0) (Fig. 3a). The least abundant class of triacylglycerol FAs, PUFAs, changed substantially only during early diapause (ED) after the cold acclimation to -3°C , as a result of a significant increase in the proportion of linoleic acid (18:2n-6) (Fig. 3a). To summarize, cold acclimation did not induce significant changes in the overall FA unsaturation (UFAs/SFAs ratio) of storage lipids during the time course of diapause, except in the mid-diapause (MD) after acclimation to -3 and -10°C , which induced a significant increase in the UFAs/SFAs ratio (Fig. 3c).

The FA composition of phospholipids did not change significantly in response to cold acclimation during any phase of diapause (Fig. 4b, c), even though the temperature of exposure during different phases of diapause induced significant changes in the relative proportions of selected FAs (Fig. 4a). Namely, in early diapause (ED), cold acclimation to -3°C induced a significant




Fig. 2 The fatty acid composition of triacylglycerols and phospholipids of cold-acclimated diapausing larvae of *O. nubilalis* to 5°C during early diapause (ED), mid-diapause (MD) and termination of diapause (TD) presenting the results considering major structural classes of FAs: saturates (SFAs), unsaturates (UFAs), monounsaturates (MUFAs) and polyunsaturates (PUFAs) (a, c), and changes in

overall FA unsaturation (UFAs/SFAs ratio) (b, d). Bars represent mean ± standard error, n=3 groups, three larvae per group. Asterisk above the bars indicate the level of statistically significant differences in the relative proportion of structural classes determined by Tukey’s HSD post hoc test in control group (5°C) during the time course of diapause (ED, MD, TD): *p<0.05

increase in the relative proportion of vaccenic (18:1n-7) and linoleic acid (18:2n-6), while acclimation to -10°C a decrease in the relative proportion of α-linolenic acid (18:3n-3) (Fig. 4a). During mid-diapause (MD) cold acclimation to -10°C induced a significant increase in the relative proportion of vaccenic (18:1n-7) and linoleic acid (18:2n-6) and a significant decrease in the relative proportion of stearic acid (18:0) (Fig. 4a). Cold acclimation during the termination of diapause (TD) did not induce any significant changes in the relative proportion of FA composition of phospholipids (Fig. 4a). Considering the overall FA unsaturation (UFAs/SFAs ratio), cold acclimation during different phases of diapause did not induce substantial changes in the FA composition of phospholipids (Fig. 4c). The UFAs/SFAs ratio showed a similar trend in both triacylglycerols and phospholipids (Figs. 3c, 4c), but with a lower overall FA unsaturation in phospholipids (Fig. 4c), which is the consequence of a more even distribution of FAs in this fraction (Fig. 4a).

Thermal changes in whole body total lipids of cold-acclimated diapausing larvae of *O. nubilalis*

Differential scanning calorimetry (DSC) was used to examine the solid–liquid phase transition properties of total lipids extracted from the whole bodies of cold-acclimated diapausing larvae of *O. nubilalis* focusing on the lipid melt transition temperatures (Fig. 5; Table 1).

Thermal analysis of total lipids detected two clearly defined melt endotherms: endotherm 1 and endotherm 2 (Fig. 5). The melt peaks were similar for cold-acclimated samples during different diapause phases; the peak of endotherm 1 varied only within the range -18.09 to -19.61°C while the peak of endotherm 2 within the range 1.64–6.54°C (Fig. 5; Table 1).




Fig. 3 The fatty acid composition of triacylglycerols of cold acclimated diapausing larvae of *O. nubilalis* to 5, -3 and -10 °C (a), during early diapause (ED), mid-diapause (MD) and termination of diapause (TD) presenting the results considering major structural classes of FAs: saturates (SFAs), unsaturates (UFAs), monounsaturates (MUFAs) and polyunsaturates (PUFAs) (b), and changes in

overall FA unsaturation (UFAs/SFAs ratio) in triacylglycerols (c). Bars represent mean \pm standard error, $n=3$ groups, three larvae per group. Asterisk above the bars indicate the level of statistically significant differences determined by Tukey's HSD post hoc test compared to control group (5 °C) during the time course of diapause (ED, MD, TD): * $p < 0.05$

Discussion

Lipid rearrangements occur in both non-dormant and dormant states in insects after exposure to low temperatures. However, in those insects in which a dormant state such as diapause is closely tied to cold hardiness, changes in the membrane lipid composition are not primarily a result of response to low temperatures, but are initiated with a photoperiodically induced diapause program and later intensified in response to cold (Kostal and Simek 1998; Hodkova et al. 1999; Michaud and Denlinger 2006; Bale and Hayward 2010; Rozsypal et al. 2014). Thus, in cold-tolerant insects it is important to clarify whether lipid rearrangements are exclusively governed by a diapause program, by cold exposure, or a combination of both.

In our previous studies on field collected larvae of *O. nubilalis*, we demonstrated that the compositional

remodeling of lipids is accompanied by changes in their biophysical properties and is a function of photoperiodically induced diapause program, rather than a result of exposure to low temperatures of diapausing larvae. Specifically, major changes in FA composition occur early in diapause, well ahead of occurrence of low temperatures during winter. A significant increase in the overall FA unsaturation (UFAs/SFAs ratio) resulted from an increase in the relative proportion of MUFAs, primarily palmitoleic (C16:1n-7) and oleic acids (C18:1n-9), and a concomitant decrease in PUFAs and SFAs content, α -linoleic (C18:2n-6) and palmitic acid (C16:0), respectively. Thermal analyses (DSC) of whole body total lipids have shown significantly lower melt transition temperatures during the time course of diapause, in comparison to non-diapause state (Vukašinović et al. 2015).




Fig. 4 The fatty acid composition of phospholipids of cold-acclimated diapausing larvae of *O. nubilalis* to 5, –3 and –10 °C (a), during early diapause (ED), mid-diapause (MD) and termination of diapause (TD) presenting the results considering major structural classes of FAs: saturates (SFAs), unsaturates (UFAs), monounsaturates (MUFAs) and polyunsaturates (PUFAs) (b), and changes in

overall FA unsaturation (UFAs/SFAs ratio) in phospholipids (c). Bars represent mean ± standard error, *n* = 3 groups, three larvae per group. Asterisk above the bars indicate the level of statistically significant differences determined by Tukey’s HSD post hoc test compared to control group (5 °C) during the time course of diapause (ED, MD, TD): **p* < 0.05

The aim of this study was to evaluate whether larvae of *O. nubilalis* are capable of responding to changes in environmental temperature during diapause or if the ability to acclimate is determined endogenously in diapause well ahead the occurrence of low winter temperature. Since *O. nubilalis* is a freeze-tolerant insect species with a high supercooling capacity in which diapause and cold hardiness are closely related, two different acclimation temperatures were chosen: –3 °C, as a potentially critical temperature in initiating adaptation to low temperatures and –10 °C as a typical low mid winter temperature.

Results compared to changes in the control group (5 °C) showed that trend in the UFAs/SFAs ratio during diapause is similar to the results published in our previous studies, where the increase in the overall unsaturation (UFAs/SFAs ratio) occurs during transition to diapause and remains stable until the termination period, when it significantly

decreases (Vukašinović et al. 2015). In the current study, the UFAs/SFAs ratio was high until the termination period (TD) when it decreased, significantly in phospholipids, mainly as a consequence of a decrease in dominant MUFAs and a concomitant increase in SFAs and PUFAs. In addition, the results of this study showed as well that cold acclimation of fifth instar *O. nubilalis* larvae during the state of diapause had modest effects further on the readjustments in FA composition of triacylglycerols and phospholipids. In this case, the overall FA unsaturation (UFAs/SFAs ratio) was without significant change, except for the triacylglycerol fraction after the exposure to –3 and –10 °C in mid-diapause (MD), where UFAs/SFAs ratio significantly increased as a result of a decrease in the relative proportion of SFAs, concretely palmitic acid (C16:0). Also, thermal analysis (DSC) of whole body total lipids revealed that cold acclimation of larvae during different phases of diapause




Fig. 5 DSC thermograms of total lipids of cold-acclimated diapausing larvae of *O. nubilalis* to 5, –3 and –10 °C, during different phases of diapause; *ED* early diapause, *MD* mid-diapause and *TD* termination of diapause. Melt endotherm 1 and 2 indicate solid–liquid phase transitions. The peak of melt endotherm 1 and the peak of melt endo-

therm 2 presented in Table 1, indicates the temperature at which the difference between the DSC curve and the interpolated baseline is at a maximum. Since the total lipids comprise complex mixtures of molecules with different level of unsaturation, different chain length, melting is a multistep process

Table 1 Melt transition temperatures (peak, onset and end set) of total lipids extracted from the whole bodies of cold acclimated diapausing larvae of *O. nubilalis* to 5, –3 and –10 °C, during different phases of diapause; *ED* early diapause, *MD* mid-diapause, *TD* termination of diapause

	Peak of melt (°C)		Onset (°C)		End set (°C)	
	1	2	1	2	1	2
ED (°C)						
5	-18.92	3.91	-26.19	-9.49	-13.82	6.42
-3	-19.33	3.19	-26.02	-10.08	-13.11	5.97
-10	-19.61	1.64	-25.35	-9.77	-15.65	4.35
MD (°C)						
5	-18.65	4.70	-26.93	-8.91	-11.83	6.74
-3	-18.92	3.63	-24.29	-9.16	-14.78	5.92
-10	-19.01	2.96	-25.01	-9.13	-13.16	5.72
TD (°C)						
5	-18.26	5.98	-25.69	-7.07	-12.26	7.55
-3	-19.08	4.83	-26.21	-9.05	-11.84	7.22
-10	-17.46	5.21	-25.16	-11.68	-9.2	8.14

The temperature at peak of melt indicates the temperature at which the difference between the DSC curve and the interpolated baseline is at maximum (Fig. 5). The melt endotherm (indicate phase transitions from solid to liquid) is determined as well by the onset temperature (temperature at which melting starts) and the end set temperature (temperature at which melting finishes)

slightly reduce the melt transition temperatures, which according to our previous findings significantly decreased during the transition to diapause, compared to the non-diapause state (Vukašinović et al. 2015). The essential compositional adjustments of lipids (FAs rearrangements) are accompanied by changes in the biophysical properties of total lipids and is a reflection of the diapause program in larvae of *O. nubilalis*. This provides the ability to survive low winter temperatures without additional adaptations to increase fluidity of lipids during overwintering.

Lipids play a key role in protection against cold stress and are involved in the initial preparations for diapause which occur prior to the onset of winter conditions. Lipids also critically influence both the progression of diapause and its stress tolerance phenotype as represent the front line protection against cold stress, representing the interface between the insect cells and their environment (Michaud and Denlinger 2006; Bale and Hayward 2010). It has been previously well documented that in some insect species winter diapause induces changes in the FA composition of triacylglycerol and phospholipid leading to a higher overall FA unsaturation (UFAs/SFAs ratio) (Bennet et al. 1997; Kostal and Simek 1998; Hodkova et al. 1999; Michaud and Denlinger 2006; Rozsypal et al. 2014). The results of this study are in accordance with these findings, i.e., cold acclimation resulted in only modest further changes in FA composition in *O. nubilalis* having changed significantly during transition to diapause (Vukašinović et al. 2015).

Since *O. nubilalis* is an atypical freeze-tolerant insect species which depresses its' supercooling point (SCP) well-below the SCP of other freeze-tolerant species, it is highly likely that changes in the composition of lipids initiated early in diapause. These changes are mostly associated with MUFAs in both membrane and storage lipids and enhance thermal tolerance of this species in environments with wide fluctuations in winter temperatures. According to Michaud and Denlinger (2006), a dominance of MUFAs in membranes is energetically more favourable than synthesis of PUFAs since it allows a wide window of fluidity without a need for continuous readjustment of FAs in response to or in preparation for low temperatures. In membrane lipids, the fluidity is maintained constitutively at a high level, probably to avoid the unregulated transition to the gel phase during cold spells. Based on the results of this study and considering our previous studies on lipids in *O. nubilalis* (Vukašinović et al. 2013, 2015), we conclude that lipid rearrangements are a function of the endogenous "diapause program" rather than a direct effect of low temperatures, which proved to have limited impact on lipid changes in diapausing larvae of *O. nubilalis*.

Acknowledgements This work was funded by the Ministry of Education, Science and Technological Development of the Republic of

Serbia, grant no. 173014., project entitled "Molecular mechanisms of redox signalling in homeostasis: adaptation and pathology". This publication is a contribution to the ECOSYSTEMS Programme of the British Antarctic Survey, UK.

References

- Bale JS, Hayward SA (2010) Insect overwintering in a changing climate. *J Exp Biol* 213:980–994
- Bennett VA, Pruitt NL, Lee RE (1997) Seasonal changes in fatty acid composition associated with cold-hardening in third instar larvae of *Eurosta solidaginis*. *J Comp Physiol B* 167:249–255
- Christie WW (1982) Lipid analysis, 2nd edn. Pergamon Press, Oxford
- Cossins AR (1994) Homeoviscous adaptation and its functional significance. In: Cossins AR (ed) Temperature adaptation of biological membranes. Portland Press, London, pp 63–75
- Cossins AR, Sinensky M (1984) Adaptations of membranes to temperature, pressure and exogenous lipids. In: Shinitzky M (ed) Physiology of membrane fluidity. CRC Press, Boca Raton, pp 1–20
- Denlinger DL (1991) Relationship between cold-hardiness and diapause. In: Lee RE, Denlinger DL (eds) Insects at low temperature. Chapman and Hall, New York, pp 174–198
- Denlinger DL (2002) Regulation of diapause. *Annu Rev Entomol* 47:93–122
- Folch J, Lees M, Sloane Stanley GH (1957) A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 226:497–509
- Grubor-Lajšić G, Block W, Palanacki V, Glumac S (1991) Cold hardiness parameters of overwintering diapause larvae of *Ostrinia nubilalis* in Vojvodina, Yugoslavia. *Cryo-Letters* 12:177–182
- Grubor-Lajšić G, Block W, Worland R (1992) Comparison of the cold hardiness of two larval Lepidoptera (Noctuidae). *Physiol Entomol* 17:148–152
- Hazel JR (1989) Cold adaptation in ectotherms: regulation of membrane function and cellular metabolism. *advance in comparative and environmental. Physiology* 4:1–50
- Hazel JR (1995) Thermal adaptation in biological membranes: is homeoviscous adaptation the explanation? *Annu Rev Physiol* 57:19–42
- Hinkle ED, Wiersma W, Jurs GS (1994) Applied statistics for behavioral sciences, 3rd edn. Houghton Mifflin Company, Boston
- Hochachka PW, Somer GN (2002) Biochemical adaptation: mechanism and process in physiological evolution. Oxford University Press, Oxford, p 560
- Hodková M, Šimek P, Zahradníčková H, Nováková O (1999) Seasonal changes in the phospholipid composition in thoracic muscles of a heteropteran, *Pyrrhocoris apterus*. *Insect Biochem Mol Biol* 29:367–376
- Jovanovic-Galovic A, Blagojevic DP, Grubor-Lajšić G, Worland MR, Spasic MB (2007) Antioxidant defense in mitochondria during diapause and postdiapause development of European corn borer (*Ostrinia nubilalis*, Hbn.). *Arch Insect Biochem Physiol* 64:111–119
- Jovanović-Galović A, Blagojević DP, Grubor-Lajšić G, Worland R, Spasić MB (2004) Role of antioxidant defense during different stages of preadult life cycle in European corn borer (*Ostrinia nubilalis*, Hubn.): diapause and metamorphosis. *Arch Insect Biochem Physiol* 55:79–89
- Kojic D, Spasojevic I, Mojovic M, Blagojevic D, Worland MR, Grubor-Lajšić G, Spasic MB (2009) Potential role of hydrogen peroxide and melanin in the cold hardiness of *Ostrinia nubilalis* (Lepidoptera: Pyralidae). *Eur J Entomol* 106:451–454

- Kojić D, Purać J, Popović ŽD, Pamer E, Grubor-Lajšić G (2010) Importance of the body water management for winter cold survival of the European corn borer *Ostrinia nubilalis* Hübn. (Lepidoptera: Pyralidae). *Biotechnol Biotechnol Equip* 24:648–654
- Kostal V, Šimek P (1998) Changes in fatty acid composition of phospholipids and triacylglycerides after cold acclimation of an aestivating insect prepupa. *J Comp Physiol B* 168:453–460
- Košťál V (2006) Eco-physiological phases of insect diapause. *J Insect Physiol* 52:113–127
- Košťál V (2010) Cell structural modifications in insects at low temperatures. In: Denlinger DL, Lee RE Jr (eds) *Low temperature biology of insects*. Cambridge University Press, Cambridge
- Košťál V, Berková P, Šimek P (2003) Remodelling of membrane phospholipids during transition to diapause and cold-acclimation in the larvae of *Chymomyza costata* (Drosophilidae). *Comp Biochem Physiol B Biochem Mol Biol* 135(3):407–419
- Košťál V, Šimek P (2000) Overwintering strategy in *Pyrrhocoris apterus* (Heteroptera): the relations between life-cycle, chill tolerance and physiological adjustments. *J Insect Physiol* 46:1321–1329
- Lee RE (1991) Principles of insect low temperature tolerance. In: Lee RE, Denlinger DL (eds) *Insects at low temperature*, Chapman, New York, pp 17–47
- MacRae TH (2010) Gene expression, metabolic regulation and stress tolerance during diapause. *Cell Mol Life Sci* 67:2405–2424
- Marshall KE, Thomas RH, Roxin Á, Chen EKY, Brown JCL, Gillies ER, Sinclair BJ (2014) Seasonal accumulation of acetylated triacylglycerols by a freeze-tolerant insect. *J Exp Biol* 217:1580–1587
- Michaud MR, Denlinger DL (2006) Oleic acid is elevated in cell membrane during rapid cold-hardening and pupal diapause in the flesh fly, *Sarcophaga crassipalpis*. *J Insect Physiol* 52:1073–1082
- Pond DW, Fallick EA, Stevens JC, Morrison JD, Dixon RD (2008) Vertebrate nutrition in a deep-sea hydrothermal vent ecosystem: fatty acid and stable isotope evidence. *Deep Sea Res Part I Oceanogr Res Pap* 55:1718–1726
- Popović ŽD, Subotić A, Nikolić TV, Radojičić R, Blagojević DP, Grubor-Lajšić G, Košťál V (2015) Expression of stress-related genes in diapause of European corn borer (*Ostrinia nubilalis* Hübn.). *Comp Biochem Physiol Part B* 186:1–7
- Purać J, Pond WD, Grubor-Lajšić G, Kojić D, Blagojević PD, Worland RM, Clark SM (2011) Cold hardening induces transfer of fatty acids between polar and nonpolar lipid pools in the Arctic collembollan *Megaphorura arctica*. *Physiol Entomol* 36:135–140
- Rinehart JP, Li A, Yocum GD, Robich RM, Hayward SAL, Denlinger DL (2007) Up-regulation of heat shock proteins is essential for cold survival during insect diapause. *Proc Natl Acad Sci USA* 104:11130–11137
- Rozsypal J, Košťál V, Berková P, Zahradníčková H, Šimek P (2014) Seasonal changes in the composition of storage and membrane lipids in overwintering larvae of the codling moth, *Cydia pomonella*. *J Therm Biol* 45:124–133
- Sinensky M (1974) Homeoviscous adaptation—a homeostatic process that regulates the viscosity of membrane lipids in *Escherichia coli*. *Proc Natl Acad Sci USA* 71:522–525
- Stanic B, Jovanovic-Galovic A, Blagojevic DP, Grubor-Lajšić G, Worland R, Spasic MB (2004) Cold hardiness in *Ostrinia nubilalis* (Lepidoptera: Pyralidae): glycerol content, hexose monophosphate shunt activity, and antioxidative defense system. *Eur J Entomol* 101:459–466
- Storey KB, Storey JM (1991) Biochemistry of cryoprotectants. In: Lee RE Jr, Denlinger DL (eds) *Insects at low temperature*. Chapman and Hall, New York, pp 64–93
- Tammariello SP, Denlinger DL (1998) G0/G1 cell cycle arrest in the brain of *Sarcophaga crassipalpis* during pupal diapause and the expression pattern of the cell cycle regulator, proliferating cell nuclear antigen. *Insect Biochem Molec Biol* 28:83–89
- Tomčala A, Tollarová M, Overgaard J, Šimek P, Košťál V (2006) Seasonal acquisition of chill tolerance and restructuring of membrane glycerophospholipids in an overwintering insect: triggering by low temperature, desiccation and diapause progression. *J Exp Biol* 209:4102–4114
- Tsvetkova NM, Quinn PJ (1994) Compatible solutes modulate membrane lipid phase behavior. In: Cossins AR (ed) *Temperature adaptation of biological membranes*, Portland Press, London, pp 49–62
- Vukašinović E, Pond WD, Worland MR, Kojić D, Purać J, Blagojević PD, Grubor-Lajšić G (2013) Diapause induces changes in the composition and biophysical properties of lipids in larvae of the European corn borer, *Ostrinia nubilalis* (Lepidoptera: Crambidae). *Comp Biochem Physiol Part B* 165:219–225
- Vukašinović E, Pond WD, Worland MR, Kojić D, Purać J, Popović DŽ, Grubor-Lajšić G (2015) Diapause induces remodeling of the fatty acid composition of membrane and storage lipids in overwintering larvae of *Ostrinia nubilalis*, Hübn. (Lepidoptera: Crambidae). *Comp Biochem Physiol Part B* 184:36–43