Erratum to: Global and regional ocean carbon uptake and climate change: sensitivity to a substantial mitigation scenario

Marcello Vichi · Elisa Manzini · Pier Giuseppe Fogli ·
Andrea Alessandri · Lavinia Patara · Enrico Scoccimarro ·
Simona Masina · Antonio Navarra

Published online: 24 July 2011
© Springer-Verlag 2011

Erratum to: Clim Dyn
DOI 10.1007/s00382-011-1079-0

In the original publication of this article, Fig. 8 was published without the vertical bars indicated in the caption. The correct figure is produced here.

The online version of the original article can be found under doi:[10.1007/s00382-011-1079-0](https://doi.org/10.1007/s00382-011-1079-0).

M. Vichi (✉) · E. Manzini · P. G. Fogli · A. Alessandri ·
L. Patara · S. Masina · A. Navarra
Centro Euro-Mediterraneo per i Cambiamenti Climatici
(CMCC), Viale Aldo Moro 44, 40127 Bologna, Italy
e-mail: marcello.vichi@cmcc.it

M. Vichi · E. Manzini · E. Scoccimarro · S. Masina ·
A. Navarra
Istituto Nazionale di Geofisica e Vulcanologia,
Bologna, Italy

Present Address:
E. Manzini
Max Planck Institute for Meteorology, Hamburg, Germany

L. Patara
Leibniz Institute of Marine Sciences (IFM-GEOMAR),
Kiel, Germany

Present Address:
A. Alessandri
ENEA, Rome, Italy

Fig. 8 Ocean carbon uptake in the 20C, A1B and E1 simulations divided into major basin contributions (11 years running means, negative values mean ocean uptake). The Southern Ocean (SO) is defined as the area southern of 50°S. The range of estimates for contemporary net CO₂ fluxes from Gruber et al. (2009) are shown as vertical bars with the relative colour for each basin