Penoplasty for buried penis in infants and children: report of 100 cases

Lena Perger · Robert S. Hanley · Neil R. Feins

Published online: 13 December 2008
© Springer-Verlag 2008

Erratum to: *Pediatr Surg Int* DOI 10.1007/s00383-008-2283-9

Due to errors in typesetting, the images in Figs. 2, 3, 4, 5, 6, 7, and 8 were all in the wrong positions, so that the illustrations did not match the legends. The correctly matched figures and legends are presented here.

Fig. 2 Buried penis in a 6-month-old boy

The online version of the original article can be found under doi:[10.1007/s00383-008-2283-9](https://doi.org/10.1007/s00383-008-2283-9).

L. Perger
Department of Surgery, University of New Mexico Health
Sciences Center, Albuquerque, NM, USA
e-mail: lperger@gmail.com

R. S. Hanley
Department of Urology, Lahey Clinic,
Tufts University School of Medicine,
Burlington, MA, USA

N. R. Feins (✉)
Department of Surgery, Children's Hospital Boston,
Harvard Medical School, Fegan 3, 300 Longwood Avenue,
Boston, MA 02115, USA
e-mail: neil.feins@childrens.harvard.edu

Fig. 3 Normal phallus exposed by application of pressure at the base of penile shaft

Fig. 5 Suturing of dorsal penile shaft base to prepubic fascia

Fig. 4 Circumferential degloving of penis and development of preputial flap

Fig. 6 Attachment of shaft skin to base of penile shaft with fullthickness vertical mattress sutures

Fig. 7 Covering of penile shaft with skin, closure of mucosal collar

Fig. 8 Improved exposure of penis 2 weeks after surgery