Erratum to: Role of endothelial Nox2 NADPH oxidase in angiotensin II-induced hypertension and vasomotor dysfunction

Colin E. Murdoch · Sara P. Alom-Ruiz ·
Minshu Wang · Min Zhang · Simon Walker ·
Bin Yu · Alison Brewer · Ajay M. Shah

Published online: 9 April 2014
© Springer-Verlag Berlin Heidelberg 2014

Erratum to: Basic Res Cardiol (2011) 106:527–538
DOI 10.1007/s00395-011-0179-7

The data symbols for Wt and Nox2Tg were mistakenly reversed in Fig. 5b. The correct figure is given below.

Fig. 5 Effect of chronic angiotensin II treatment on blood pressure in Tg and WT mice. Ambulatory blood pressure was measured in WT and Tg mice before and after implantation of osmotic pumps delivering either **a** saline, **b** low dose AngII (0.3 mg/kg/day) or **c** high dose AngII (1.1 mg/kg/day). Systolic, diastolic and mean blood pressure and heart rate were averaged over 24 h periods. $n = 6$, $*P < 0.05$ for a significant interaction by two-way ANOVA

The online version of the original article can be found under doi:10.1007/s00395-011-0179-7.

C. E. Murdoch · S. P. Alom-Ruiz · M. Wang · M. Zhang ·
S. Walker · B. Yu · A. Brewer · A. M. Shah
Cardiovascular Division, King's College London British Heart
Foundation Centre, London SE5 9PJ, UK

A. M. Shah (✉)
Cardiovascular Division, The James Black Centre,
King's College London, 125 Coldharbour Lane,
London SE5 9NU, UK
e-mail: ajay.shah@kcl.ac.uk