Tune B. Bonn
Karin Ldtke
Rainer Jordan
Petr tpnek
Christine M. Papadakis

Aggregation behavior of amphiphilic poly(2-alkyl-2-oxazoline) diblock copolymers in aqueous solution studied by fluorescence correlation spectroscopy

Published online: 15 September 2004
 Springer-Verlag 2004

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00396-004-1131-2>

T. B. Bonn · C. M. Papadakis ()
Fakultt fr Physik und
Geowissenschaften, Universitt Leipzig,
Linnstr. 5, 04103 Leipzig, Germany
E-mail: Christine.Papadakis@ph.tum.de

K. Ldtke · R. Jordan
Chemie Department,
Technische Universitt Mnchen,
Lichtenbergstr. 4, 85747 Garching,
Germany

P. tpnek
Institute of Macromolecular Chemistry,
Academy of Sciences of the Czech
Republic, Heyrovsky Sq. 2,
CZ-16206 Prague, Czech Republic

Present address: T. B. Bonn
C. M. Papadakis
Physik Department E13,
Technische Universitt Mnchen,
James-Franck-Str. 1, 85747 Garching,
Germany

R. Jordan
Department of Chemistry,
Chemical Engineering and Materials
Science, Polytechnic University,
Six Metrotech Center, Brooklyn,
NY 11201, USA

**Colloid Polym Sci (2004)
282: 833–843**

Due to an editorial error, the following changes should be made:

In the original paper on p. 841, left column, row 21, ‘the mutual diffusion micellar coefficient’ should be replaced by ‘the micellar mutual diffusion coefficient’.

On the same page, left column, row 3 from below, ‘ $r_H = (11.9 \pm 0.7) \text{ nm}$ ’ should be replaced by ‘ $r_H = (13 \pm 2) \text{ nm}$ ’.

In Table 1, row 4, column 8, the value of the PDI of sample PMOX₂₆ of 2.21 should be replaced by ‘1.21’. The corrected Table reads:

Table 1 Analytical values of the polymers

Polymer	DP _n [M1] ₀ /[I] ₀ 1st block ^a	DP _n [M2] ₀ /[I] ₀ 2nd block ^a	DP 1st block ^b	DP 2nd block ^b	M _n ¹ H-NMR ^c	Yield/% ^d	PDI ^e
P[(MOx) ₄₀ (NOx) ₇]	30	6	40	7	4034	89	1.20
P[(NOx) ₁₀ (MOx) ₃₂]	7	40	10	32	4796	96	1.07
PMOX ₂₆	30	–	26	–	2313	87	1.21

^aDegree of polymerization (DP_n) calculated from initial monomer/initiator feed

^bDegree of polymerization calculated from ¹H-NMR spectra (from end group analysis)

^cMolar mass from ¹H-NMR spectra

^dYield vs initial monomer feed

^ePolydispersity index (PDI: $\bar{M}_w/\bar{M}_n$) as measured by GPC