I. M. Okhapkin
A. R. Khokhlov
E. E. Makhaeva

Two-dimensional classification of amphiphilic monomers based on interfacial and partitioning properties. 1. Monomers of synthetic water-soluble polymers

Published online: 22 August 2006
© Springer-Verlag 2006

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00396-005-1342-1> of the article with published errors.

I. M. Okhapkin · A. R. Khokhlov (✉)
Department of Polymer Science, University of Ulm,
Albert-Einstein-Allee 11,
89069 Ulm, Germany
e-mail: khokhlov@polly.phys.msu.ru
Tel.: +7-095-9392959
Fax: +7-095-9392988

I. M. Okhapkin · A. R. Khokhlov
Nesmeyanov Institute of Organoelement Compounds,
Vavilov Street 28,
119991 Moscow, Russia

A. R. Khokhlov · E. E. Makhaeva
Physics Department,
Moscow State University,
119992 Moscow, Russia

Colloid Polym Sci (2005) 284:117–123

On page 120, Fig. 4 is missing as, unfortunately, Fig. 5 was published twice. The correct Fig. 4 is the following. The publisher apologizes for this mistake.

Fig. 4 Two-dimensional diagram of interfacial activity and phase affinity, general view