p73 expression in medulloblastoma: TAp73/ Δ Np73 transcript detection and possible association of p73 α / Δ Np73 immunoreactivity with survival

**Karel Zitterbart · Iva Zavrelova · Jitka Kadlecova ·
Renata Spesna · Adela Kratochvilova ·
Zdenek Pavelka · Jaroslav Sterba**

Published online: 27 September 2008
© Springer-Verlag 2008

**Erratum to: Acta Neuropathol (2007) 114:641–650
DOI 10.1007/s00401-007-0298-2**

Unfortunately, lettering of Fig. 3 was incorrect. This mistake does not affect the conclusions. The correct figure and figure legend are shown here.

The online version of the original article can be found under
doi:[10.1007/s00401-007-0298-2](https://doi.org/10.1007/s00401-007-0298-2).

K. Zitterbart (✉) · Z. Pavelka · J. Sterba
Department of Pediatric Oncology,
University Hospital Brno and Masaryk University,
Cernopolni 9, 613 00 Brno, Czech Republic
e-mail: kzitter@med.muni.cz

I. Zavrelova
Department of Pathology,
University Hospital Brno and Masaryk University,
Brno, Czech Republic

J. Kadlecova · R. Spesna · A. Kratochvilova
Department of Medical Genetics,
University Hospital Brno, Brno, Czech Republic

Fig. 3 p73 α (**a, c, e**) and Δ Np73 (**b, d, f**) immunolabeling of representative medulloblastoma samples. Both diffuse (**a**) and clustered (**c, e**) staining was found in a case of p73 α -positivity, the principle pattern of immunoreactivity was combined nuclear and cytoplasmic. Δ Np73-positivity was ascertained in a subset of p73 α -reactive medulloblastomas only; it was limited to the cytoplasm, and with one exception (**f**), to a lesser extent and intensity (**b, d**), as compared to the case of p73 α . Scale bar: 200 μ m