WenChieh Chen · Chao-Chun Yang · Guan-Yu Chen
Meng-Chie Wu · Hamm-Ming Sheu · Tzong-Shin Tzai

Patients with a large prostate show a higher prevalence of androgenetic alopecia

Published online: 23 September 2005
© Springer-Verlag 2005

Arch Dermatol Res (2004) 296:245-249

In the fifth sentence, the words “serum albumin” were incorrect. The sentence should read as follows:

They fulfilling the clinical diagnosis of BPH defined as (1) prostate volume $> 30 \text{ cm}^3$, measured by transrectal ultrasound, (2) maximal urine flow rate $< 15 \text{ ml/s}$ and mean urine flow rate $< 10 \text{ ml/s}$, and (3) prostate specific antigen $< 10 \text{ ng/ml}$.

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00403-004-0514-z>

WenChieh Chen (✉)
Department of Dermatology, Chang Gung Memorial Hospital,
Kaohsiung, Chang Gung University, Ta-Pei Road 123, Niao-Sung
Hisang, 833 Kaohsiung, Taiwan
E-mail: wchen@cgmh.org.tw

C.-C. Yang · M.-C. Wu · H.-M. Sheu
Department of Dermatology, College of Medicine, National Cheng
Kung University, Sheng-Li Road 138, 704 Tainan, Taiwan

G.-Y. Chen
Department of Dermatology, Armed Forces Taichung General
Hospital, Taichung, Taiwan

T.-S. Tzai
Department of Urology, College of Medicine, National Cheng
Kung University, Sheng-Li Road 138, 704 Tainan, Taiwan