Oxidative stress as determined by glutathione (GSH) concentrations in venous cord blood in elective cesarean delivery versus uncomplicated vaginal delivery

**Orit Paamoni-Keren · Tali Silberstein · Ariela Burg ·
Iris Raz · Moshe Mazor · Oshra Saphier ·
Adi Yekuda Weintraub**

Published online: 3 August 2007
© Springer-Verlag 2007

**Erratum to: Arch Gynecol Obstet (2007) 276:43–46
DOI 10.1007/s00404-006-0304-2**

Due to an unfortunate error, the seventh author name was deleted. The complete list of author names is given below:

Orit Paamoni-Keren, Tali Silberstein, Ariela Burg, Iris Raz,
Moshe Mazor, Oshra Saphier, and Adi Yekuda Weintraub.

The online version of the original article can be found under
doi:[10.1007/s00404-006-0304-2](https://doi.org/10.1007/s00404-006-0304-2).

O. Paamoni-Keren · T. Silberstein · I. Raz ·
M. Mazor · A. Y. Weintraub
Department of Obstetrics and Gynecology,
Soroka University Medical Center,
Faculty of Health Sciences,
Ben-Gurion University of the Negev,
Beer-Sheva, Israel

A. Burg · O. Saphier (✉)
Department Chemical Engineering,
Faculty of Chemical Engineering,
Sami Shamoon College of Engineering,
Beer Sheva 84100, Israel
e-mail: oshras@sce.ac.il