REPLY

Reply to “Questions about and speculations on the incidence of idiopathic polyhydramnios by fetal gender”

Cahit Birdir¹ · Roxana Banica²

Received: 4 March 2015 / Accepted: 9 March 2015 / Published online: 19 March 2015
© Springer-Verlag Berlin Heidelberg 2015

Dear Editor,

I would like to thank Kim et al. [1] on behalf of all authors of our paper for their interest and suggestions.

A multiple regression analysis has not been performed in our study, since our single parameter was the fetal gender to compare with polyhydramnios. Therefore, we do not believe that a multiple regression analysis would be defining the relation of polyhydramnios statistically further.

The measurements of the amniotic fluid index (AFI) in our database have been carried out during the daytime. We did not use any protocol to measure the AFI at night. We, therefore, do not have any collected data of AFI measurements at nighttime.

Although there has been an interesting hypothesis raised, we do not believe that the differences in AFI

measurements can be in any help to predict the nocturnal enuresis. There have been no documented acute or daily changes of the AFI levels in the literature. We could not find any relationship between idiopathic polyhydramnios and postnatal bladder capacity.

Conflict of interest None.

Reference

1. Kim TH, Kim JM, Lee HH (2015) Questions about and speculations on the incidence of idiopathic polyhydramnios by fetal gender. Arch Gynecol Obstet. doi:[10.1007/s00404-015-3688-z](https://doi.org/10.1007/s00404-015-3688-z)

This reply refers to the comment available at
doi:[10.1007/s00404-015-3688-z](https://doi.org/10.1007/s00404-015-3688-z).

✉ Cahit Birdir
cahit.birdir@uk-essen.de

¹ Department of Obstetrics and Gynecology, University Hospital of Essen, Hufelandstr. 55, 45147 Essen, Germany

² Bucur Maternity, Saint John Clinical Emergency Hospital, University of Medicine and Pharmacy, Bucharest, Romania