CORRESPONDENCE

Comments on “Dietary vitamin B2 intake and breast cancer risk: a systematic review and meta-analysis”

Jun Zhang¹

Received: 24 July 2017 / Accepted: 19 September 2017 / Published online: 25 September 2017
© Springer-Verlag GmbH Germany 2017

To the editor,

We read with great interest of the recently published article named “Dietary vitamin B2 intake and breast cancer risk: a systematic review and meta-analysis” by Yu et al. [1], and would like to discuss it with our own opinion. Generally, the systematic review was clearly designed and conducted, while several points may influence the quality of this study.

First, we notice that the authors only searched Medline for eligible studies, this is not recommended due to the potential risk of missing detection of studies. As the Cochrane handbook suggested, at least two databases (e.g. Medline, Embase) should be searched when conducting a systematic review or meta-analysis [2]. Second, the authors claimed they used PRISMA-P [3] to undertake their meta-analysis, while this is not appropriate since PRISMA-P is designed for guiding authors to develop a protocol of systematic review or meta-analysis instead of systematic review or meta-analysis itself [3]. Third, the authors combined the OR/RR from the highest quantile to detect the potential influence of vitamin B2 on breast cancer and got statistically significant pooled results (RR = 0.85, 95% CI: 0.76, 0.95). This method, to some extent, is invalid because the “level” of highest quantile of vitamin B2 varies across studies (from 1.16 to 9.52) making the pooled result of heterogeneous. Fourth, in the table the authors listed all the HR/RR corresponding to each quantile of vitamin B2 on breast cancer, and only one study by Cancarini et al. [4] showed that dietary vitamin B2 may reduce the risk of breast cancer but not other nine studies.

However, the final pooled results are statistically significant and the authors concluded that “dietary vitamin B2 intake is weakly related to the reduced risk of breast cancer”. This should be treated with caution because the results would be unstable—when omitting the study by Cancarini et al. [4], the results may change to negative. The heterogeneous results of subgroup analysis by the authors also support our concern. Though the authors employed dose–response analysis to cover the shortage, this is helpless because they only used a simple linear regression without testing whether this is appropriate (e.g. Ward test [5]).

In conclusion, this is an interesting systematic review and meta-analysis, but the results should be treated with caution. Further strictly designed meta-analysis is needed.

Author contributions JZ: come up with the idea and draft the manuscript

Compliance with ethical standards

Conflict of interest The author declares no conflict of interest.

References

1. Yu L, Tan Y, Zhu L (2017) Dietary vitamin B2 intake and breast cancer risk: a systematic review and meta-analysis. Arch Gynecol Obstet 295(3):721–729
2. Higgins Julian PT, Green S (eds) (2011) Cochrane handbook for systematic reviews of interventions, vol 4. Wiley, Hoboken
3. Moher D, Shamseer L, Clarke M et al (2015) Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. Syst Rev 4(1):1
4. Cancarini I, Krogh V, Agnoli C et al (2015) Micronutrients involved in one-carbon metabolism and risk of breast cancer subtypes. PLoS One 10(9):e0138318
5. Castle JL, Hendry DF (2010) A low-dimension portmanteau test for non-linearity. J Econom 158(2):231–245

✉ Jun Zhang
zhangj1978@yeah.net

¹ Department of Gynaecology and Obstetrics, The First People’s Hospital of Jiangxia District, Wuhan, China