UPPP combined with radiofrequency thermotherapy of the tongue base for the treatment of obstructive sleep apnea syndrome

Emke van den Broek · Wietske Richard ·
Harm van Tinteren · Nico de Vries

Published online: 20 September 2008
© Springer-Verlag 2008

Erratum to: Eur Arch Otorhinolaryngol
DOI 10.1007/s00405-008-0640-x

The third sentence of the second paragraph in “Discussion” should read:

Adding RFTB to UPPP gave an increase in objective success rate of 17.2%, from 37.9 to 55.1% for Friedman stage II patients and an increase of 24.9%, from 8.1 to 33.0% for stage III patients.

The online version of the original article can be found under doi: [10.1007/s00405-008-0640-x](https://doi.org/10.1007/s00405-008-0640-x).

E. van den Broek · W. Richard · N. de Vries (✉)
Department of Otorhinolaryngology, Head and Neck Surgery,
St. Lucas Andreas Hospital, Jan Tooropstraat 164,
1061 AE Amsterdam, The Netherlands
e-mail: n.vries@slaz.nl

E. van den Broek
e-mail: emke_broek@hotmail.com

H. van Tinteren
Department of Biostatistics,
Antoni van Leeuwenhoek Hospital,
Amsterdam, The Netherlands