An animal model for the analysis of cochlear blood flow disturbance and hearing threshold in vivo

**Martin Canis · Warangkana Arpornchayanon ·
Catalina Messmer · Markus Suckfuell ·
Bernhard Olzowy · Sebastian Strieth**

Published online: 22 September 2009
© Springer-Verlag 2009

**Erratum to: Eur Arch Otorhinolaryngol
DOI 10.1007/s00405-009-1036-2**

This article was unfortunately submitted for publication with an error in the title. The correct title is given at the head of this page.

The online version of the original article can be found under doi: [10.1007/s00405-009-1036-2](https://doi.org/10.1007/s00405-009-1036-2).

M. Canis · W. Arpornchayanon · M. Suckfuell · B. Olzowy · S. Strieth
Department of Otorhinolaryngology, Head and Neck Surgery,
University of Munich (LMU), Campus Grosshadern,
Munich, Germany

M. Canis · W. Arpornchayanon · C. Messmer · B. Olzowy · S. Strieth
Walter-Brendel-Centre for Experimental Medicine (WBex),
University of Munich (LMU), Campus Grosshadern,
Munich, Germany

M. Canis (✉)
Ludwig-Maximilians-Universität München,
Klinikum Großhadern, Marchioninstr. 15,
81377 Munich, Germany
e-mail: martin.canis@med.uni-muenchen.de