D. A. Zedgenizov · B. Harte · Edinburgh Ion
Microprobe Facility (EIMF) · V. S. Shatsky
A. A. Politov · G. M. Rylov · N. V. Sobolev

Directional chemical variations in diamonds showing octahedral following cuboid growth

Published online: 14 January 2006
© Springer-Verlag 2006

Contrib Mineral Petrol (2006).
DOI 10.1007/s00410-005-0044-5

In the original article, Fig. 6 was unfortunately incorrect. The correct version is shown here.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00410-005-0044-5>

D. A. Zedgenizov · V. S. Shatsky · G. M. Rylov · N. V. Sobolev
Institute of Mineralogy and Petrography, pr. ak. Koptiyuga 3,
Novosibirsk 630090, Russia

B. Harte (✉)
Grant Institute of Earth Sciences, School of Geosciences,
University of Edinburgh, Edinburgh EH9 3JW, UK
E-mail: ben.harte@ed.ac.uk
Tel.: +44-131-6508528
Fax: +44-131-6683184

A. A. Politov
Institute of Solid State Chemistry and Mechanochemistry,
Kutateladze 18, Novosibirsk 630128, Russia

Fig. 6 Carbon isotope and nitrogen abundance trends for diamonds showing octahedral following cuboid growth; *solid circular* symbols for compositions of cuboid cores; *open diamond* symbols

for outer octahedral regions. Fields for worldwide diamond data from Cartigny et al. (2001) are shown, together with the field of coated diamonds (with cuboid coat on outside of diamond)