A practical method for assessment of dose conversion coefficients for aquatic biota

A. Ulanovsky · G. Pröhl

Published online: 18 October 2006
© Springer-Verlag 2006

Erratum to: Radiat Environ Biophys 45:203–214 DOI 10.1007/s00411-006-0061-4

1. Unfortunately, during the article preparation the fourth column in Table 1 has been corrupted and therefore, published wrong. The corrected values are given below in bold font.

2. Another mistake can be found in the equation in the beginning of the second paragraph of the Appendix. The beginning of the first sentence of this paragraph should read as follows:

“Let $R_0 = \sqrt[3]{abc}$ be a radius of sphere of ...”

Table 1 Shapes and parameters of the aquatic organisms considered in the present study

Surface	Semi-axes	Proportions (a:b:c)	ξ	χ	η
Sphere	$a = b = c$	(1:1:1)	1.0	1.0	1.0
Prolate ovoid	$a = b < c$	(1:1:10)	0.1	0.1	0.59
		(1:1:50)	0.02	0.02	0.35
Oblate ovoid	$a < b = c$	(1:10:10)	1.0	0.1	0.41
		(1:50:50)	1.0	0.02	0.15
Ellipsoid	$a \neq b \neq c$	(1:2:3)	0.33	0.67	0.85
		(1:3:5)	0.6	0.2	0.70
		(1:5:10)	0.5	0.1	0.51
		(1:10:50)	0.2	0.02	0.25

Semi-axes notations follow a general quadric equation: $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$. Parameters ξ , χ , and η are dimensionless

The online version of the original article can be found at
<http://www.dx.doi.org/10.1007/s00411-006-0061-4>.

A. Ulanovsky (✉) · G. Pröhl
GSF—Forschungszentrum für Umwelt und Gesundheit, GmbH,
Ingolstädter Landstraße 1,
85764 Neuherberg, Germany
e-mail: ulanovsky@gsf.de