In focus in HCB

Douglas J. Taatjes¹ · Jürgen Roth²

Accepted: 16 June 2017 / Published online: 22 June 2017
© Springer-Verlag GmbH Germany 2017

Visualization of endothelial barrier function damage

Histochemical determinations of atherosclerosis are often-times assessed by imaging of oil red O-stained cryostat sections (Paigen et al. 1987; Taatjes et al. 2000, 2008; Wadsworth et al. 2002). This useful method can be quantitative and delineates atherosclerotic lesions in various stages of progression, though the earliest stages indicated by endothelial barrier function damage will not be recognized. Bömmel et al. (2017) have now developed a novel, simple staining protocol aimed at assessing this initial phase of atherosclerosis formation in large vessels. Evans blue dye was injected intracardially into mice following washout of blood. The aortae were then removed, fixed, and frozen. Endothelial barrier permeability disruptions were indicated by the presence of Evans Blue dye in the intimal region of the aorta from three mouse models known to develop atherosclerotic lesions with differing degrees of severity (ApoE^{-/-}, LDLr^{-/-}, Ccl^{-/-}). Evans blue deposition in the aortic intima was found in all three mouse models prior to light microscopically visualized morphological changes, and could be easily quantitated using standard image analysis parameters. The authors also documented the presence of endothelial cell barrier disfunctions using several in vitro approaches. Overall, this novel histochemical technique promises to provide a simple and inexpensive

assay for evaluating the earliest stages of atherosclerotic lesion formation and the potential effects of pharmaceutical interventions.

Epithelial changes in amikacin-damaged organ of Corti

The organ of Corti located in the cochlear of the inner ear contains the hair cells, which serve as the sensory cells for hearing. Many types of insults, including acoustic trauma from loud noise and ototoxic drugs can damage hair cells resulting in hearing loss. Following this trauma, epithelial reorganization occurs in the cochlear whereby specialized columnar epithelial cells are replaced by an unspecialized flattened epithelium (Oesterle 2013). Ladrech et al. (2017) have now used a model of amikacin (an aminoglycoside antibiotic) induced cochlear remodeling following hair cell loss to study these processes using immunohistochemistry. Using samples available from prior studies, they investigated in a spatio-temporal fashion proteins involved in (1) maintaining cell shape (actin and tau); (2) cell adhesion (E-cadherin, polysialylated NCAM, laminin); and (3) cell migration (vimentin, GFAP), and (4) enzymatic activities (PKC isoforms). Moreover, morphological changes in the cochlear induced by amikacin poisoning were investigated by both scanning and transmission electron microscopy. Based upon their immunofluorescence and electron microscopic investigations, the authors determined that following severe hair cell loss, the reorganization of the organ of Corti is dependent upon several cellular processes, including epithelial-to-mesenchymal transition as well as individual and collective cellular migration. Moreover, structural reorganization appeared to involve both migratory root cells and infiltrating fibroblasts. Finally, their results suggest that the

✉ Douglas J. Taatjes
douglas.taates@uvm.edu

¹ Department of Pathology and Laboratory Medicine, Larner College of Medicine, University of Vermont, Burlington, VT 05405, USA

² University of Zurich, 8091 Zurich, Switzerland

unspecialized flattened epithelium arising following hair cell damage may represent a potential therapeutic target for sensory cell regeneration.

Visualization of genetic heterogeneity in morphologically homogenous colorectal cancer

Colorectal cancer may be composed of subpopulations of morphological distinct cancer cells harboring different genetic alterations. This cancer cell/tumor heterogeneity is considered a major cause for therapy resistance. Because of the lack of analogous data, El-Heliebi et al. (2017) have investigated morphologically homogenous colorectal carcinomas with regard to their genetic landscape. Using next-generation sequencing, they identified in a first step mutational profiles of four microdissected areas of six morphologically homogenous colorectal cancers. In three out of the six studied carcinomas, genetic heterogeneity was obvious. Afterwards, they applied mRNA-based in situ mutation detection technology based on padlock probes for direct mutation detection in sections of formaldehyde-fixed and paraffin-embedded cancer tissue. The padlock probe technique is highly specific and permits the detection of point mutations in situ in tissue sections (Larsson et al. 2010). Using this in situ mutation detection, El-Heliebi et al. (2017) were able for instance to clearly localize not only FANCD2 wild type, but also mutated transcripts in certain tumor areas. They concluded that genetic heterogeneity of morphologically homogenous colorectal cancer may be frequent and that the padlock probe technique is a powerful in situ tool for their visualization.

Mucosal barrier in human chagasic-megacolon mucosa

Chagas disease is an illness caused by the protozoan parasite *Trypanosoma cruzi*. During the chronic phase, the parasite exists mainly in the heart and muscle of the digestive system, causing cardiomyopathy and/or intestinal dilatation such as megacolon (Rassi et al. 2010). Patients suffering from chagasic megacolon have an intact mucosal barrier, and while the selective survival of VIP-ergic nerve fibres within the lamina propria has been demonstrated, a loss of cholinergic neurons occurs in the myenteric and submucosal plexus (Jabari et al. 2014a, b). In the present investigation Koch et al. (2017) have analyzed goblet cells [including their markers mucin 2 (MUC2) and trefoil factor 3 (TFF3)], as well as enterocytes, and the relation of VIP-immunoreactive fibers to these epithelial cell types. Goblet cells accounted for 38.4% of all epithelial cells in normal colon, and increased to 58.9% in chagasic-megacolon

mucosa. In megacolon mucosa, TFF3 was confined to the goblet cells in the base of the crypts and MUC2 was detectable only in luminal goblet cells, in contrast to their uniform expression in normal colon. Moreover, in contrast to normal colon, gelsolin-positive goblet cells were almost undetectable in megacolon mucosa. The surviving VIP-ergic nerve fibers were always in close proximity to goblet cells. The mitosis rate was significantly increased in the megacolon and anal parts. The observed changes may be contributing factors to maintain an intact mucosal barrier in chagasic megacolon for decades.

References

- Bömmel H, Kleefeldt F, Zerneck A, Ghavampour S, Wagner N, Kuersten S, Ergun S (2017) Visualization of endothelial barrier damage prior to formation of atherosclerotic plaques. *Histochem Cell Biol*. doi:[10.1007/s00418-017-1562-8](https://doi.org/10.1007/s00418-017-1562-8)
- El-Heliebi A, Kashofer K, Fuchs J, Jahn SW, Viertler C, Matak A, Sedlmayr P, Hoefler G (2017) Visualization of tumor heterogeneity by in situ padlock probe technology in colorectal cancer. *Histochem Cell Biol*. doi:[10.1007/s00418-017-1557-5](https://doi.org/10.1007/s00418-017-1557-5)
- Jabari S, da Silveira AB, de Oliveira EC, Quint K, Wirries A, Neuhuber W, Brehmer A (2014a) Mucosal layers and related nerve fibres in non-chagasic and chagasic human colon—a quantitative immunohistochemical study. *Cell Tissue Res* 358:75–83. doi:[10.1007/s00441-014-1934-5](https://doi.org/10.1007/s00441-014-1934-5)
- Jabari S, de Oliveira EC, Brehmer A, da Silveira AB (2014b) Chagasic megacolon: enteric neuron and related structures. *Histochem Cell Biol* 142:235–244. doi:[10.1007/s00418-014-1250-x](https://doi.org/10.1007/s00418-014-1250-x)
- Koch C, da Silveira ABM, de Oliveira EC, Quint K, Neuhuber W, Brehmer A, Jabari S (2017) Epithelial cell types and their proposed roles in maintaining the mucosal barrier in human chagasic-megacolon mucosa. *Histochem Cell Biol*. doi:[10.1007/s00418-017-1563-7](https://doi.org/10.1007/s00418-017-1563-7)
- Ladrech S, Eybalin M, Puel J-L, Lenoir M (2017) Epithelial–mesenchymal transition, and collective and individual cell migration regulate epithelial changes in the amikacin-damaged organ of Corti. *Histochem Cell Biol*. doi:[10.1007/s00418-017-1548-6](https://doi.org/10.1007/s00418-017-1548-6)
- Larsson C, Grundberg I, Söderberg O, Nilsson M (2010) In situ detection and genotyping of individual mRNA molecules. *Nat Methods* 7:395–397. doi:[10.1038/nmeth.1448](https://doi.org/10.1038/nmeth.1448)
- Oesterle EC (2013) Changes in the adult vertebrate auditory sensory epithelium after trauma. *Hear Res* 297:91–98
- Paigen B, Morrow A, Holmes PA, Mitchell D, Williams RA (1987) Quantitative assessment of atherosclerotic lesions in mice. *Atherosclerosis* 68:231–240
- Rassi A Jr, Rassi A, Marin-neto JA (2010) Chagas disease. *Lancet* 375:1388–1402. doi:[10.1016/S0140-6736\(10\)60061-X](https://doi.org/10.1016/S0140-6736(10)60061-X)
- Taatjes DJ, Wadsworth MP, Schneider DJ, Sobel BE (2000) Improved quantitative characterization of atherosclerotic plaque composition with immunohistochemistry, confocal fluorescence microscopy, and computer-assisted image analysis. *Histochem Cell Biol* 113:161–173
- Taatjes DJ, Wadsworth MP, Quinn AS, Rand JH, Bovill EG, Sobel BE (2008) Imaging aspects of cardiovascular disease at the cell and molecular level. *Histochem Cell Biol* 130:235–245
- Wadsworth MP, Sobel BE, Schneider DJ, Taatjes DJ (2002) Delineation of the evolution of compositional changes in atheroma. *Histochem Cell Biol* 118:59–68