Erratum to: Case–control study of knee osteoarthritis and lifestyle factors considering their interaction with physical workload

Ilias Vrezas · Gine Elsner · Ulrich Bolm-Audorff ·
Nasreddin Abolmaali · Andreas Seidler

Published online: 6 May 2010
© Springer-Verlag 2010

**Erratum to: Int Arch Occup Environ Health (2010)
83:291–300
DOI 10.1007/s00420-009-0486-6**

In the original paper, there is a mistake in the calculation of population-attributable risks: Applying formula (3) as reported by Coughlin et al. (1994) [$AR = P(E/D) * ((RR - 1)/RR)$, where $P(E/D)$ is the proportion of exposed cases], for persons with elevated BMI in combination with

moderate to high exposure to occupational kneeling/squatting, the population attributable risk (PAR) was not 4% (as stated in the abstract, the results section, and the discussion), but 19%. Furthermore, the PAR for elevated BMI in combination with moderate to high exposure to occupational lifting/carrying of loads was not 7%, but 24%. With correct PAR values, the last part of the “Results” section “Population attributable risks (PAR) for BMI and physical workload” should read as follows:

- The adjusted population attributable risk (PAR) for a BMI of 22.86 or more compared with a BMI of less than 22.86 was 59% (no table). The adjusted PAR for kneeling/squatting for 4,757 h or more was 17% (no table). The adjusted PAR for occupational lifting and carrying of weights $\geq 5,120$ kg*hours was 23%.
- When population attributable risks were calculated for the combination of BMI elevations and occupational exposures, for persons with a BMI ≥ 24.92 kg/m² exposed to kneeling/squatting for 4,757 h or more, the PAR was 19%. The population attributable risk for the combined exposure to BMI ≥ 24.92 kg/m² and occupational lifting/carrying of weights $\geq 5,120$ kg*hours was 24%. When persons with occupational lifting/carrying of weights ≥ 630 kg*hours were regarded as exposed, the population attributable risk for the combined exposure to BMI ≥ 24.92 kg/m² and occupational lifting/carrying increased to 37%.

The online version of the original article can be found under doi:[10.1007/s00420-009-0486-6](https://doi.org/10.1007/s00420-009-0486-6).

I. Vrezas
Cyprus International Institute for Environmental and Public Health in Association with Harvard School of Public Health, Cyprus University of Technology, Corner of Athinon Road & N. Xiouta Street, 3041 Lemesos, Cyprus

G. Elsner
Institute of Occupational Medicine,
Johann Wolfgang Goethe-University,
Theodor-Stern-Kai 7, 60590 Frankfurt/Main, Germany

U. Bolm-Audorff
Labour Inspection, Division of Occupational Health,
Simone-Veil-Str. 5, 65197 Wiesbaden, Germany,
and Associated Professor of Occupational Health,
Giessen University, Giessen, Germany

N. Abolmaali
OncoRay, Center for Radiation Research in Oncology,
Fetscherstraße 74, P.O. Box 86, 01307 Dresden, Germany

A. Seidler (✉)
Institute and Policlinic for Occupational and Social Medicine,
Fetscherstr. 74, 01307 Dresden, Germany
e-mail: Andreas.Seidler@mailbox.tu-dresden.de

Reference

Coughlin SS, Benichou J, Weed DL (1994) Attributable risk estimation in case-control studies. *Epidemiol Rev* 16:51–64