ERRATUM

Takahide Kato · Yoshinori Matsumura
Atsuko Tsukanaka · Takeshi Harada
Mitsuo Kosaka · Nobuo Matsui

Effect of low oxygen inhalation on changes in blood pH, lactate, and ammonia due to exercise

Published online: 31 March 2004
© Springer-Verlag 2004

Eur J Appl Physiol (2004) 91:296–302

In Table 1, the values were not correctly aligned. The correct version is given below.

Table 1 Exercise, cardiorespiratory and blood gas parameters influenced by exhaustive work at normoxia (*MAX*) and at hypoxia (*HYPOXIA*), as well as at the workload imposed during hypoxia but carried out at normoxia (*SUBMAX*) in six subjects. Mean values with standard deviations

Indicated are parameters determined in the state of *HYPOXIA*, which differ significantly ($P < 0.05$):
*from the state of *MAX*,
§from the state of *SUBMAX*

		MAX	HYPOXIA	SUBMAX
Performance time (min)		14''48 (2''10)	11''01 (0''59)*	11''01 (0''59)*
Maximal workload (watt)		290 (45)	213 (33)*	213 (33)*
$\dot{V}O_2$ max ($\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$)		45.25 (2.12)	36.73 (5.03)*	39.12 (3.17)*
Heart rate ($\text{beats}\cdot\text{min}^{-1}$)	Pre-exercise	71.7 (4.9)	82.5 (9.7)*§	73.0 (6.7)
	Exhaustion	195.3 (4.7)	188.2 (3.0)*§	174.5 (7.4)*
	At 30 min post-exercise	102.2 (9.2)	97.5 (10.6)	82.7 (18.2)*
V_E ($\text{l}\cdot\text{min}^{-1}$)	Pre-exercise	10.3 (1.0)	12.8 (3.0)*§	9.5 (0.9)
	Exhaustion	118.1 (17.0)	116.5 (22.0)§	75.3 (11.1)*
	At 30 min post-exercise	16.1 (3.3)	14.7 (3.2)§	11.2 (1.7)*
Blood pH	Pre-exercise	7.401 (0.016)	7.437 (0.053)	7.396 (0.034)
	Minimum value	7.178 (0.059)	7.246 (0.067)*	7.295 (0.033)*
	At 30 min post-exercise	7.337 (0.038)	7.407 (0.029)*	7.381 (0.027)*
Blood HCO_3^- (mM)	Pre-exercise	24.0 (1.5)	27.1 (2.1)*	24.9 (3.0)
	Minimum value	12.8 (2.7)	13.4 (2.8)§	18.4 (1.4)*
	At 30 min post-exercise	18.7 (3.1)	19.5 (2.3)§	23.1 (2.5)*
Base excess (mEq/l)	Pre-exercise	0.2 (1.3)	3.3 (0.8)*	0.6 (2.1)
	Minimum value	-15.4 (3.2)	-12.8 (4.0)§	-7.5 (2.1)*
	At 30 min post-exercise	-5.6 (3.3)	-3.5 (2.2)	-1.2 (2.5)*

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00421-003-0975-3>

T. Kato (✉) · Y. Matsumura · A. Tsukanaka · T. Harada
M. Kosaka · N. Matsui
Laboratory for Exercise Physiology and Biomechanics,
Chukyo University, 101 Tokodachi, Kaizu-cho Toyota,
470-0393 Aichi, Japan
E-mail: tkato@seiken.sass.chukyo-u.ac.jp
Tel.: +81-565-461211
Fax: +81-565-461272