Murat Karabulut · Scott E. Crouter
David R. Bassett Jr

Comparison of two waist-mounted and two ankle-mounted electronic pedometers

Published online: 20 January 2006
© Springer-Verlag 2006

Eur J Appl Physiol (2005): 335–343

Parts c and d of Fig. 2 were missing. The complete figure is given below.

The online version of the original article can be found at <http://dx.xoi.org/10.1007/s00421-005-0018-3>

M. Karabulut
Department of Health and Exercise Science,
University of Oklahoma, Norman, OK, USA

D. R. Bassett Jr
Department of Exercise, Sport, and Leisure Studies,
The University of Tennessee, 1914 Andy Holt Avenue,
Knoxville, TN 37996, USA

S. E. Crouter (✉)
Cornell University, Division of Nutritional Sciences,
279 MVR, Ithaca, NY 14853, USA
E-mail: sec62@cornell.edu
Tel.: +1-607-2546403
Fax: +1-607-2548375

Fig. 2 Modified Bland–Altman plots depicting error scores (actual steps per minute minus pedometer steps per minute) for each pedometer across all walking speeds: **a** Step Watch 3 (SW-3_{Ankle}),

b AMP-331 (AMP_{Ankle}), **c** New Lifestyle NL-2000 (NL_{Waist}), **d** Digiwalker SW-701 (SW-701_{Waist}). *Dashed lines* represent mean difference; *solid lines* represent 95% prediction intervals