Kunihiko Nomura · Yoshiaki Takei · Masaki Yoshida
Yasuyoshi Yanagida

Phase-dependent chronotropic response of the heart during running in humans

Published online: 21 April 2006
© Springer-Verlag 2006

Eur J Appl Physiol (2006)
DOI 10.1007/s00421-005-0103-7

The equation to obtain the instantaneous phase was described in the Data analyses section of the Methods of this article as $\phi(t) = 2\pi (t - t_k/t_{k+1} - t_k) + 2\pi k$. This equation should have been:

$$\phi(t) = 2\pi \frac{t - t_k}{t_{k+1} - t_k} + 2\pi k.$$

Furthermore, the equation to obtain the generalized relative phase was described in the Data analyses as $\Psi_a - b = 1/2\pi (\phi_b(t_k) \bmod 2\pi)$. This should have been as follows:

$$\Psi_{a-b} = \frac{1}{2\pi} (\phi_b(t_k) \bmod 2\pi).$$

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00421-005-0103-7>

K. Nomura (✉) · M. Yoshida
Department of Biomedical Engineering,
Osaka Electro-Communication University,
1130-70 Kiyotaki Shijonawate-shi, Osaka 575-0063, Japan
E-mail: n-kuni@isc.osakac.ac.jp
Tel.: +81-72-8765185
Fax: +81-72-8765199

Y. Takei · Y. Yanagida
Faculty of Human Development, Kobe University,
3-11 Tsurukabuto Nada-ku Kobe-shi, Hyogo 657-8501, Japan