ERRATUM

Erratum to: Eicosapentaenoic and docosahexaenoic acids-rich fish oil supplementation attenuates strength loss and limited joint range of motion after eccentric contractions: a randomized, double-blind, placebo-controlled, parallel-group trial

Yosuke Tsuchiya¹ · Kenichi Yanagimoto² · Koichi Nakazato³ · Kohsuke Hayamizu⁴ · Eisuke Ochi⁵

Published online: 21 July 2016
© Springer-Verlag Berlin Heidelberg 2016

**Erratum to: Eur J Appl Physiol (2016) 116:
1179–1188
DOI 10.1007/s00421-016-3373-3**

The author would like to correct the following errors in the print publication of the article.

In page 1181, under the supplement section, the first sentence should read as follows:

The EPA group consumed eight 300 mg EPA and DHA-rich fish oil softgel capsules (Nippon Suisan Kai-sha Ltd., Tokyo, Japan) per day (containing 600 mg EPA and 260 mg DHA in a total of 2400 mg)

Unfortunately, the Fig. 2 was published incorrectly in the original publication of the article. The correct figure is given below:

The online version of the original article can be found under doi:[10.1007/s00421-016-3373-3](https://doi.org/10.1007/s00421-016-3373-3).

✉ Eisuke Ochi
ochi@hosei.ac.jp

- ¹ Laboratory of Health and Sports Sciences, Meiji Gakuin University, Kanagawa, Japan
- ² Human Life Science R&D Center, Nippon Suisan Kaisha, Ltd., Tokyo, Japan
- ³ Graduate School of Health and Sport Science, Nippon Sport Science University, Tokyo, Japan
- ⁴ General Health Medical Center, Yokohama University of Pharmacy, Kanagawa, Japan
- ⁵ Faculty of Bioscience and Applied Chemistry, Hosei University, Tokyo, Japan