ERRATUM

Kanji Mori · Tokuhiro Chano · Ryoji Kushima
Sinsuke Hukuda · Hidetoshi Okabe

Expression of E-cadherin in chordomas: diagnostic marker and possible role of tumor cell affinity

Published online: 1 June 2002
© Springer-Verlag 2002

Virchows Arch (2002) 440:123–127

The following Table 1 should have been included:

Table 1 Clinicopathological features in seven cases of chordoma

Case	Age	Gender	Location	Primary (P) or recurrent (R) chordoma, or metastasis (M)	Pathological findings		
					Cytokeratin	E-cadherin	Ep-CAM
1	60	F	Sacrum	P	++	+	–
2	75	M	Sacrum	R	++	+	–
3	60	M	C3	P	++	+	–
4	69	M	Intracranial	R	++	+	–
5	37	M	Intracranial	P	++	+	–
6	38	F	Intracranial ^a	P	+ ^b	+	–
7	21	F	Intracranial ^a	P	+ ^b	+	–
			Spinal cord (C3–7)	M	++	+	–
			Thoracic wall	M	++	+	–

^a Chondroid chordoma

^b Only a few cells were positive

The online version of the original article can be found at <http://dx.doi.org/10.1007/s004280100525>

K. Mori (✉) · S. Hukuda
Department of Orthopedic Surgery,
Shiga University of Medical Science, Tsukinowa-cho, Seta, Otsu,
Shiga, 520-2192, Japan
e-mail: kanchi@belle.shiga-med.ac.jp
Tel.: +81-77-5482252, Fax: +81-77-5482254

T. Chano · R. Kushima · H. Okabe
Department of Clinical Laboratory Medicine,
Shiga University of Medical Science, Tsukinowa-cho, Seta, Otsu,
Shiga, 520-2192, Japan