Expression of estrogen receptor co-regulators NCoR and PELP1 in epithelial cells and myofibroblasts of colorectal carcinomas. Cytoplasmic translocation of NCoR in epithelial cells correlates with better prognosis

Vassiliki Tzelepi · Petros Grivas ·
Zinovia Kefalopoulou · Haralabos Kalofonos ·
John N. Varakis · Georgia Sotiropoulou-Bonikou

Published online: 23 December 2008
© Springer-Verlag 2009

Erratum to: Virchows Arch
DOI 10.1007/s00428-008-0708-4

The title of the article was rendered incorrectly.
The correct title is as given above.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00428-008-0708-4>.

V. Tzelepi · Z. Kefalopoulou · J. N. Varakis ·
G. Sotiropoulou-Bonikou (✉)
Department of Anatomy and Histology-Embryology,
University of Patras,
26500 Rion, Greece
e-mail: sotiropouloug@yahoo.com

V. Tzelepi · Z. Kefalopoulou
Department of Pathology, University of Patras,
26500 Rion, Greece

P. Grivas · H. Kalofonos
Department of Internal Medicine-Division of Oncology,
Medical School, University of Patras,
26500 Rion, Greece

P. Grivas
Department of Internal Medicine, Hahnemann University
Hospital/Drexel University College of Medicine,
Philadelphia, PA 19102, USA