Erratum to: Respiratory hospitalizations and respiratory syncytial virus prophylaxis in special populations

B. Paes · I. Mitchell · A. Li · K. L. Lanctôt

Published online: 20 June 2012
© Springer-Verlag 2012

Erratum to: Eur J Pediatr
DOI 10.1007/s00431-011-1654-8

On page 835, in the first paragraph of the Utilization and Compliance section, the overall average number of injections should be 3.9 ± 1.5 , not 3.6 ± 1.5 . As well, the average injections for Special Populations (Group 2) and Premature Infants (Group 1) were reversed. On average, Special Populations (Group 2) had 4.2 ± 1.6 injections and Premature Infants (Group 1) had 3.8 ± 1.7 injections.

In Figure 2B, the caption is missing the following: “Note: In figure b the solid and dotted lines overlap and appear as a single entity.”

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00431-011-1654-8>.

B. Paes
Department of Pediatrics, McMaster University,
Hamilton, Ontario, Canada

I. Mitchell
Department of Pediatrics, University of Calgary,
Calgary, Alberta, Canada

A. Li · K. L. Lanctôt (✉)
Medical Outcomes and Research in Economics (MORE®),
Research Group, Sunnybrook Health Sciences Centre,
University of Toronto,
2075 Bayview Avenue, Room FG-05,
Toronto, Ontario M4N 3M5, Canada
e-mail: krista.lanctot@sunnybrook.ca