ERRATUM

Erratum to: Long-term liposteroid therapy for idiopathic pulmonary hemosiderosis

Takehiko Doi¹ · Shouichi Ohga^{1,2} · Masataka Ishimura¹ · Hidetoshi Takada¹ · Kanako Ishii¹ · Kenji Ihara¹ · Hideyuki Nagai³ · Toshiro Hara¹

Published online: 30 September 2015
© Springer-Verlag Berlin Heidelberg 2015

Erratum to: Eur J Pediatr (2013) 172:1475–1481
DOI 10.1007/s00431-013-2065-9

The errata are noted below.

Abstract

The sentence, “Liposteroid was intravenously infused at 0.8 mg/kg/day for three consecutive days at the time of acute bleeding.” was corrected to “Liposteroid was intravenously infused at 0.05–0.08 mg (of dexamethasone)/kg/day for three consecutive days at the time of acute bleeding.”

Page 1476, Methods, Liposteroid therapy, line 4–5

The sentence, “It was intravenously infused at 0.8 mg/kg/day for three consecutive days at the time of acute bleeding, after the obtainment of informed consent from parents.” was corrected to “It was intravenously infused at 0.05–0.08 mg (of dexamethasone)/kg/day for three consecutive days at the time of acute bleeding, after the obtainment of informed consent from parents.”

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00431-013-2065-9>.

✉ Shouichi Ohga
ohgas@pediatr.med.kyushu-u.ac.jp

- ¹ Department of Pediatrics, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan
- ² Department of Perinatal and Pediatric Medicine, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan
- ³ Division of Pediatrics, Kyoto Yamashiro General Medical Center, 27-1 Ikeda, Kizu, Kizukawa, Kyoto 619-0214, Japan