Larval convergence in a colonial tunicate: the organization of the sarcotubular complex in *Ecteinascidia turbinata* (Perophoridae, Phlebobranchiata, Tunicata, Chordata)

Thomas Stach · Andreas Kirbach

Published online: 9 January 2009
© Springer-Verlag 2008

Erratum to: Zoomorphology
DOI 10.1007/s00435-008-0069-1

Due to an unfortunate error the legend of the Fig. 7 was incorrect. The correct legend is given here.

The online version of the original article can be found under
doi: [10.1007/s00435-008-0069-1](https://doi.org/10.1007/s00435-008-0069-1).

T. Stach (✉) · A. Kirbach
Institut für Biologie, Chemie, Pharmazie,
Zoologie, Evolution und Systematik der Tiere,
Freie Universität Berlin, Königin-Luise-Straße 1-3,
14195 Berlin, Germany
e-mail: tstach@zoosyst-berlin.de

Fig. 7 Schematic representation of different types of the sarcotubular complex found in chordates. See Table 2 for information on taxonomic distribution, meristic variation, and data source. **a** Flat, tubular invaginations. **b** Tubular invaginations. **c** Laminar invagination. **d** Invaginations

lacking. **e** Invaginations are also lacking in the specialized muscle plates of cephalochordates. **f** Tubular invaginations in craniates. *mf* myofibril; *pc* peripheral coupling; *si* invagination of the sarcolemma; *sl* sarcolemma; *sr* sarcoplasmic reticulum