Gereon Schares · Alfred Otto Heydorn · Arnd Cüppers
Heinz Mehlhorn · Lutz Geue · Martin Peters
Franz Josef Conraths

In contrast to dogs, red foxes (*Vulpes vulpes*) did not shed *Neospora caninum* upon feeding of intermediate host tissues

Published online: 27 March 2002
© Springer-Verlag 2002

Parasitol Res (2002) 88:44–52

The alteration of one word during editing the above mentioned article led to a wrong statement at the end of the discussion (last paragraph, first sentence). The mistake remained un-observed during proof-reading. The sentence “By bioassay, we can **conclude** that there were more than 3,700 and 200 oocysts of our

N. caninum-isolate among the *Hammondia* sp.-like oocysts collected during the first and the second fox-feeding experiments, respectively.” has to be corrected into “By bioassay, we can **exclude** that there were more than 3,700 and 200 oocysts of our *N. caninum*-isolate among the *Hammondia* sp.-like oocysts collected during the first and the second fox-feeding experiments, respectively.”

The online version of the original article can be found at <http://dx.doi.org/10.1007/s004360100496>

G. Schares (✉) · L. Geue · F.J. Conraths
Institute for Epidemiological Diagnostics,
Federal Research Centre for Virus Diseases of Animals,
Seestrasse 55, 16868 Wusterhausen, Germany
E-mail: g.schares@wus.bfav.de
Fax: +49-33979-80222

A.O. Heydorn
Institute for Parasitology
and Tropical Veterinary Medicine,
Freie Universität Berlin,
Königsberg 67, 14163 Berlin, Germany

A. Cüppers · H. Mehlhorn
Institut für Zoomorphologie,
Zellbiologie und Parasitologie,
Heinrich-Heine-Universität,
Universitätsstrasse 1, 40225 Düsseldorf, Germany

M. Peters
Institute for Pathology,
Tierärztliche Hochschule Hannover,
Bünteweg 17, 30559 Hannover, Germany