ERRATUM

Koichi Koyama

Dendritic cell expansion occurs in mesenteric lymph nodes of B10.BR mice infected with the murine nematode parasite *Trichuris muris*

Published online: 11 August 2005
© Springer-Verlag 2005

Parasitol Res (2005) DOI 10.1007/s00436-005-1427-2

Unfortunately, Table 1 was published with errors. The correct Table 1 is given here.

Table 1 Kinetics of CD11c⁺ B220⁺ cells in MLNs of *Trichuris muris*-infected B10.BR mice

Days p.i.	Percentage of positive staining cells ^a				Total cells isolated (×10 ⁷) ^b
	CD11c ⁺ B220 ⁺	CD4 ⁺	CD8 ⁺	B220 ⁺	
Uninfected	1.0 ± 0.2 ^c	39.2 ± 1.4	23.4 ± 2.5	32.0 ± 2.9	2.70 ± 0.77
14	0.9 ± 0.2	39.1 ± 2.5	21.6 ± 2.0	35.3 ± 4.9	4.05 ± 0.79*
20	2.0 ± 0.2**	30.7 ± 3.0**	18.0 ± 1.3**	44.3 ± 2.3**	6.40 ± 1.18**
25	1.0 ± 0.3	31.8 ± 3.2***	18.6 ± 2.3*	42.9 ± 4.1***	4.55 ± 0.65**
32	0.9 ± 0.1	36.7 ± 3.3	20.1 ± 2.2	38.6 ± 4.7*	4.76 ± 1.34*

^aMLNCs were prepared from the MLNs of uninfected or *Trichuris muris*-infected B10.BR mice on days 14, 20, 25, and 32 p.i. MLNCs were stained with PE-anti-CD11c and FITC-anti-B220 MoAbs, or PE-anti-CD4 and FITC-anti-CD8 MoAbs. Stained cells were then analyzed using a FACSscan

^bTotal numbers of mononuclear cells isolated from the MLNs of uninfected and infected mice

^cResults represent the mean ± SD for five mice at each time-point and are representative of three independent experiments that gave the similar results

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$ compared with uninfected controls

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00436-005-1427-2>

K. Koyama
Department of Parasitology, Kitasato University School
of Medicine, 1-15-1 Kitasato, Sagamihara Kanagawa,
228-8555, Japan
E-mail: koyama@kitasato-u.ac.jp
Fax: +81-42-7789312