Hagit Gill · Ilan Paperna

Leucocytozoonosis in the Israeli sparrow, *Passer domesticus biblicus* Hartert 1904

Published online: 20 September 2005
© Springer-Verlag 2005

Parasitol Res (2005) 96:373–377

In the online PDF and the print version, Figs. 5–8 are duplicates of Figs. 1–4. The correct Figs. 5–8 are given here.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00436-005-1352-4>

H. Gill · I. Paperna (✉)
Department of Animal Sciences, Faculty of Agricultural,
Food and Environmental Quality Sciences, The Hebrew University
of Jerusalem, Rehovot, 76100, Israel
E-mail: paperna@agri.huji.ac.il
Tel.: +972-8-9489945
Fax: +972-8-9465763

Figs. 5-8 Kidney histology of *Leucocytozoon*-infected sparrow; MHE, same scale bar 15 μ m

Fig. 5 Congestion of blood vessels by gametocytes (C) with gradual vacuolar degeneration of the tubular and interstitial renal tissue (V). One tubule contains merogonic stages (Z)

Fig. 6 Assembly of degenerated and necrotic kidney tubules (arrows); meronts present in the intact tubular tissue (Z)

Fig. 7 Longitudinal view of necrotized tubule (D), adjoining congested blood vessel (C); cells of the kidney tubule contain *Leucocytozoon* young meronts (arrows)

Fig. 8 *Leucocytozoon* gametocytes nested in the blood vessels (fine arrows) of the glomerulus (bold arrow)