Erratum to: Haemin enhances the in vivo efficacy of artemether against juvenile and adult *Schistosoma mansoni* in mice

Naglaa Mohamed El-Lakkany · Sayed Hassan Seif el-Din

Published online: 18 April 2013
© Springer-Verlag Berlin Heidelberg 2013

Erratum to: Parasitol Res
DOI 10.1007/s00436-013-3358-7

The original version of this article inadvertently contained a mistake. Two values found in the column entries of Tables 1 and 4 were incorrect. Please see below for the correct entries.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00436-013-3358-7>.

N. M. El-Lakkany (✉) · S. H. Seif el-Din
Department of Pharmacology, Theodor Bilharz Research Institute,
Warrak El-Hadar, Imbaba, PO Box 30, Giza 12411, Egypt
e-mail: naglaellakkany@yahoo.com

Table 1 Effect of haemin on artemether efficacy against juvenile (28 days PI) and adult (49 days PI) *S. mansoni* worms in relation to mice hemoglobin concentration

Animal groups	Juvenile <i>S. mansoni</i>				Adult <i>S. mansoni</i>			
	Hb (g/dL)		Art efficacy (3 weeks post Art treatment)		Hb (g/dL)		Art efficacy (3 weeks post Art treatment)	
	72 h post-Art treatment	3 weeks post-Art treatment	Total females	Total worms	72 h post-Art treatment	3 weeks post-Art treatment	Total females	Total worms
Uninfected untreated	11.85±1.11	10.39±0.88	-	-	12.76±2.54	11.64±1.76	-	-
Infected untreated	8.17±0.82 ^a	6.43±0.67 ^b	11.58±1.46	27.43±3.56	5.47±0.44 ^a	4.11±0.63 ^b	12.43±2.23	26.13±3.68
Haemin	10.46±1.47	7.12±1.13 ^a	11.35±2.17 (2)	27.02±3.15 (1.50)	9.16±1.88	7.89±1.32 ^c	12.26±1.88 (1.37)	25.95±2.14 (0.69)
Art	7.43±0.95 ^b	6.32±0.75 ^b	1.51±0.17 ^c (87.04)	6.86±0.95 ^c (75)	6.34±0.98 ^a	5.34±0.86 ^b	3.36±0.21 ^d (72.97)	10.71±1.89 ^d (59.01)
Haemin + Art	11.23±1.11 ^{c, f}	8.76±1.14	0.35±0.01 ^{e, h} (96.98)	2.47±0.16 ^{e, h} (91)	11.76±2.01 ^{d, f}	8.37±0.89 ^{d, f}	1.06±0.09 ^{e, h} (91.47)	4.34±0.38 ^{e, g} (83.39)

Values are presented as mean ± SEM. Number of animals/group = 8. Numbers between parentheses indicate the percentage of reduction from infected untreated group

^a $p<0.05$, ^b $p<0.01$ (significant difference from uninfected untreated); ^c $p<0.05$, ^d $p<0.001$ (significant difference from infected untreated); ^e $p<0.05$, ^f $p<0.01$, ^g $p<0.001$ (significant difference from Art-treated)

Table 4 Glutathione peroxidase (GPx) activities in juvenile (28 days PI) and adult (49 days PI) *S. mansoni* worms recovered 72 h and 3 weeks post-artemether treatment, from mice treated with artemether alone and in addition to haemin

Animal groups	Juvenile <i>S. mansoni</i>				Adult <i>S. mansoni</i>			
	72 h post-Art treatment		3 weeks post-Art treatment (7 weeks PI)		72 h post-Art treatment		3 weeks post-Art treatment (10 weeks PI)	
	Male	Female	Male	Female	Male	Female	Male	Female
Infected untreated	10.75±1.24	7.11±1.17	21.28±3.75	13.22±2.13	23.86±2.14	16.33±1.22	34.23±4.54	23.22±2.57
Haemin	8.56±1.23 (20.37)	5.49±0.66 (22.78)	19.75±2.36 (7.19)	11.51±2.18 (12.94)	21.44±3.24 (10.14)	14.75±1.62 (9.68)	31.65±4.12 (7.54)	21.13±1.84 (9.00)
Art	6.33±0.95 ^a (41.12)	3.18±0.41 ^b (55.27)	17.45±2.33 (18)	10.21±1.06 (22.77)	16.33±1.73 ^a (31.56)	9.38±1.26 ^b (42.56)	29.26±3.11 (14.52)	18.64±2.11 (19.72)
Haemin + Art	3.66±0.42 ^{c, d} (65.95)	1.48±0.22 ^{c, e} (79.18)	15.91±2.34 (25.23)	8.92±1.41 (32.53)	11.65±1.83 ^b (51.17)	3.93±0.72 ^{c, e} (75.93)	27.93±3.72 (18.40)	16.57±2.63 (28.64)

Values are presented as mean ± SEM. Number of samples/group = 8 (each sample consisted of ten male or ten female worms). GPx was expressed by the formation of 1 μmole of NADPH/min/worm. Numbers between parentheses indicate the percentage of reduction from infected untreated group

^a $p<0.05$, ^b $p<0.01$, ^c $p<0.001$ (significant difference from infected untreated); ^d $p<0.05$, ^e $p<0.01$ (significant difference from Art-treated)