Timothy J. Langlois · Marti J. Anderson
Russell C. Babcock · Shin Kato

Marine reserves demonstrate trophic interactions across habitats

Published: 20 December 2005
© Springer-Verlag 2005

Oecologia (2005). DOI: 10.1007/s00442-005-0148-7

Figures 1 and 3 were inadvertently interchanged; their legends were not. These figures are given again here, each with its correct number and legend

Electronic Supplementary Material Additionally, a new version of the electronic supplementary material is provided at <http://dx.doi.org/10.1007/s00442-005-0237-7>

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00442-005-0148-3>

T. J. Langlois (✉)
Leigh Marine Laboratory, University of Auckland,
PO Box 349, Warkworth, New Zealand
E-mail: timothy.langlois@gmail.com
Tel.: +64-9-4226111
Fax: +64-9-4226113

M. J. Anderson
Department of Statistics, University of Auckland,
Private Bag 92019, Auckland, New Zealand

R. C. Babcock
CSIRO Marine Research Floreat, Private Bag No. 5,
Wembley, WA 6913, Australia

S. Kato
Jinbe-kan, 540-9 Kakinouchi, Iburi, Tosashimizu,
Kochi, Japan

Fig. 1 **a** Map of Hauraki Gulf and environs showing the location of the two reserves used in this study. **b** The insets show the reserve boundaries (dashed lines) and experimental sites (filled circles) at the two locations

Fig. 3 Proportion of dead *Dosinia subrosea* found at the experimental sites, within each reserve status, that exhibited marks consistent with predation by large lobster (Cracked), small lobster (Nibbled) or had no attributable marks (Ambiguous). Locations were pooled