Coevolution of foraging behavior with intrinsic growth rate: risk-taking in naturally and artificially selected growth genotypes of *Menidia menidia*

Susumu Chiba · Stephen A. Arnott · David O. Conover

Published online: 9 May 2008
© Springer-Verlag Berlin Heidelberg 2008

Erratum to: *Oecologia* (2007) 154:237–246 DOI 10.1007/s00442-007-0825-9

In the “Discussion” under the heading “Artificially selected genotypes” one paragraph was unfortunately omitted in the conversion process.

The missing paragraph is provided here:

There was a significant difference between treatments regarding the frequency with which fear-related behavior was expressed (Fisher’s exact test in 2×4 contingency table, $P = 0.002$). In the absence of food, fear-related behavior was observed in three of the seven NYFG groups

and six of the seven SC groups (Tukey’s WSD test, $P > 0.05$). However, in the presence of food, no NYFG groups expressed fear-related behavior, whereas six of the seven NYSG groups did (Tukey’s WSD test, $P < 0.01$).

In all seven of the NYFG trials where food was provided, every individual consumed food within 18 min (mean $\pm$ SD, 10.16 min $\pm$ 6.68) of reemergence. Feeding was also observed in all of the NYSG trials, although, in four cases, not all of the individuals accessed the food. There was a significant difference between genotypes in the proportion of individuals that fed (two-tailed t -test, $t = -2.63$, $P = 0.039$).

The online version of the original article can be found under doi:[10.1007/s00442-007-0825-9](https://doi.org/10.1007/s00442-007-0825-9).

S. Chiba · S. A. Arnott · D. O. Conover
Marine Sciences Research Center, Stony Brook University,
Stony Brook, NY 11794-5000, USA

Present Address:

S. Chiba (✉)
Department of Aquatic Bioscience,
Tokyo University of Agriculture, Hokkaido 099-2493, Japan
e-mail: s2chiba@bioindustry.nodai.ac.jp

Present Address:

S. A. Arnott
Beckman Center, B305, Stanford University,
Stanford, CA 94305, USA