Atle Mysterud · F. Javier Pérez-Barbería
Iain J. Gordon

The effect of season, sex and feeding style on home range area versus body mass scaling in temperate ruminants

Published online: 17 May 2001
© Springer-Verlag 2001

Oecologia (2001) 127(1):30–39

The first three sentences of the Introduction should read:

Considerable effort has been made to determine how the home range area (HR) of mammals scales with body mass (M) for a wide range of taxonomic groups ($HR=aM^b$; McNab 1963; Harestad and Bunnell 1979;

Damuth 1981; Mace and Harvey 1983; Lindstedt et al. 1986; Swihart 1986; Swihart et al. 1988; Gompper and Gittleman 1991). For energetic demands the scaling coefficient (b) is expected to be 0.75 (McNab 1963). However, most studies have recorded a slope steeper than this ($1 < b < 1.4$; e.g. Harestad and Bunnell 1979; Mace and Harvey 1983; Lindstedt et al. 1986).

The online version of the original article can be found at <http://dx.doi.org/10.1007/s004420000562>

A. Mysterud (✉)
Department of Biology, Division of Zoology, University of Oslo,
P.O. Box 1050 Blindern, 0316 Oslo, Norway
e-mail: atle.mysterud@bio.uio.no
Tel.: +47-22-854045, Fax: +47-22-854605

F.J. Pérez-Barbería · I.J. Gordon
The Macaulay Land Use Research Institute, Craigiebuckler,
Aberdeen, AB15 8QH, Scotland, UK

F.J. Pérez-Barbería
Instituto de Recursos Naturales y Ordenación del Territorio,
13 Independencia, Oviedo, 33071, Spain