Elementary Moves on Triangulations*

Monika Ludwig and Matthias Reitzner

Institut für Diskrete Mathematik und Geometrie, Technische Universität Wien,
Wiedner Hauptstraße 8-10/104, 1040 Wien, Austria
{monika.ludwig,matthias.reitzner}@tuwien.ac.at

Abstract. It is proved that a triangulation of a polyhedron can always be transformed into any other triangulation of the polyhedron using only elementary moves. One consequence is that an additive function (valuation) defined only on simplices may always be extended to an additive function on all polyhedra.

Introduction and Statement of Results

An n -polyhedron P in $\mathbb{R}^N$, $1 \leq n \leq N$, is a finite union of n -dimensional polytopes, where a polytope is the compact convex hull of finitely many points in $\mathbb{R}^N$. A finite set of n -simplices αP is a *triangulation* of P if no pair of simplices intersects in a set of dimension n and their union equals P . We investigate transformations of triangulations by elementary moves. Here an *elementary move* applied to αP is one of the two following operations: a simplex $T \in \alpha P$ is dissected into two n -simplices T_1, T_2 by a hyperplane containing an $(n - 2)$ -dimensional face of T ; or the reverse, that is, two simplices $T_1, T_2 \in \alpha P$ are replaced by $T = T_1 \cup T_2$ if T is again a simplex. We say that triangulations αP and βP are equivalent by elementary moves, and write $\alpha P \sim \beta P$, if there are finitely many elementary moves that transform αP into βP . The main object of this note is to show the following result.

Theorem 1. *If αP and βP are triangulations of the n -polyhedron P , then $\alpha P \sim \beta P$.*

A triangulation αP with the additional property that any pair of simplices intersects in a common face gives rise to a simplicial complex $\hat{\alpha} P$. It is a classical result of algebraic topology due to Alexander [5] and Newman [26], [27] (see also [20]) that a simplicial

* The research of both authors was supported in part by the European Network PHD, MCRN-511953.

complex $\hat{\alpha}P$ can always be transformed into any other simplicial complex $\hat{\beta}P$ with the same underlying polyhedron by using only finitely many stellar moves. Here a *stellar move* is a suitable sequence of elementary moves followed by a simplicial isomorphism, where a *simplicial isomorphism* between two complexes is a bijection between their vertices that induces a bijection between their k -dimensional simplices for $1 \leq k \leq n$. For precise definitions, see [20] and for related results, see [10], [28], and [29]. The new feature of Theorem 1 is that simplicial isomorphisms are not allowed. So our theorem belongs to metric geometry whereas the Alexander–Newman theorem is a topological result.

As an application of Theorem 1 we obtain the following results on valuations. Here a function $\mu: \mathcal{S} \rightarrow \mathbb{R}$ defined on a class $\mathcal{S}$ of sets is called a *valuation* or *additive* if $\mu(\emptyset) = 0$, where $\emptyset$ is the empty set, and if

$$\mu(S) + \mu(T) = \mu(S \cup T) + \mu(S \cap T),$$

for all $S, T \in \mathcal{S}$ such that $S \cup T, S \cap T \in \mathcal{S}$ as well.

Let $\mathcal{T}^n$ be the set of simplices of dimension at most n in $\mathbb{R}^N$. Note the following connection between the definition of valuations on simplices and elementary moves. If $S, T \in \mathcal{T}^n$, then $S \cup T \in \mathcal{T}^n$ implies that S and T can be obtained from $S \cup T$ by elementary moves and the most basic case is when $S \cup T$ is dissected by an elementary move into S and T . Let $\mathcal{Q}^n$ be the set of polyhedra of dimension at most n in $\mathbb{R}^N$.

Theorem 2. *Every valuation on $\mathcal{T}^n$ has a unique extension to a valuation on $\mathcal{Q}^n$.*

As a corollary we obtain the analogous theorem for polytopes. Let $\mathcal{P}^n$ be the set of polytopes of dimension at most n in $\mathbb{R}^N$.

Corollary 3. *Every valuation on $\mathcal{T}^n$ has a unique extension to a valuation on $\mathcal{P}^n$.*

A version of Corollary 3 is stated and used in [19]. Note that in the proof of this result in [19] the Alexander–Newman theorem has to be replaced by Theorem 1 of the present paper.

Valuations on polyhedra are classical and they played a critical role in Dehn’s solution of Hilbert’s Third Problem. Results regarding the classification and characterization of invariant valuations are central to convex and integral geometry; see [12], [18], [24], and [25]. In recent years, many new results on valuations have been obtained, see, for example, [1]–[4], [14]–[17], [21]–[23], and [34]. Also extension questions for general valuations are classical. Volland [37] and Perles and Sallee [30] proved that every valuation on polytopes has a unique extension to a valuation on polyhedra. Their results were generalized by Groemer [11]. Volume is the most basic example of a valuation on polyhedra. The question of how to extend volume from simplices to polyhedra is closely connected with the quest for an elementary definition of volume for polytopes. In his Third Problem, Hilbert [13] asked whether volume on polyhedra can be defined by using only scissors congruences, that is, if two polyhedra P and Q of the same volume can each be cut into a finite number of pieces $P_1, \dots, P_m$ and $Q_1, \dots, Q_m$ with P_i congruent to Q_i by a rigid motion for each i ? This question was answered in the negative by Dehn

[8] (see also [7] and [31]). However, there are simple definitions of volume for simplices. So the volume of n -simplices can be defined as height times the $(n - 1)$ -dimensional volume of its base divided by n . A simple geometric argument (see [35]) shows that this does not depend on the choice of the base and volume defined in this way is a valuation on simplices. Schatunovsky [33] (for $n = 3$) and Süss [35] (for general dimensions) proved that there is a unique extension of this valuation from simplices to polyhedra. In that way they obtained an elementary definition of volume for polyhedra. Theorem 2 is the extension of their result to general valuations.

Theorem 1 has already been used by a number of authors. In fact, results more general than Theorem 1 have already been used. For example, Lemma 2.2 in Sah's book on Hilbert's Third Problem [31] states that any two triangulations have a common refinement by elementary moves using only dissections. However, this is well known to be an important open problem in algebraic topology, see [20]. Later, Sah [32] replaced his lemma by Theorem 1 of the present paper. While Sah observes that Theorem 1 suffices for the constructions in his book, he neglects to provide a proof of Theorem 1. For related results, see also [9].

1. Proof of Theorem 1

A triangulation αP of an n -polytope P is called a *starring* at $a \in P$, if every n -simplex in αP has a vertex at a . Note that every n -polytope P has a starring at every $a \in P$. This can be seen by using induction on n . It is trivial for $n = 1$. Suppose that every $(n - 1)$ -polytope has a starring. Let P be an n -polytope and $a \in P$. For every $(n - 1)$ -dimensional face F_j of P with $a \notin F_j$, we choose a starring $\alpha_j F_j$. Then the convex hulls of a and the $(n - 1)$ -simplices in $\alpha_j F_j$ are a starring of P at a .

A triangulation γP of an n -polyhedron P is a *refinement* of αP if every simplex of γP is contained in a simplex of αP . Note that any two triangulations αP and βP have a common refinement. To see this, let $P_1, \dots, P_l$ be the polytopes $S \cap T$, $S \in \alpha P$, $T \in \beta P$, that are n -dimensional. For $j = 1, \dots, l$, we choose a triangulation $\gamma_j P_j$ of P_j , for example, we can take a starring of P_j at any point $a_j \in P_j$. Then $\gamma P = \gamma_1 P_1 \cup \dots \cup \gamma_l P_l$ is a triangulation of P . Since every simplex in γP is contained in a suitable P_j , γP is a common refinement of αP and βP .

Let P be an n -polyhedron. To show that $\alpha P \sim \beta P$ for any two triangulations αP and βP of P , we prove that any refinement of a given triangulation can be obtained from the original triangulation by finitely many elementary moves. To show this, it is enough to prove the following proposition.

Proposition. *If αT is a triangulation of the n -simplex T , then $\alpha T \sim T$.*

Here we write T (instead of $\{T\}$) for the trivial triangulation of T .

The rest of this section is devoted to the proof of this proposition. We follow the classical approach of Alexander and Newman as presented by Lickorish [20]. The main new feature of the proof is Lemma 2.

We use induction on the dimension n . The case $n = 1$ is trivial. Assume that the proposition is true for dimensions less than or equal to n , that is, for every triangulation

αS of a simplex S ,

$$\alpha S \sim S \quad \text{for} \quad \dim S < n, \quad (1)$$

where $\dim$ stands for dimension.

Note that if S is a simplex, then every elementary subdivision of S induces an elementary subdivision of the simplices in the boundary of S . The following observation is used several times. Let $P = [Q, v]$ be a pyramid with apex v and base Q . Let $\alpha Q = \{S_1, \dots, S_k\}$ and $\beta Q = \{T_1, \dots, T_l\}$ be triangulations of the $(n-1)$ -polytope Q . Let $[\alpha Q, v] = \{[S_1, v], \dots, [S_k, v]\}$ and $[\beta Q, v] = \{[T_1, v], \dots, [T_l, v]\}$. Then $[\alpha Q, v]$ and $[\beta Q, v]$ are triangulations of P and

$$\alpha Q \sim \beta Q \quad \Rightarrow \quad [\alpha Q, v] \sim [\beta Q, v]. \quad (2)$$

Lemma 1. *If αT is a starring of an n -simplex T , then $\alpha T \sim T$.*

Proof. Assume $T \subset \mathbb{R}^n$. First, let T be starred at $a \in \partial T$. Let $S_0, \dots, S_n$ be the facets of T and let $a \in S_0$. Then

$$\alpha T = [\alpha_1 S_1, a] \cup \dots \cup [\alpha_n S_n, a],$$

where $\alpha_i S_i$ is a triangulation of S_i . The induction hypothesis (1) and (2) imply that $[\alpha_i S_i, a] \sim [S_i, a]$. Thus,

$$\alpha T \sim [S_1, a] \cup \dots \cup [S_n, a].$$

Let v be the vertex of T opposite to S_0 . Then there is a starring $\alpha_0 S_0$ of S_0 at a such that

$$[S_1, a] \cup \dots \cup [S_n, a] = [\alpha_0 S_0, v].$$

The induction hypothesis (1) and (2) imply that $[\alpha_0 S_0, v] \sim [S_0, v]$. Consequently,

$$\alpha T \sim T \quad \text{for} \quad a \in \partial T. \quad (3)$$

Now, let a be a point in the interior of T . Then

$$\alpha T = [\alpha_0 S_0, a] \cup \dots \cup [\alpha_n S_n, a],$$

where $\alpha_i S_i$ is a triangulation of the facet S_i of T . By (1) and (2),

$$\alpha T \sim [S_0, a] \cup \dots \cup [S_n, a].$$

We write aT for the starring at a of the n -simplex T if every simplex in aT has a facet of T as its base. We dissect T by a hyperplane H through a and an $(n-2)$ -dimensional face of T into two simplices T^+, T^- . This is an elementary move. Thus by (3),

$$T \sim T^+ \cup T^- \sim aT^+ \cup aT^-.$$

If a facet S is subdivided by H into S^+ and S^- , then $[S^+, a] \cup [S^-, a] \sim [S, a]$. Thus

$$aT^+ \cup aT^- \sim aT.$$

This completes the proof of the lemma. $\square$

Note that, for $a \in P$ fixed, (1) and (2) imply that any two starrings at a are equivalent by elementary moves. We write aP for a starring of a polytope P at a point $a \in P$.

Lemma 2. *For every n -polytope P and $a, b \in P$, $aP \sim bP$.*

Proof. Assume $P \subset \mathbb{R}^n$. We use induction on the number m of vertices of P . If $m = n + 1$, then P is an n -simplex and the statement is true by Lemma 1. Suppose the lemma is true for polytopes with at most m vertices. Write $[A_1, \dots, A_l]$ for the convex hull of sets $A_1, \dots, A_l$.

Let P be a polytope with vertices $v_1, \dots, v_m, v$. Let $a \in P^- = [v_1, \dots, v_m]$. We say that a facet F of P^- is visible from v if for every $x \in F$, $[v, x] \cap P^- = \{x\}$. By starring P^- at a , the facets of P^- that are visible from v are subdivided into $(n - 1)$ -simplices S_i , $i = 1, \dots, l$. Note that aP^- contains the simplices $[S_i, a]$, $i = 1, \dots, l$, that are n -dimensional. Let $\mathcal{V}_a$ be the set of the n -simplices $[S_i, v]$, $i = 1, \dots, l$.

The main step is to show that

$$aP \sim aP^- \cup \mathcal{V}_a. \quad (4)$$

If (4) holds and if $b \in P^-$, then the induction hypothesis on the number of vertices implies that $aP^- \sim bP^-$. By (1) and (2) we obtain $\mathcal{V}_a \sim \mathcal{V}_b$. Thus

$$aP \sim aP^- \cup \mathcal{V}_a \sim bP^- \cup \mathcal{V}_b \sim bP.$$

If there is no vertex v such that $a, b \in [v_1, \dots, v_m]$, then there are vertices v, w of P such that $a \in [v_1, \dots, v_{m-1}, w]$, $b \in [v_1, \dots, v_{m-1}, v]$, and $P = [v_1, \dots, v_{m-1}, v, w]$. Choose $c \in [v_1, \dots, v_{m-1}, w] \cap [v_1, \dots, v_{m-1}, v]$ in the interior of P . Then by the above argument

$$aP \sim cP \quad \text{and} \quad cP \sim bP.$$

Thus it is enough to show (4) to prove the lemma.

Let H_j , $j = 1, \dots, k$, be the affine hulls of S_i , $i = 1, \dots, l$. Denote the intersection point of the hyperplane H_j with the segment $[a, v]$ by x_j (these points are not necessarily distinct). Without loss of generality assume that the hyperplanes H_j are numbered such that $[a, v]$ is dissected into $[a, x_1]$, $[x_1, x_2]$, $\dots$, $[x_{k-1}, x_k]$, $[x_k, x_{k+1} = v]$. Let $\mathcal{A}_1$ be the set of those simplices $[S_i, a]$, $i = 1, \dots, l$, that are n -dimensional. Set $\mathcal{V}_1 = \mathcal{V}_a$.

First, assume that $a \neq x_1$ and $x_k \neq v$. Starting with $j = 1$, we construct $\mathcal{A}_{j+1}$ and $\mathcal{V}_{j+1}$ from $\mathcal{A}_j$ and $\mathcal{V}_j$ in the following way. The simplices in $\mathcal{A}_j$ have a as a vertex. Let $\mathcal{A}_j(H_j) \subset \mathcal{A}_j$ be the subset of those simplices that have their $(n - 1)$ -dimensional base in H_j . The simplices in $\mathcal{V}_j$ have v as a vertex. Let $\mathcal{V}_j(H_j) \subset \mathcal{V}_j$ be the subset of those simplices that have their $(n - 1)$ -dimensional base in H_j . Note that the set of these $(n - 1)$ -dimensional bases coincides for $\mathcal{A}_j(H_j)$ and $\mathcal{V}_j(H_j)$. These bases form a facet F , say, of $[P^-, x_j]$. By the induction hypothesis, the triangulation of F by these $(n - 1)$ -dimensional bases is equivalent by elementary moves to a starring of F at x_j . The moves applied in H_j induce moves on $\mathcal{A}_j(H_j)$ and $\mathcal{V}_j(H_j)$. Denote the sets obtained by these moves by $\mathcal{A}'_j(H_j)$ and $\mathcal{V}'_j(H_j)$. Note that the set of $(n - 1)$ -dimensional bases coincides for $\mathcal{A}'_j(H_j)$ and $\mathcal{V}'_j(H_j)$. We replace $\mathcal{A}_j(H_j)$ by $\mathcal{A}'_j(H_j)$ and $\mathcal{V}_j(H_j)$ by $\mathcal{V}'_j(H_j)$ and set $\mathcal{A}'_j = (\mathcal{A}_j \setminus \mathcal{A}_j(H_j)) \cup \mathcal{A}'_j(H_j)$ and $\mathcal{V}'_j = (\mathcal{V}_j \setminus \mathcal{V}_j(H_j)) \cup \mathcal{V}'_j(H_j)$.

In the next step we do the following. If $x_j = x_{j+1}$, we set $\mathcal{A}_{j+1} = \mathcal{A}'_j$ and $\mathcal{V}_{j+1} = \mathcal{V}'_j$ and have

$$\mathcal{A}_j \cup \mathcal{V}_j \sim \mathcal{A}_{j+1} \cup \mathcal{V}_{j+1}.$$

If $x_j \neq x_{j+1}$, denote by $\mathcal{A}'_j(x_j) \subset \mathcal{A}'_j$ the subset of those simplices which contain x_j as a vertex and by $\mathcal{V}'_j(x_j) \subset \mathcal{V}'_j$ the subset of those simplices which contain x_j as a vertex. Each simplex in $\mathcal{V}'_j(x_j)$ is the convex hull of x_j, v , and an $(n-2)$ -dimensional face B , say. The convex hull $[B, x_j, v]$ is in $\mathcal{V}'_j(x_j)$ and $[B, a, x_j]$ in $\mathcal{A}'_j(x_j)$. We subdivide the simplex $[B, x_j, v]$ into the simplices $[B, x_j, x_{j+1}]$ and $[B, x_{j+1}, v]$. Then we take the union of $[B, a, x_j]$ and $[B, x_j, x_{j+1}]$ and obtain the simplex $[B, a, x_{j+1}]$. These operations are elementary moves. We do this with every simplex with vertex x_j . We obtain the new set $\mathcal{A}'_j(x_{j+1})$ which contains the simplices $[B, a, x_{j+1}]$ and the new set $\mathcal{V}'_j(x_{j+1})$ which contains the simplices $[B, x_{j+1}, v]$. Let $\mathcal{A}_{j+1} = (\mathcal{A}'_j \setminus \mathcal{A}'_j(x_j)) \cup \mathcal{A}'_j(x_{j+1})$ for $j \leq k$ and $\mathcal{V}_{j+1} = (\mathcal{V}'_j \setminus \mathcal{V}'_j(x_j)) \cup \mathcal{V}'_j(x_{j+1})$ for $j < k$ and $\mathcal{V}_{k+1} = \emptyset$. We have

$$\mathcal{A}_j \cup \mathcal{V}_j \sim \mathcal{A}_{j+1} \cup \mathcal{V}_{j+1}.$$

Next, we consider the case $a = x_1 = \dots = x_i$. Note that each facet of P^- that contains a is already starred at a . Let $\mathcal{V}_1(a) \subset \mathcal{V}_1$ be the subset of those simplices which contain a as a vertex. Each simplex in $\mathcal{V}_1(a)$ is the convex hull of a, v , and an $(n-2)$ -dimensional face B . We subdivide the simplex $[B, a, v]$ into the simplices $[B, a, x_{i+1}]$ and $[B, x_{i+1}, v]$. These operations are elementary moves. We do this with every simplex with vertex a . We obtain the new set $\mathcal{A}_{i+1}$ which contains the simplices $[B, a, x_{i+1}]$ and the new set $\mathcal{V}_{i+1}$ which contains the simplices $[B, x_{i+1}, v]$ and the simplices in $\mathcal{V}_1 \setminus \mathcal{V}_1(a)$. We have $\mathcal{V}_1 \sim \mathcal{A}_{i+1} \cup \mathcal{V}_{i+1}$. Starting with $j = i+1$, we construct $\mathcal{A}_{j+1}$ and $\mathcal{V}_{j+1}$ by the algorithm described above. If $x_{i'} = \dots = x_k = v$, then the above algorithm implies that $\mathcal{V}_{i'}$ contains only lower-dimensional simplices. Therefore we set $\mathcal{V}_{i'} = \emptyset, \dots, \mathcal{V}_{k+1} = \emptyset$ and $\mathcal{A}_{i'+1} = \mathcal{A}_{i'}, \dots, \mathcal{A}_{k+1} = \mathcal{A}_{i'}$.

Thus in all cases

$$\mathcal{A}_1 \cup \mathcal{V}_1 \sim \dots \sim \mathcal{A}_{k+1} \cup \mathcal{V}_{k+1} = \mathcal{A}_{k+1}.$$

Since $aP \setminus \mathcal{A}_{k+1} = aP^- \setminus \mathcal{A}_1$, this proves (4). $\square$

Let T be an n -simplex and let αT be a triangulation of T . Assume $T \subset \mathbb{R}^n$. Let $H_1, \dots, H_l$ be the affine hulls of the facets of the simplices in αT . Let $\zeta_j T$ be the dissection into polytopes (in general not simplices) of T by the hyperplanes $H_1, \dots, H_j$ and let $\zeta_0 T = T$. We dissect each polytope P of $\zeta_j T$ into simplices by starring P at an arbitrary interior point. Let $\beta_j T$ be a triangulation of T obtained from $\zeta_j T$ in this way.

We use induction on the number k of hyperplanes. By Lemma 1, we have $T \sim \beta_0 T$. For $k \geq 1$ assume that

$$T \sim \beta_j T \quad \text{for } j < k. \quad (5)$$

Note that we obtain $\zeta_k T$ from $\zeta_{k-1} T$ by cutting by H_k . Every cell of $\zeta_{k-1} T$ is either unchanged or it is cut into two pieces. So let $P \in \zeta_{k-1} T$ be cut into pieces P_1, P_2 . We show that for $a \in P$ and $a_i \in P_i, i = 1, 2$,

$$aP \sim a_1 P_1 \cup a_2 P_2. \quad (6)$$

By Lemma 2, $aP \sim bP$ for every $b \in H_k \cap P$. Again by Lemma 2, $a_1P_1 \sim bP_1$ and $a_2P_2 \sim bP$. Since any two starrings of P at $a \in P$ are equivalent by elementary moves this implies (6). By our definition of $\beta_{k-1}T$ and β_kT , (6) implies that

$$\beta_{k-1}T \sim \beta_kT.$$

Thus, by induction,

$$T \sim \beta_0T \sim \dots \sim \beta_lT. \quad (7)$$

Let αT consist of the simplices $S_1, \dots, S_m$. Let $\zeta_j S_i$ be the dissection into polytopes of S_i by the hyperplanes $H_1, \dots, H_j$. Let $\beta_j S_i$ be a triangulation that is obtained by starring each cell of $\zeta_j S_i$ at a point in that cell. By Lemma 2 these triangulations for different starring points are equivalent. Thus we obtain as before that

$$S_i \sim \beta_0 S_i \sim \dots \sim \beta_l S_i.$$

Consequently,

$$\alpha T = (S_1 \cup \dots \cup S_m) \sim \dots \sim (\beta_l S_1 \cup \dots \cup \beta_l S_m) = \beta_l T.$$

Combined with (7) this completes the proof of the proposition.

2. Proof of Theorem 2

We use induction on n . The case $n = 0$ is straightforward. Suppose that every valuation on $\mathcal{T}^{n-1}$ has a unique extension to a valuation on $\mathcal{Q}^{n-1}$. Set $\mathcal{R}^n = \{T \cup Q \mid T \in \mathcal{T}^n, Q \in \mathcal{Q}^{n-1}\}$. By the induction hypothesis, it is straightforward that μ has a unique extension from $\mathcal{T}^n$ to $\mathcal{R}^n$. We prove that μ has a unique extension from $\mathcal{R}^n$ to a valuation on $\mathcal{Q}^n$.

Let $Q \in \mathcal{Q}^n$ and let $\delta Q = \{R_1, \dots, R_k\}$, $R_i \in \mathcal{R}^n$, be a dissection of Q ; that is, the intersection of a pair of elements of δQ has dimension less than n and their union equals Q . For an ordered j -tuple $I = \{i_1, \dots, i_j\}$, $1 \leq i_1 < \dots < i_j \leq k$, $1 \leq j \leq k$, set $R_I = R_{i_1} \cap \dots \cap R_{i_j}$ and $|I| = j$. The set $\mathcal{Q}^n$ is a lattice. Thus, if μ can be extended to a valuation on $\mathcal{Q}^n$, then by the inclusion-exclusion principle

$$\mu(Q) = \sum_{I \subset \{1, \dots, k\}} (-1)^{|I|-1} \mu(R_I). \quad (8)$$

We show that if $Q \in \mathcal{Q}^n$ and if $\delta Q = \{R_1, \dots, R_k\}$, $R_i \in \mathcal{R}^n$, is a dissection of Q , then (8) can be used as a definition of $\mu(Q)$. Note that $R_I \in \mathcal{R}^n$. Therefore the induction hypothesis implies that the right side of (8) is well defined.

We show that $\mu(Q)$ as defined by (8) does not depend on the choice of δQ . First, let Q be an n -polyhedron and let $\alpha Q = \{T_1, \dots, T_k\}$ be a triangulation of Q . By Theorem 1, all triangulations of an n -polyhedron are equivalent by elementary moves. Thus it is sufficient to show that applying an elementary move to αQ does not change $\mu(Q)$. Since an elementary move is either the dissection of a simplex or the reverse, it suffices to show the following. If $S_i = T_i$, $i = 1, \dots, k-1$, and if T_k is subdivided into S_k and S_{k+1} , then

$$\sum_{I \subset \{1, \dots, k\}} (-1)^{|I|-1} \mu(T_I) = \sum_{J \subset \{1, \dots, k+1\}} (-1)^{|J|-1} \mu(S_J). \quad (9)$$

Set $I' = I \setminus \{k\}$. Since μ is a valuation on $\mathcal{R}^n$, we have for an ordered j -tuple I with $k \in I$,

$$\begin{aligned}\mu(T_I) &= \mu(T_{I'} \cap T_k) = \mu(S_{I'} \cap (S_k \cup S_{k+1})) = \mu((S_{I'} \cap S_k) \cup (S_{I'} \cap S_{k+1})) \\ &= \mu(S_{I'} \cap S_k) + \mu(S_{I'} \cap S_{k+1}) - \mu(S_{I'} \cap S_k \cap S_{k+1}) \\ &= \mu(S_I) + \mu(S_{I'} \cap S_{k+1}) - \mu(S_I \cap S_{k+1}).\end{aligned}$$

It follows that

$$\begin{aligned}\sum_J (-1)^{|J|-1} \mu(S_J) &= \sum_{k, k+1 \notin J} (-1)^{|J|-1} \mu(S_J) + \sum_{k \in J, k+1 \notin J} (-1)^{|J|-1} \mu(S_J) \\ &\quad + \sum_{k \notin J, k+1 \in J} (-1)^{|J|-1} \mu(S_J) + \sum_{k, k+1 \in J} (-1)^{|J|-1} \mu(S_J) \\ &= \sum_{k \notin I} (-1)^{|I|-1} \mu(S_I) + \sum_{k \in I} (-1)^{|I|-1} \mu(S_I) \\ &\quad + \sum_{k \in I} (-1)^{|I|-1} \mu(S_{I'} \cap S_{k+1}) - \sum_{k \in I} (-1)^{|I|-1} \mu(S_I \cap S_{k+1}) \\ &= \sum_{k \notin I} (-1)^{|I|-1} \mu(T_I) + \sum_{k \in I} (-1)^{|I|-1} \mu(T_I) \\ &= \sum_I (-1)^{|I|-1} \mu(T_I).\end{aligned}$$

Thus (9) holds and there is a unique extension of μ to the set of n -polyhedra.

Next, let $P \in \mathcal{Q}^n$ be decomposed into Q and R , where Q is a uniquely defined n -polyhedron (or the empty set) and $R \in \mathcal{Q}^{n-1}$. Let $\delta P = \alpha Q \cup R$, where $\alpha Q = \{R_1, \dots, R_{k-1}\}$ is a triangulation of Q , and set $R_k = R$. This is a dissection of P . By (8),

$$\begin{aligned}\mu(P) &= \sum_I (-1)^{|I|-1} \mu(R_I) = \sum_{k \notin I} (-1)^{|I|-1} \mu(R_I) + \sum_{k \in I} (-1)^{|I|-1} \mu(R_I) \\ &= \mu(Q) + \mu(R) - \mu(Q \cap R).\end{aligned}$$

Since Q is an n -polyhedron and since $R, Q \cap R \in \mathcal{Q}^{n-1}$, the terms $\mu(Q)$, $\mu(R)$, and $\mu(Q \cap R)$ are well defined. Hence $\mu(P)$ does not depend on δP either.

Finally, we show that μ as defined by (8) is a valuation on $\mathcal{Q}^n$. Let $P, Q \in \mathcal{Q}^n$. We choose a dissection $\{R_1, \dots, R_m\}$ of $P \cup Q$ such that, for every R_i , we have $R_i \subset P$ or $R_i \subset Q$. Then for every ordered j -tuple I ,

$$\mu(P \cap R_I) + \mu(Q \cap R_I) = \mu((P \cup Q) \cap R_I) + \mu((P \cap Q) \cap R_I).$$

By (8), it follows that μ is a valuation.

3. Proof of Corollary 3

The proof presented here relies essentially on a theorem of Tverberg [36]. Note that it is also possible to prove the corollary by the extension theorems of Volland [37] or Perles and Sallee [30] or Groemer [11].

The main step is to prove that there is at most one extension. Let μ_1 and μ_2 be valuations on $\mathcal{P}^n$ such that $\mu_1 = \mu_2$ on $\mathcal{T}^n$. Using induction on n we show that $\mu_1 = \mu_2$ also on $\mathcal{P}^n$. The cases $n = 0, 1$ are trivial. Suppose that $\mu_1 = \mu_2$ on $\mathcal{P}^{n-1}$.

A *binary space partition* is formed by partitioning $\mathbb{R}^N$ by a hyperplane H into two closed halfspaces H^+ , H^- , and then recursively partitioning each of the two resulting halfspaces; the result is a hierarchical decomposition of space into closed convex cells (see [6]). Let P be an n -polytope. Since μ_1 and μ_2 are valuations, after the first step of a binary space partition we have for $i = 1, 2$,

$$\mu_i(P) = \mu_i(P \cap H^+) + \mu_i(P \cap H^-) - \mu_i(P \cap H).$$

Note that $P \cap H \in \mathcal{P}^{n-1}$ for $P \not\subset H$ and thus $\mu_1(P \cap H) = \mu_2(P \cap H)$. Hence to prove $\mu_1(P) = \mu_2(P)$ it suffices to prove $\mu_1(Q) = \mu_2(Q)$ for Q equal to $P \cap H^+$ and $P \cap H^-$. In the next step of the binary space partition the convex polytopes $P \cap H^+$ and $P \cap H^-$ are dissected by suitable hyperplanes and the same argument applies, etc.

Tverberg [36] showed that, given an n -polytope P , one can find a binary space partition that decomposes P in finitely many steps into n -simplices. Since $\mu_1 = \mu_2$ on $\mathcal{T}^n$, this implies $\mu_1(P) = \mu_2(P)$ and the uniqueness is established.

By Theorem 2 there is at least one extension of μ to a valuation on $\mathcal{P}^n$, the one which can be further extended to $\mathcal{Q}^n$. Combined with the uniqueness result this proves the corollary.

Acknowledgments

The authors thank Professors G. Kuperberg, W. B. R. Lickorish, and P. McMullen for help in clarifying which statements and proofs are available in the literature and Professors P. M. Gruber and E. Lutwak and the anonymous referees for their helpful comments.

References

1. S. Alesker, Continuous rotation invariant valuations on convex sets, *Ann. of Math. (2)* **149** (1999), 977–1005.
2. S. Alesker, Description of translation invariant valuations on convex sets with solution of P. McMullen's conjecture, *Geom. Funct. Anal.* **11** (2001), 244–272.
3. S. Alesker, Hard Lefschetz theorem for valuations, complex integral geometry, and unitarily invariant valuations, *J. Differential Geom.* **63** (2003), 63–95.
4. S. Alesker, The multiplicative structure on continuous polynomial valuations, *Geom. Funct. Anal.* **14** (2004), 1–26.
5. J. Alexander, The combinatorial theory of complexes, *Ann. of Math. (2)* **31** (1930), 292–320.
6. M. de Berg, M. van Kreveld, M. Overmars, and O. Schwarzkopf, *Computational Geometry*, Springer-Verlag, Berlin, 2000.
7. V. Boltianskii, *Hilbert's Third Problem*, Wiley, New York, 1978.
8. M. Dehn, Über den Rauminhalt, *Math. Ann.* **55** (1901), 465–478.
9. J. L. Dupont, Algebra of polytopes and homology of flag complexes, *Osaka J. Math.* **19** (1982), 599–641.
10. G. Ewald and G. C. Shephard, Stellar subdivisions of boundary complexes of convex polytopes, *Math. Ann.* **210** (1974), 7–16.
11. H. Groemer, On the extension of additive functionals on classes of convex sets, *Pacific J. Math.* **75** (1978), 397–410.

12. H. Hadwiger, *Vorlesungen über Inhalt, Oberfläche und Isoperimetrie*, Springer, Berlin, 1957.
13. D. Hilbert, Mathematical Problems. Lecture delivered before the International Congress of Mathematicians in Paris, 1900. Translated by M. W. Newson, *Bull. Amer. Math. Soc.* **8** (1902), 437–479.
14. D. A. Klain, A short proof of Hadwiger's characterization theorem, *Mathematika* **42** (1995), 329–339.
15. D. A. Klain, Star valuations and dual mixed volumes, *Adv. Math.* **121** (1996), 80–101.
16. D. A. Klain, Invariant valuations on star-shaped sets, *Adv. Math.* **125** (1997), 95–113.
17. D. A. Klain, Even valuations on convex bodies, *Trans. Amer. Math. Soc.* **352** (2000), 71–93.
18. D. A. Klain and G.-C. Rota, *Introduction to Geometric Probability*, Cambridge University Press, Cambridge, 1997.
19. G. Kuperberg, A generalization of Filliman duality, *Proc. Amer. Math. Soc.* **131** (2003), 3893–3899.
20. W. B. R. Lickorish, Simplicial moves on complexes and manifolds, *Proceedings of the Kirbyfest (Berkeley, CA, 1998)*, Geometry of Topology Monographs, vol. 2, Geometry of Topology, Coventry, 1999, pp. 299–320.
21. M. Ludwig, Valuations on polytopes containing the origin in their interiors, *Adv. Math.* **170** (2002), 239–256.
22. M. Ludwig, Ellipsoids and matrix valued valuations, *Duke Math. J.* **119** (2003), 159–188.
23. M. Ludwig and M. Reitzner, A characterization of affine surface area, *Adv. Math.* **147** (1999), 138–172.
24. P. McMullen, Valuations and dissections, in *Handbook of Convex Geometry*, Vol. B (P. M. Gruber and J. M. Wills, eds.), North-Holland, Amsterdam, 1993, pp. 933–990.
25. P. McMullen and R. Schneider, Valuations on convex bodies, *Convexity and Its Applications* (P. M. Gruber and J. M. Wills, eds.), Birkhäuser, Basel, 1983, pp. 170–247.
26. M. H. A. Newman, On the foundations of combinatory analysis situs. I, II, *Proc. Amsterdam* **29** (1926), 611–626, 627–641.
27. M. H. A. Newman, A theorem in combinatory topology, *J. London Math. Soc.* **6** (1931), 186–192.
28. U. Pachner, Über die bistellare Äquivalenz simplizialer Sphären und Polytope, *Math. Z.* **176** (1981), 565–576.
29. U. Pachner, Shellings of simplicial balls and p.l. manifolds with boundary, *Discrete Math.* **81** (1990), 37–47.
30. M. A. Perles and G. T. Sallee, Cell complexes, valuations, and the Euler relation, *Canad. J. Math.* **22** (1970), 235–241.
31. C. H. Sah, *Hilbert's Third Problem: Scissors Congruence*, Research Notes in Mathematics, vol. 33, Pitman (Advanced Publishing Program), Boston, MA, 1979.
32. C. H. Sah, Scissors congruences. I. The Gauss–Bonnet map, *Math. Scand.* **49**(2) (1981), 181–210 (1982).
33. S. O. Schatunovsky, Über den Rauminhalt der Polyeder, *Math. Ann.* **57** (1903), 496–508.
34. R. Schneider, Simple valuations on convex bodies, *Mathematika* **43** (1996), 32–39.
35. W. Süss, Über das Inhaltsmass bei mehrdimensionalen Polyedern, *Tohoku Math. J.* **38** (1933), 252–261.
36. H. Tverberg, How to cut a convex polytope into simplices, *Geom. Dedicata* **3** (1974), 239–240.
37. W. Volland, Ein Fortsetzungssatz für additive Eipolyederfunktionale im euklidischen Raum, *Arch. Math.* **8** (1957), 144–149.

Received June 28, 2005, and in revised form December 28, 2005, and January 9, 2006.

Online publication April 18, 2006.