Affinely Regular Polygons as Extremals of Area Functionals

Paolo Gronchi · Marco Longinetti

Received: 5 August 2005 / Revised: 29 May 2006 /

Published online: 11 September 2007

© Springer Science+Business Media, LLC 2007

Abstract For any convex n -gon P we consider the polygons obtained by dropping a vertex or an edge of P . The area distance of P to such $(n - 1)$ -gons, divided by the area of P , is an affinely invariant functional on n -gons whose maximizers coincide with the affinely regular polygons. We provide a complete proof of this result.

We extend these area functionals to planar convex bodies and we present connections with the affine isoperimetric inequality and parallel X-ray tomography.

Keywords Affinely regular polygons · Geometric tomography · Affine length

1 Introduction

Given a convex polygon P with n vertices z_j ordered counterclockwise, we define $W_j(P)$ to be the triangle $z_{j-1}z_jz_{j+1}$ and $T_j(P)$ to be the (possibly unbounded) triangle outside P bounded by the side z_jz_{j+1} and the continuations of the two adjacent sides (see Fig. 1). Henceforth, the index j is taken modulo n , and $|C|$ denotes the area of C .

In this paper we consider the following affinely invariant functionals defined on the class $\mathcal{P}_n$ of planar convex n -gons, i.e., polygons with *exactly* n vertices:

P. Gronchi (✉)

Dipartimento di Matematica e Applicazioni per l'Architettura, Università degli Studi di Firenze,
Piazza Ghiberti 27, 50122 Firenze, Italy
e-mail: paolo@fi.iac.cnr.it

M. Longinetti

Dipartimento di Ingegneria Agraria e Forestale, Università degli Studi di Firenze, P. le delle
Cascine 15, 50139 Firenze, Italy
e-mail: marco.longinetti@unifi.it

Fig. 1 Triangles W_j and T_j of P

$$F(P) = \min_{j=1,\dots,n} \frac{|W_j(P)|}{|P|}, \quad (1)$$

$$G(P) = \min_{j=1,\dots,n} \frac{|T_j(P)|}{|P|}, \quad (2)$$

and we are interested in the maximizers of these functionals.

In Theorem 1.8 it is shown that *the maximizers of the above functionals are affinely regular n -gons, i.e., affine images of regular n -gons*. This class, denoted by $\mathcal{R}_n$, often appears in geometric problems with affine invariance [1, 3, 5, 17, 20].

A characterization of $\mathcal{R}_n$ as extremals of area functionals was obtained by Renyi and Sulanke [20]. They proved in Satz 2 that $\prod_{i=1}^n |W_i(P)|/|P|^n$ attains its maximum on $\mathcal{R}_n$.

The functional F was first introduced by Lopez and Reisner [17] in connection with algorithms for the approximation of a convex set by polygons. They showed that Theorem 1.8 for the functional F is a consequence of the result by Renyi and Sulanke.

The functional G was first introduced by Longinetti [15], where Theorem 1.8 is proved for $n = 5, 6$, via elementary geometric arguments. The functional G and a similar functional (not affinely invariant) considered in [14] are related to Hammer's X-ray problem for planar convex bodies proposed in [10]: How many X-ray pictures of a convex body must be taken in order to permit its reconstruction? The solution of this problem is given by Gardner and McMullen [6]. We refer to [7, Chap. 1] for an overview of this topic. In Sect. 6 we present in detail the connection between the functional G and the stability of the reconstruction in the Hammer's problem. In Sect. 5, we discuss some extensions of functionals F and G to the class of planar convex bodies related to the affine length of a convex body and to the affine isoperimetric inequality. These functionals are also related to the approximation of planar convex bodies by polygons [9, 17]. In particular, F is related to the approximation of an n -gon P by $(n-1)$ -gons contained in P . Similarly, G is related with the approximation of P by $(n-1)$ -gons containing P . Because of this we use the word *inner* or *outer* in connection with F or G , respectively. In higher dimension, similar function-

als involving polytopes obtained by dropping a vertex or a facet were investigated by Reisner et al. [19].

As a first remark we deal with the trivial cases $n = 3, 4$. For $n = 3$, we have $F(P) = 1$ and $G(P) = \infty$, for every P . For $n = 4$, by elementary arguments, one can prove that the maximizers of F have diagonals which divide them into triangles of equal area, and $G(P) = \infty$ only for parallelograms. Hence, all maximizers of F and G are parallelograms, and vice versa. As mentioned, some instances of Theorem 1.8 were already proved. In this paper we complete this result in a common framework to all n , for both the inner and outer functional.

We now provide a guide to the proof of Theorem 1.8. The functionals F and G are continuous with respect to the Hausdorff metric in $\mathcal{P}_n$, which is not compact, since n -gons can converge to polygons with fewer vertices. Hence, we have first to prove the existence of the maximizers in Lemma 2.1.

A first step towards the characterization of these maximizers is to show that they satisfy an inner (or outer) *equal-area property* which has some interest by itself. Consider the following classes:

$$\Phi_n = \{P \in \mathcal{P}_n : |W_1(P)| = |W_2(P)| = \cdots = |W_n(P)|\}, \quad (3)$$

$$\Gamma_n = \{P \in \mathcal{P}_n : |T_1(P)| = |T_2(P)| = \cdots = |T_n(P)|\}. \quad (4)$$

Henceforth, we say that a polygon P has the inner (outer) equal-area property when P belongs to Φ_n (Γ_n , respectively). Section 2.1 contains the proof of the following proposition.

Proposition 1.1 *F and G attain their maximum on Φ_n and Γ_n , respectively.*

It can be proved that the classes Φ_5 and Γ_5 coincide with the class of affinely regular pentagons. For $n > 5$, it is easy to see that the above class is larger than the class $\mathcal{R}_n$ of affinely regular polygons. For example, hexagons in Γ_6 are, up to an affine transformation, the intersection of two concentric equilateral triangles (see the shaded polygon in Fig. 2). A proof can be found in [15]. Similarly, hexagons in Φ_6 are, up to an affine transformation, equiangular (see the larger polygon in Fig. 2). In [11] the polygons of Φ_n are considered and a larger class of Φ_n containing not necessarily

Fig. 2 Polygons from Φ_6 and Γ_6

convex n -gons is explicitly parametrized by $n - 5$ real parameters modulo the action of the affine group.

Roughly speaking, (3) or (4) involve $n - 1$ independent constraints. Since n -gons depend on the $2n$ coordinates of its vertices, it follows that Φ_n and Γ_n depend on $n + 1$ parameters, and therefore, modulo the action of the affine group, depend on $n - 5$ parameters. So, in order to show that all maximizers of F and G are in $\mathcal{R}_n$ other significant properties must be proved. A subclass of Φ_{2m} was characterized by Bianchi and Longinetti [2, Lemma 1].

Let l_j be the length of the side $z_j z_{j+1}$ of P , i.e. $l_j = \|z_j - z_{j+1}\|$ and d_j the length of the diagonal $z_{j-1} z_{j+2}$.

Definition 1.2 We define $\mathcal{F}_n$ as the class of convex n -gons such that

$$\left(\frac{d_{j+1} - l_{j+1}}{d_{j+1}} \right) \left(\frac{d_j - l_j}{l_j} \right) = \left(\frac{d_{j-2} - l_{j-2}}{d_{j-2}} \right) \left(\frac{d_{j-1} - l_{j-1}}{l_{j-1}} \right) \quad \text{for } j = 1, \dots, n. \quad (5)$$

We say that P has the *inner-ratio property* when $P \in \mathcal{F}_n$. Assuming that all triangles T_j of P are bounded, we define v_j as the vertex of T_j not in P . Also, set $s_j = \|v_j - z_j\|$ and $p_j = \|z_{j+1} - v_j\|$. Let e_j be the length of the segment joining the outer points v_{j+1}, v_{j-1} . Notice that $e_j = p_{j-1} + l_j + s_{j+1}$.

Definition 1.3 We define $\mathcal{G}_n$ as the class of convex n -gons such that

$$\frac{s_j e_{j-1}}{l_{j-1}(p_{j-2} + l_{j-1})} = \frac{p_j e_{j+1}}{l_{j+1}(s_{j+2} + l_{j+1})} \quad \text{for } j = 1, \dots, n. \quad (6)$$

We call this the *outer-ratio property*. Notice that the outer-ratio property is an equality between ratios of lengths of segments on the two lines through v_j in Fig. 1.

The following theorems, proved in Sects. 2.1, 2.2, are important steps toward the goal.

Theorem 1.4 If P^* is a maximizer of F in $\mathcal{P}_n$, then $P^* \in \mathcal{F}_n$.

Theorem 1.5 If P^* is a maximizer of G in $\mathcal{P}_n$, then $P^* \in \mathcal{G}_n$.

In Sect. 3, through an algebraic manipulation, we will prove that in (5) the ratios $\lambda_j = d_j/l_j$ are independent of j . Analogously, for the outer problem, we will prove that in (6) the ratios $\zeta_j = s_j/l_{j-1} = p_{j-1}/l_j$ are independent of j . In Sect. 4 we go back to planar geometry and prove the following theorems.

Theorem 1.6

$$\Phi_n \cap \mathcal{F}_n = \mathcal{R}_n.$$

Theorem 1.7

$$\Gamma_n \cap \mathcal{G}_n = \mathcal{R}_n.$$

These results permit us to obtain the goal of the paper.

Theorem 1.8 *All maximizers of F or G on $\mathcal{P}_n$ are affinely regular polygons.*

2 First Variations at Extremal Polygons

Observe that for each $P \in \mathcal{P}_n$ the triangles $W_j(P)$ have positive area, hence $F(P) > 0$, but if P is close to a polygon with $n - 1$ vertices, then $F(P)$ is close to zero. Therefore $\inf_{P \in \mathcal{P}_n} F(P) = 0$ and F has no minimum in $\mathcal{P}_n$. Similarly, $\inf_{P \in \mathcal{P}_n} G(P) = 0$ and G has no minimum in $\mathcal{P}_n$. Then *all the extremals of the functionals F and G in $\mathcal{P}_n$ are maximizers*. In this section we obtain the more significant properties of these polygons. We use only elementary arguments of Euclidean geometry in the proof of Proposition 1.1. The idea of the proof is to consider a maximizer P^* of each functional and a suitable local variation P_ε^* of one or two vertices. An analysis of the sign of the area difference Δ_ε yields the results.

To prove directly Theorems 1.4 and 1.5, which are the principal goal of this section, a more complicated perturbation P_ε^* of P^* involving five consecutive vertices of P^* can be carried out. In this case one has to take into account only the first order terms of Δ_ε . This computation, using Proposition 1.1, can be explicitly obtained in terms of the sides and angles of P^* .

Here, we prefer to give a different proof, less geometric, via partial differentiation of area functionals (Lemmas 2.2 and 2.3) with respect to the vertices z_j . This yields the Lagrange multiplier systems (21) and (28) for the area $|P|$ under the corresponding equal-area property constraints. At the end of each subsection, an algebraic manipulation of such systems will give the proof of Theorem 1.4 for the inner problem and the proof of Theorem 1.5 for the outer one.

We begin with the following result.

Lemma 2.1 *The functionals F and G have maxima in $\mathcal{P}_n$.*

Proof From the well-known John theorem about the maximal ellipse contained in P , see [12], we can restrict to n -gons with fixed area 1, whose boundaries are contained in a circular annulus A of radii r and $2r$. Clearly $r \geq 1/\sqrt{4\pi}$ and the diameter of P is less or equal to $D = 4/\sqrt{\pi}$. Represent each polygon in $\mathcal{P}_n$ as a point in $\mathbb{R}^{2n}$ with coordinates the coordinates of its vertices. With respect to the standard metric on $\mathbb{R}^{2n}$, F is continuous and the class of k -gons, $k \leq n$, with vertices contained in A is compact. Since $F > 0$, it is trivial to show that a convergent maximizing sequence on $\mathcal{P}_n$ has vertices which converge to n distinct points, no three collinear.

Turning to G , some difficulties arise since $|T_j|$ may be infinite. Let α_j be the exterior angle of P at the vertex z_j . If $T_j(P)$ is bounded then

$$|T_j(P)| = \frac{1}{2} l_j^2 (\cot \alpha_{j+1} + \cot \alpha_j)^{-1}. \quad (7)$$

The function $\cot x$ is convex in $(0, \pi/2)$ and symmetric with respect to the point $\pi/2$. Thus

$$\cot\left(\frac{x_1 + x_2}{2}\right) \leq \frac{\cot x_1 + \cot x_2}{2} \quad \text{if } x_1, x_2 > 0 \text{ and } x_1 + x_2 < \pi. \quad (8)$$

Since $\alpha_j > 0$ and $\sum_{j=1}^n \alpha_j = 2\pi$, there exists j such that

$$\frac{\alpha_{j+i} + \alpha_j}{2} \leq \frac{2\pi}{n}.$$

Therefore, by (7) and (8) it follows that

$$\min_{j=1, \dots, n} |T_j(P)| < \frac{1}{4} D^2 \tan \frac{2\pi}{n}.$$

Hence, G is bounded from above on $\mathcal{P}_n$, for $n \geq 5$.

Now consider a maximizing sequence $\{P^m\}$ of n -gons with vertices in the circular annulus A , which converges to P^* . It remains to show that P^* has exactly n vertices. We can suppose

$$\min_{j=1, \dots, n} |T_j(P^m)| > \frac{1}{2} \sup_{P \in \mathcal{P}_n} G(P) = \mu > 0, \quad \text{for all } m. \quad (9)$$

For each P with vertices in A , consider a triangle similar to $T_j(P)$ bounded by the continuations of its sides not in P and a line parallel to the side $z_{j+1}z_j$ through the center of A . If h_j denotes the altitude of $T_j(P)$ to the side $z_{j+1}z_j$, then this larger triangle has an altitude smaller than $h_j + 2r$ and a base larger than $2r$. Hence

$$\frac{l_j}{h_j} \geq \frac{2r}{h_j + 2r},$$

i.e., $l_j \geq (2r - l_j)h_j/2r$. Since $l_j h_j = 2|T_j(P)| \geq 2\mu$, we deduce $l_j^2 \geq (2r - l_j)\mu/r$ and consequently that the sides of the polygons P^m are uniformly larger than a positive constant. This implies that n distinct points z_j^* of P^* are limits of the sequences of the vertices of P^m . Moreover, from (9), the limit of the area $|T_j(P^m)|$ is positive. Hence, no three consecutive points z_j^* are collinear and they are all distinct vertices of P^* . $\square$

Proof of Proposition 1.1 The claim that G attains its maximum on Γ_n was already proved in [15, Theorem 1].

For the inner case, assume that P^* is a maximizer of F . Let W_r be a triangle of maximal area among the n triangles W_j . Moving the vertex z_r towards the interior of P^* reduces the area of P^* . Since F cannot increase, the value $\min |W_j|$ has to decrease. Hence, either W_{r-1} or W_{r+1} is of minimal area. The freedom we have in choosing the direction along which z_r moves easily implies that they are both of minimal area. Therefore,

$$\begin{aligned} |z_{r-1}z_{r-2}z_{r-3}| &= |W_{r-2}| \geq |W_{r-1}| = |z_r z_{r-1} z_{r-2}|, \\ |z_{r+1}z_{r+2}z_{r+3}| &= |W_{r+2}| \geq |W_{r+1}| = |z_r z_{r+1} z_{r+2}|. \end{aligned}$$

Then, the distance of z_{r-3} from the line containing the edge $z_{r-2}z_{r-1}$ is larger or equal to that of z_r . Analogously, the distance of z_{r+3} from the line containing $z_{r+2}z_{r+1}$ is larger or equal to that of z_r . From the convexity of P^* , the triangle $U = z_{r-3}z_rz_{r+3}$ is ordered counterclockwise and contained in P^* . Therefore, any movement of z_r inside U increases the area of both W_{r-1} and W_{r+1} and decreases the area of P . The maximality of P^* implies that $|W_r|$ cannot be larger than $\min |W_j|$. $\square$

We represent with complex variable $x_j + iy_j$ the vertices z_j , i.e. z_j represents both a point in the plane and a complex number; so the area functionals $|T_j|$, $|W_j|$ and $|P|$ are real functions of complex variables z_j . We use partial derivatives of functions with respect to complex variables with the notation

$$f_{\bar{z}} = \frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right).$$

Lemma 2.2 *If T is the triangle with vertices a, b, c ordered counterclockwise, then*

$$\frac{4}{i} \frac{\partial |T|}{\partial \bar{a}} = c - b. \quad (10)$$

Proof If $a = x_a + iy_a$, $b = x_b + iy_b$, and $c = x_c + iy_c$ the result is obtained by an elementary computation starting from the formula

$$\begin{aligned} \frac{4}{i} |T| &= \frac{2}{i} ((x_c - x_b)(y_a - y_b) - (x_a - x_b)(y_c - y_b)) \\ &= (c - b)(\bar{a} - \bar{b}) - (\bar{c} - \bar{b})(a - b). \end{aligned} \quad (11)$$

$\square$

Lemma 2.3 *Let b, c, d and e be the vertices of a convex quadrilateral ordered clockwise. Let abc be the triangle T outside the quadrilateral bounded by bc and the continuations of the two sides eb and dc . The area of $|T|$ depending on b, c, d, e , satisfies:*

$$-4i |T|_{\bar{b}} = (a - c) + (a - b)|ae|/|be|, \quad (12)$$

$$-4i |T|_{\bar{c}} = (b - a) + (c - a)|ad|/|cd|, \quad (13)$$

$$-4i |T|_{\bar{d}} = (a - c)|ac|/|cd|, \quad (14)$$

$$-4i |T|_{\bar{e}} = (b - a)|ab|/|be|. \quad (15)$$

Proof Observe that the vertex a is a function of b, c, d, e and each proof starts by differentiating (11) with respect to these variables. For example, in order to get (13) we have

$$4i |T|_{\bar{c}} = -(c - b)\bar{a}_{\bar{c}} + (\bar{c} - \bar{b})a_{\bar{c}} + (a - b). \quad (16)$$

To compute the partial derivatives of a and $\bar{a}$ with respect to $\bar{c}$ we consider the collinear conditions of a with b and e and with c and d , i.e.

$$(a - b)(\bar{e} - \bar{b}) - (e - b)(\bar{a} - \bar{b}) = 0, \quad (17)$$

$$(a - c)(\bar{c} - \bar{d}) - (c - d)(\bar{a} - \bar{c}) = 0, \quad (18)$$

and, by differentiating, we obtain

$$(\bar{e} - \bar{b})a_{\bar{c}} - (e - b)\bar{a}_{\bar{c}} = 0,$$

$$(\bar{c} - \bar{d})a_{\bar{c}} - (c - d)\bar{a}_{\bar{c}} = d - a.$$

Solving for $a_{\bar{c}}, \bar{a}_{\bar{c}}$ in the previous equations we obtain

$$a_{\bar{c}} = (e - b)(d - a) / ((e - b)(\bar{c} - \bar{d}) - (\bar{e} - \bar{b})(c - d)),$$

$$\bar{a}_{\bar{c}} = (\bar{e} - \bar{b})(d - a) / ((e - b)(\bar{c} - \bar{d}) - (\bar{e} - \bar{b})(c - d)).$$

By substituting in (16) we get

$$\frac{4}{i} \frac{\partial |T|}{\partial \bar{c}} = (b - a) + (d - a) \frac{(c - b)(\bar{e} - \bar{b}) - (\bar{c} - \bar{b})(e - b)}{(e - b)(\bar{c} - \bar{d}) - (\bar{e} - \bar{b})(c - d)}.$$

The proof of (13) is obtained via the formula

$$(c - b)(\bar{e} - \bar{b}) - (\bar{c} - \bar{b})(e - b) = \frac{|ac|}{|cd|} ((e - b)(\bar{c} - \bar{d}) - (\bar{e} - \bar{b})(c - d)). \quad (19)$$

Indeed, by subtracting the left-hand side of (17) from the left hand side of (19) we deduce

$$(c - b)(\bar{e} - \bar{b}) - (\bar{c} - \bar{b})(e - b) = (c - a)(\bar{e} - \bar{b}) - (\bar{c} - \bar{a})(e - b).$$

Since a, c, d are collinear the vector $(c - a)$ is proportional to the vector $(d - c)$ by the factor $|ac|/|cd|$. This proves (19).

The proof of (12) can be obtained in a similar way or simply interchanging b and c, e and d , by a reflection. We remark that such a reflection changes the sign as in formula (11).

The proof of (14) follows from similar computations. More explicitly, by differentiating (11) with respect to $\bar{d}$ we obtain

$$4i|T|_{\bar{d}} = -(c - b)\bar{a}_{\bar{d}} + (\bar{c} - \bar{b})a_{\bar{d}}. \quad (20)$$

By differentiating with respect to $\bar{d}$ the constraints (17) and (18), we get

$$(\bar{e} - \bar{b})a_{\bar{d}} - (e - b)\bar{a}_{\bar{d}} = 0,$$

$$(\bar{c} - \bar{d})a_{\bar{d}} - (c - d)\bar{a}_{\bar{d}} = a - c.$$

This implies

$$a_{\bar{d}} = (e - b)(a - c) / ((e - b)(\bar{c} - \bar{d}) - (\bar{e} - \bar{b})(c - d)),$$

$$\bar{a}_{\bar{d}} = (\bar{e} - \bar{b})(a - c) / ((e - b)(\bar{c} - \bar{d}) - (\bar{e} - \bar{b})(c - d)).$$

By substituting in (20) we get

$$\frac{4}{i} \frac{\partial |T|}{\partial \bar{d}} = (a - c) \frac{(c - b)(\bar{e} - \bar{b}) - (\bar{c} - \bar{b})(e - b)}{(e - b)(\bar{c} - \bar{d}) - (\bar{e} - \bar{b})(c - d)},$$

and from (19) we prove (14). A similar argument proves (15). $\square$

2.1 Inner Case

Proposition 2.4 *Let P^* be a maximizer of F on $\mathcal{P}_n$. Then there exist real numbers $\mu_1, \dots, \mu_n$ such that the following equations hold for the vertices of P^* :*

$$(z_{j-1} - z_{j+1}) = \mu_{j-1}(z_{j-1} - z_{j-2}) + \mu_j(z_{j-1} - z_{j+1}) + \mu_{j+1}(z_{j+2} - z_{j+1}), \quad (21)$$

for $j = 1, \dots, n$.

Proof By Proposition 1.1, each maximizer P^* satisfies the inner equal-area property. Choosing a suitable affine transformation we can assume that $|W_k(P)| = 1$ for $k = 1, \dots, n$. By the Lagrange multipliers argument we have that at P^* the gradient of the area functional $|P|$ is a linear combination of the gradients of the constraints $|W_k(P)| = 1$, i.e. there exist real multipliers μ_k such that at P^* ,

$$\frac{\partial |P|}{\partial \bar{z}_j} = \sum_k \mu_k \frac{\partial |W_k|}{\partial \bar{z}_j} \quad \text{for } j = 1, \dots, n.$$

Since W_k depends only on z_{k-1}, z_k, z_{k+1} we have that

$$\frac{\partial |W_k|}{\partial \bar{z}_j} = 0 \quad \text{for } j \notin \{k-1, k, k+1\}.$$

Now we apply Lemma 2.2 to the triangles W_{j-1}, W_j, W_{j+1} with respect to the vertex z_j and find that

$$\begin{aligned} \frac{4}{i} \frac{\partial |W_{j-1}|}{\partial \bar{z}_j} &= z_{j-1} - z_{j-2}, & \frac{4}{i} \frac{\partial |W_j|}{\partial \bar{z}_j} &= z_{j-1} - z_{j+1}, \\ \frac{4}{i} \frac{\partial |W_{j+1}|}{\partial \bar{z}_j} &= z_{j+2} - z_{j+1}. \end{aligned}$$

Since P can be decomposed in the disjoint subsets $W_j, P \setminus W_j$ and the latter does not depend on the vertex z_j , we have also

$$\frac{4}{i} \frac{\partial |P|}{\partial \bar{z}_j} = \frac{4}{i} \frac{\partial |W_j|}{\partial \bar{z}_j} = z_{j-1} - z_{j+1}. \quad (22)$$

The six previous equations prove (21). $\square$

The system (21) is a system of $2n$ real equations in the $3n$ real unknowns z_j, μ_j , and so it cannot determine the maximizers. The inner equal-area property adds more information. In particular, we have that for each j the vector $(z_{j+2} - z_{j-1})$ is parallel to $(z_{j+1} - z_j)$. So any maximizer P^* also satisfies the following system for suitable positive λ_j :

$$(z_{j+2} - z_{j-1}) = \lambda_j(z_{j+1} - z_j) \quad \text{for } j = 1, \dots, n. \quad (23)$$

We notice that the λ_j 's are well defined and positive because all vertices of P^* are distinct and P^* is convex. Moreover, we can prove that

$$\lambda_j > 1 \quad \text{for } j = 1, \dots, n. \quad (24)$$

Indeed, suppose that there exists a $\lambda_j \leq 1$. Up to an affine transformation we can assume that z_{j-1}, z_j, z_{j+1} are three consecutive vertices of a square Q . Since $\lambda_j \leq 1$, by (23), z_{j+2} belongs to Q . Using (23) again, we obtain that $z_{j+3} - z_j$ has the direction of $z_{j+2} - z_{j+1}$. This means that z_{j+3} belongs to a line supporting Q at z_j . Any choice of z_{j+3} on such a line gives a contradiction, since either z_{j-1} belongs to the convex hull of the other vertices, or the side $z_{j+2}z_{j+3}$ intersects the boundary of Q .

By the definition (23) of λ_j , we have $\lambda_j = d_j/l_j$ and then the inner-ratio property (5) can be rewritten as

$$\frac{(\lambda_{j-2} - 1)(\lambda_{j-1} - 1)}{\lambda_{j-2}} = \frac{(\lambda_{j+1} - 1)(\lambda_j - 1)}{\lambda_{j+1}} \quad \text{for } j = 1, \dots, n. \quad (25)$$

This is the form we get in the following proof.

Proof of Theorem 1.4 From (23) we get

$$\begin{aligned} z_{j+2} &= z_{j-1} + \lambda_j(z_{j+1} - z_j) \quad \text{and} \\ z_{j-2} &= z_{j+1} - \lambda_{j-1}(z_j - z_{j-1}). \end{aligned}$$

By substituting in (21) and rearranging the terms we obtain

$$\begin{aligned} &(z_{j+1} - z_j)(\mu_{j-1} + \mu_j + \mu_{j+1} - 1 - \lambda_j \mu_{j+1}) \\ &+ (z_j - z_{j-1})(\mu_{j-1} + \mu_j + \mu_{j+1} - 1 - \lambda_{j-1} \mu_{j-1}) = 0. \end{aligned}$$

Since $P^* \in \mathcal{P}_n$ the vertices z_j are distinct and no three collinear. This implies that the vectors $(z_{j+1} - z_j), (z_j - z_{j-1})$ are linearly independent and their coefficients in the previous equation must be zero. Hence, we get for $j = 1, \dots, n$,

$$\begin{cases} \mu_{j-1} + \mu_j + \mu_{j+1} - 1 = \lambda_j \mu_{j+1}, \\ \mu_{j-1} + \mu_j + \mu_{j+1} - 1 = \lambda_{j-1} \mu_{j-1}. \end{cases} \quad (26)$$

We infer that the λ_j 's are such that the linear system (26) has a solution $\mu_1, \mu_2, \dots, \mu_n$. Taking into account the six equations involving only the five unknowns

$\mu_{j-2}, \mu_{j-1}, \mu_j, \mu_{j+1}$, and μ_{j+2} , we deduce that the determinant of the matrix

$$\begin{pmatrix} 1 & 1 & 1 - \lambda_{j-1} & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 - \lambda_j & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 - \lambda_{j+1} & 1 \\ 1 - \lambda_{j-2} & 1 & 1 & 0 & 0 & 1 \\ 0 & 1 - \lambda_{j-1} & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 - \lambda_j & 1 & 1 & 1 \end{pmatrix}$$

has to be zero. After some manipulations, we get the following equation

$$\lambda_j \lambda_{j-2} (1 - \lambda_{j+1}) \lambda_{j-1} (1 - \lambda_j) - \lambda_{j-1} \lambda_{j+1} (1 - \lambda_{j-2}) \lambda_j (1 - \lambda_{j-1}) = 0. \quad (27)$$

Since all $\lambda_j > 0$, the previous equation can be simplified as in (25). $\square$

2.2 Outer Case

In order to get the Lagrange multipliers system for the outer case in a simple way we recall that v_j is the intersection of the two lines through the consecutive vertices z_{j-1}, z_j and the vertices z_{j+1}, z_{j+2} , (see Fig. 1), $l_j = \|z_{j+1} - z_j\|$, i.e., the length of the j th side, and $s_j = \|v_j - z_j\|$, $p_j = \|z_{j+1} - v_j\|$.

Proposition 2.5 *Let P^* be a maximizer of the functional G on $\mathcal{P}_n$, $n \geq 5$. Then there exist real numbers $\eta_1, \dots, \eta_n$ such that the following equations hold for the vertices of P^* , for $j = 1, \dots, n$:*

$$\begin{aligned} (z_{j-1} - z_{j+1}) &= \eta_{j-2} (z_{j-1} - v_{j-2}) \frac{p_{j-2}}{l_{j-1}} \\ &\quad + \eta_{j-1} \left((v_{j-1} - z_{j-1}) + (v_{j-1} - z_{j+1}) \frac{p_{j-1}}{l_j} \right) \\ &\quad + \eta_j \left((z_{j+1} - v_j) + (z_{j-1} - v_j) \frac{s_j}{l_{j-1}} \right) \\ &\quad + \eta_{j+1} (v_{j+1} - z_{j+1}) \frac{s_{j+1}}{l_j}. \end{aligned} \quad (28)$$

Proof By Proposition 1.1 each maximizer P^* satisfies the outer equivalent triangle property. Choosing a suitable affine transformation we can assume that $|T_j(P)| = 1$, for $j = 1, \dots, n$. By the Lagrange multipliers argument, at P^* the gradient of the area functional $|P|$ is a linear combination of the gradients of the constraints $|T_k(P)| = 1$, i.e. there exist real multipliers η_j such that, at P^* ,

$$\frac{\partial |P|}{\partial \bar{z}_j} = \sum_k \eta_k \frac{\partial |T_k|}{\partial \bar{z}_j} \quad \text{for } j = 1, \dots, n.$$

Since $|T_k|$ depends only on $z_{k-1}, z_k, z_{k+1}, z_{k+2}$ we get

$$\frac{4}{i} \frac{\partial |T_k|}{\partial \bar{z}_j} = 0 \quad \text{for } k \notin \{j-2, j-1, j, j+1\}.$$

By Lemma 2.3, we obtain

$$\begin{aligned}\frac{4}{i}|T_{j-2}|\overline{z_j} &= (z_{j-1} - v_{j-2})\frac{p_{j-2}}{l_{j-1}}, \\ \frac{4}{i}|T_{j-1}|\overline{z_j} &= (v_{j-1} - z_{j-1}) + (v_{j-1} - z_{j+1})\frac{p_{j-1}}{l_j}, \\ \frac{4}{i}|T_j|\overline{z_j} &= (z_{j+1} - v_j) + (z_{j-1} - v_j)\frac{s_j}{l_{j-1}}, \\ \frac{4}{i}|T_{j+1}|\overline{z_j} &= (v_{j+1} - z_{j+1})\frac{s_{j+1}}{l_j}.\end{aligned}$$

The six previous equations and (22) yield (28). $\square$

Since the lengths p_j, l_j, s_j are functions of the vertices z_j , (28) is a system of $2n$ real equations in the unknowns z_j, η_j , i.e. $3n$ real unknowns, and so it cannot determine the maximizers. From the outer equivalent triangle property of any maximizer P^* we have that, for each j , the vector $(v_j - v_{j-1})$ is parallel to $(z_{j+1} - z_{j-1})$. Hence P^* satisfies also the following system for suitable positive ζ_j :

$$(v_j - v_{j-1}) = \zeta_j(z_{j+1} - z_{j-1}) \quad \text{for } j = 1, \dots, n. \quad (29)$$

We notice that the triangles v_j, z_j, v_{j-1} and z_{j+1}, z_j, z_{j-1} are similar and then

$$\zeta_j = \frac{|v_j - v_{j-1}|}{|z_{j+1} - z_{j-1}|} = \frac{s_j}{l_{j-1}} = \frac{p_{j-1}}{l_j}. \quad (30)$$

Thus, the outer-ratio property (6) can be rewritten as

$$\frac{(1 + \zeta_{j+2})}{\zeta_{j+1}(1 + \zeta_{j+1} + \zeta_{j+2})} = \frac{(1 + \zeta_{j-1})}{\zeta_j(1 + \zeta_j + \zeta_{j-1})} \quad \text{for } j = 1, \dots, n. \quad (31)$$

Proof of Theorem 1.5 Since v_j, z_j , and z_{j-1} are collinear, (30) implies that we can express v_j in terms of z_j, z_{j-1} and positive ζ_j

$$v_j = z_j + \zeta_j(z_j - z_{j-1}).$$

Similarly, (30) also implies that

$$\begin{aligned}v_{j+1} &= z_{j+1} + \zeta_{j+1}(z_{j+1} - z_j), \\ v_{j-1} &= z_j + \zeta_{j-1}(z_j - z_{j+1}), \\ v_{j-2} &= z_{j-1} + \zeta_{j-1}(z_{j-1} - z_j).\end{aligned}$$

This permits us to eliminate in (28) all the vectors v_j , i.e.,

$$\begin{aligned}(z_{j-1} - z_{j+1}) &= \eta_{j-2}\zeta_{j-1}^2(z_j - z_{j-1}) \\ &\quad + \eta_{j-1}[(z_j - z_{j-1}) + \zeta_j(z_j - z_{j+1}) + (z_j - z_{j+1})(1 + \zeta_j)\zeta_j]\end{aligned}$$

$$\begin{aligned}
& + \eta_j[(z_{j+1} - z_j) - \zeta_j(z_j - z_{j-1}) + (z_{j-1} - z_j)(1 + \zeta_j)\zeta_j] \\
& + \eta_{j+1}(z_{j+1} - z_j)\zeta_{j+1}^2.
\end{aligned} \tag{32}$$

Rearranging the terms we obtain

$$\begin{aligned}
& (z_{j-1} - z_j)(1 + \eta_{j-2}\zeta_{j-1}^2 + \eta_{j-1} - \eta_j(2\zeta_j + \zeta_j^2)) \\
& + (z_j - z_{j+1})(1 - \eta_{j-1}(2\zeta_j + \zeta_j^2) + \eta_j + \eta_{j+1}\zeta_{j+1}^2) = 0.
\end{aligned}$$

Since $P^* \in \mathcal{P}_n$, the vertices z_j are distinct and not collinear. This implies that the vectors $(z_{j+1} - z_j)$, $(z_j - z_{j-1})$ are linearly independent and their coefficients in the previous equation must be zero, i.e.,

$$\begin{cases} \eta_{j-2}\zeta_{j-1}^2 + \eta_{j-1} - \eta_j(2\zeta_j + \zeta_j^2) = -1, \\ -\eta_{j-1}(2\zeta_j + \zeta_j^2) + \eta_j + \eta_{j+1}\zeta_{j+1}^2 = -1 \end{cases} \tag{33}$$

for $j = 1, \dots, n$. We shift by one the index in the first equation and adding to and subtracting from the second one, we obtain

$$\begin{cases} \eta_{j-1}\zeta_j - \eta_j + \eta_{j+1}\zeta_{j+1} = 1, \\ \eta_{j-1}\zeta_j(1 + \zeta_j) - \eta_{j+1}\zeta_{j+1}(1 + \zeta_{j+1}) = 0 \end{cases} \tag{34}$$

for $j = 1, \dots, n$.

We infer that the ζ_j 's are such that the linear system (34) has a solution $\eta_1, \eta_2, \dots, \eta_n$. Taking into account the six equations involving only the five unknowns $\eta_{j-2}, \dots, \eta_{j+2}$, we deduce that the determinant of the matrix

$$\begin{pmatrix} \zeta_{j-1} & -1 & \zeta_j & 0 & 0 & 1 \\ 0 & \zeta_j & -1 & \zeta_{j+1} & 0 & 1 \\ 0 & 0 & \zeta_{j+1} & -1 & \zeta_{j+2} & 1 \\ \zeta_{j-1} + \zeta_{j-1}^2 & 0 & -\zeta_j - \zeta_j^2 & 0 & 0 & 0 \\ 0 & \zeta_j + \zeta_j^2 & 0 & -\zeta_{j+1} - \zeta_{j+1}^2 & 0 & 0 \\ 0 & 0 & \zeta_{j+1} + \zeta_{j+1}^2 & 0 & -\zeta_{j+2} - \zeta_{j+2}^2 & 0 \end{pmatrix}$$

has to be zero. After some manipulations we obtain

$$\begin{aligned}
& \zeta_{j-1}(1 + \zeta_j)(1 + \zeta_{j+1})\zeta_{j+2}[\zeta_j(1 + \zeta_{j+2})(1 + \zeta_j + \zeta_{j-1}) \\
& - \zeta_{j+1}(1 + \zeta_{j-1})(1 + \zeta_{j+1} + \zeta_{j+2})] = 0.
\end{aligned}$$

Since $\zeta_j > 0$, the previous equation can be simplified as in the statement (31) and we conclude the proof. $\square$

3 Circulant Systems

In this section we have collected the steps of the proofs which are not strictly related to the geometry of the problem. We focus on the systems (25), (31) and we prove that the only suitable solutions are those independent of j .

First we deal with the inner problem.

Proposition 3.1 *Any solution of the system of equations*

$$\frac{(\lambda_{j-2} - 1)(\lambda_{j-1} - 1)}{\lambda_{j-2}} = \frac{(\lambda_{j+1} - 1)(\lambda_j - 1)}{\lambda_{j+1}},$$

for $j = 1, 2, \dots, n$, with $\lambda_j > 1$ for all j , satisfies $\lambda_i = \lambda_j$ for all i and j .

Proof Set $\gamma_j = 1/(\lambda_j - 1)$, for all j . Notice that $\lambda_j > 1$ implies $\gamma_j > 0$. Then (25) reduce to

$$\gamma_{j-1}(1 + \gamma_{j-2}) = \gamma_j(1 + \gamma_{j+1}). \quad (35)$$

Consider the maximum number among the expressions $\gamma_j(1 + \gamma_{j-1})$, for all j . Assume this maximum is attained by $\gamma_M(1 + \gamma_{M-1})$. From (35) we obtain

$$\gamma_M(1 + \gamma_{M-1}) \geq \gamma_j(1 + \gamma_{j+1}),$$

for every $j = 1, 2, \dots, n$. Now, from $\gamma_M(1 + \gamma_{M-1}) \geq \gamma_{M-1}(1 + \gamma_M)$ we have

$$\gamma_M \geq \gamma_{M-1}. \quad (36)$$

From $\gamma_M(1 + \gamma_{M-1}) \geq \gamma_M(1 + \gamma_{M+1})$ we find

$$\gamma_{M-1} \geq \gamma_{M+1}. \quad (37)$$

From (35) we deduce that $\gamma_M(1 + \gamma_{M-1}) = \gamma_{M+1}(1 + \gamma_{M+2})$ and so we can also write $\gamma_{M+1}(1 + \gamma_{M+2}) \geq \gamma_{M+1}(1 + \gamma_M)$, which implies that

$$\gamma_{M+2} \geq \gamma_M, \quad (38)$$

and $\gamma_{M+1}(1 + \gamma_{M+2}) \geq \gamma_{M+2}(1 + \gamma_{M+1})$, which yields

$$\gamma_{M+1} \geq \gamma_{M+2}. \quad (39)$$

By combining inequalities (36–39) we discover $\gamma_{M+1} \geq \gamma_{M+2} \geq \gamma_M \geq \gamma_{M-1} \geq \gamma_{M+1}$ and so $\gamma_{M-1} = \gamma_M = \gamma_{M+1} = \gamma_{M+2}$. Taking into account (35) again yields $\gamma_i = \gamma_j$, for all i and j . $\square$

We now turn to the outer problem.

Proposition 3.2 *Any solution of the system of equations*

$$\frac{1 + \zeta_{j+2}}{\zeta_{j+1}(1 + \zeta_{j+1} + \zeta_{j+2})} = \frac{1 + \zeta_{j-1}}{\zeta_j(1 + \zeta_j + \zeta_{j-1})}$$

for $j = 1, 2, \dots, n$, with $\zeta_j > 0$ for all j , satisfies $\zeta_i = \zeta_j$ for all i and j .

Proof Set

$$a_j = \frac{\zeta_j}{1 + \zeta_j + \zeta_{j+1}}, \quad b_j = \frac{\zeta_{j+1}}{1 + \zeta_j + \zeta_{j+1}}. \quad (40)$$

Notice that $\zeta_j > 0$, for all j , implies that all a_j and b_j are positive and that

$$a_j + b_j < 1. \quad (41)$$

The system (31) can be rewritten in terms of a_j, b_j as

$$\frac{1 - a_{j+1}}{b_j} = \frac{1 - b_{j-1}}{a_j}. \quad (42)$$

Furthermore, it is easy to check that the relation

$$\frac{1 - a_j}{b_j} = \frac{1 - b_{j+1}}{a_{j+1}} \quad (43)$$

also holds for every $j = 1, 2, \dots, n$.

Now consider the maximum number among the expressions $(1 - a_{j+1})/b_j$ or $(1 - a_j)/b_j$ involved in (42) and (43) and call it C . Assume first that C appears in (42), i.e., $C = \frac{1 - a_{M+1}}{b_M}$, for some M . From $\frac{1 - a_{M+1}}{b_M} \geq \frac{1 - a_M}{b_M}$ we infer

$$a_M \geq a_{M+1}. \quad (44)$$

From

$$\frac{1 - b_{M-1}}{a_M} = \frac{1 - a_{M+1}}{b_M} \geq \frac{1 - b_M}{a_M}$$

we deduce

$$b_M \geq b_{M-1}. \quad (45)$$

From

$$\frac{1 - a_{M+1}}{b_M} = \frac{1 - b_{M-1}}{a_M} \geq \frac{1 - a_M}{b_{M-1}}$$

we deduce $b_{M-1} - b_{M-1}^2 \geq a_M - a_M^2$ and then

$$(b_{M-1} - a_M)(1 - b_{M-1} - a_M) \geq 0.$$

Since $b_{M-1} \leq b_M$ by (45), inequality (41) implies that $(1 - b_{M-1} - a_M) > 0$. Therefore

$$b_{M-1} \geq a_M. \quad (46)$$

Analogously, from

$$\frac{1 - a_{M+1}}{b_M} \geq \frac{1 - a_{M+2}}{b_{M+1}} = \frac{1 - b_M}{a_{M+1}}$$

we infer

$$(a_{M+1} - b_M)(1 - a_{M+1} - b_M) \geq 0.$$

Inequalities (41), (44) ensure that $(1 - a_{M+1} - b_M) > 0$ and so

$$a_{M+1} \geq b_M. \quad (47)$$

Combining inequalities (44–47) we obtain

$$b_{M-1} \geq a_M \geq a_{M+1} \geq b_M \geq b_{M-1}$$

and so

$$b_{M-1} = a_M = a_{M+1} = b_M.$$

Now it is easy to use (42) and (43) to deduce that all a_j and b_j have to be equal, and from this fact and (40) that all the ζ_j 's are equal.

Assume now that C appears in (43), i.e., $C = (1 - a_M)/b_M$, for some M . From $(1 - a_M)/b_M \geq (1 - b_M)/a_M$ and (41) we deduce

$$a_M \geq b_M. \quad (48)$$

From $\frac{1-b_{M+1}}{a_{M+1}} = \frac{1-a_M}{b_M} \geq \frac{1-a_{M+1}}{b_{M+1}}$ and (41) we deduce

$$b_{M+1} \geq a_{M+1}. \quad (49)$$

From $\frac{1-b_{M+1}}{a_{M+1}} = \frac{1-a_M}{b_M} \geq \frac{1-b_M}{a_{M+1}}$ we infer

$$b_M \geq b_{M+1}. \quad (50)$$

From $\frac{1-a_M}{b_M} \geq \frac{1-a_{M+1}}{b_M}$ we obtain

$$a_{M+1} \geq a_M. \quad (51)$$

Once again we have obtained a sequence of inequalities that can be satisfied only if $a_M = a_{M+1} = b_M = b_{M+1}$. As in the previous case, it is now easy to conclude that all ζ_j have to be equal. $\square$

G. Ottaviani has analyzed the case $n = 7$ for the inner and outer problem. By using the computer algebra system Macaulay, he found that in the complex variables $\gamma_1, \dots, \gamma_7$, the system (35) has a solution set which consists of the trivial line $\gamma_1 = \dots = \gamma_7$ and of an algebraic surface of degree 14. It is interesting that the line and the surface are disjoint components. Similarly, he found that for the outer problem the solution set of the system (31) contains the trivial line and a surface of degree 71.

4 Proofs of Theorems 1.6 and 1.7

In this section we complete the characterization of the maximizers of F and G , thus proving Theorems 1.6, 1.7. To do this we turn back to geometry.

Proof of Theorem 1.6 As noted in Sect. 2.1, each polygon P contained in $\Phi_n \cap \mathcal{F}_n$ satisfies (23–25). By Proposition 3.1 we deduce that there exists a $\lambda > 1$ such that

$$z_{j+2} - z_{j-1} = \lambda(z_{j+1} - z_j), \quad \text{for all } j. \quad (52)$$

Such a condition is satisfied only by affinely regular n -gons. This is proved in [5, Statement 3, Theorem 1], where the result is attributed to Coxeter [3]. We provide a proof for completeness and for the convenience of the reader.

Up to an affine transformation, we can assume that three consecutive edges of P , z_1z_2 , z_2z_3 and z_3z_4 , say, all have length one. By (52), the quadrilateral $z_1z_2z_3z_4$ is an isosceles trapezium, and so the angles of P at z_2 and z_3 are also equal. Denote by r_2 the bisector of the angle at z_2 . By assumption, r_2 is an axis of symmetry of the segment z_1z_3 . Since the vertex z_n , consecutive to z_1 , is determined by the relation $z_n - z_3 = \lambda(z_1 - z_2)$ and z_4 by $z_4 - z_1 = \lambda(z_3 - z_2)$, it is easily seen that r_2 is an axis of symmetry of the segment z_4z_n . Similarly, z_{n-1} and z_5 are uniquely determined by the previous vertices (and λ) and then r_2 is also an axis of symmetry of the segment z_5z_{n-1} . This argument shows that r_2 is a symmetry axis of P . Analogously, considering the vertex z_3 of the trapezium $z_1z_2z_3z_4$, the bisector r_3 of the angle at z_3 is a symmetry axis of P . This clearly implies that P is a regular polygon. $\square$

Proof of Theorem 1.7 As noted in Sect. 2.2, each polygon P contained in $\Gamma_n \cap \mathcal{G}_n$ satisfies systems (29), (31). By Proposition 3.2 we deduce that there exists a $\zeta > 0$ such that

$$v_j - v_{j-1} = \zeta(z_{j+1} - z_{j-1}), \quad \text{for all } j. \quad (53)$$

This means that the pairs of triangles $v_{j-1}v_jz_j$, $z_{j+1}z_{j-1}z_j$ are all similar with the same ratio of similarity. Hence,

$$v_j - z_j = \zeta(z_j - z_{j-1}), \quad v_j - z_{j+1} = \zeta(z_{j+1} - z_{j+2}).$$

Therefore the triangles $v_jz_{j+1}z_j$, $v_jz_{j+2}z_{j-1}$ are also similar and

$$z_{j+2} - z_{j-1} = (1 + \zeta)(z_{j+1} - z_j), \quad \text{for all } j. \quad (54)$$

Such a condition is just like (52) and the argument of the previous proof implies that P is an affinely regular polygon. $\square$

Theorems 1.6 and 1.7 were the last ingredients in the proof of Theorem 1.8, whose quantitative form is the following.

Theorem 4.1 *Let $P \in \mathcal{P}_n$, $n \geq 5$, then*

$$\min_{j=1,\dots,n} |T_j(P)| \leq |P| \frac{2 \sin^2(\pi/n)}{n \cos(2\pi/n)}, \quad (55)$$

$$\min_{j=1,\dots,n} |W_j(P)| \leq |P| \frac{4 \sin^2(\pi/n)}{n}, \quad (56)$$

and equality holds if and only if P is an affinely regular n -gon.

Notice that (55) implies the inequality in [17, Lemma 3].

5 Extensions and Affine Length

Let C be a planar convex body, and let γ be a connected closed proper subset of ∂C , i.e., a closed arc with endpoints a and b . Define $C(\gamma)$ as the intersection of all half-planes supporting C at boundary points in $\partial C \setminus \gamma$. The boundary of $C(\gamma)$ is obtained by extending $\partial C \setminus \gamma$ with the continuations of the half-lines tangent to $\partial C \setminus \gamma$ at a and b . Consider the region $I(C, \gamma) = C(\gamma) \setminus C$ and the area ratio $|I(C, \gamma)|/|C|$. Now let Γ_n be a finite family of connected closed subsets γ_i of ∂C with disjoint interiors:

$$\Gamma_n = \{\gamma_1, \dots, \gamma_n: \gamma_i \subset \partial C, \text{int}(\gamma_i) \cap \text{int}(\gamma_j) = \emptyset \text{ for } i \neq j, i, j = 1, \dots, n\}.$$

Define

$$G(C, \Gamma_n) = \min_{j=1, \dots, n} \frac{|I(C, \gamma_j)|}{|C|}, \quad (57)$$

and

$$G_n(C) = \max_{\Gamma_n} G(C, \Gamma_n). \quad (58)$$

Theorem 5.1 *For $n > 4$ and for any planar convex body C we have*

$$G_n(C) \leq \max_{P \in \mathcal{P}_n} G(P). \quad (59)$$

Equality holds if and only if C is an affinely regular n -gon.

Proof Let C be a planar convex body and let Γ be a finite family of arcs γ_i on ∂C with endpoints a_i, b_i . We can consider families of arcs which partition ∂C , i.e., such that $a_{i+1} = b_i$ for all i . This can be proved by taking a partition Γ^* with arcs γ_i^* so that $\gamma_i \subset \gamma_i^*$. Since $I(C, \gamma_i) \subset I(C, \gamma_i^*)$ we have $G(C, \Gamma) \leq G(C, \Gamma^*)$.

Let P be the convex polygon whose vertices are the endpoints a_i, b_i of the arcs γ_i of Γ . Clearly, $P \subset C$ and P has exactly n edges. Moreover, if $T_i(P)$ is the outer triangle defined in the introduction corresponding to the edge with vertices a_i, b_i , then $I(C, \gamma_i) \subset T_i(P)$, and

$$G(C, \Gamma) \leq G(P).$$

This immediately yields (59) and the equality conditions. $\square$

Corollary 5.2 *Let K and K' be two planar convex bodies such that their symmetric difference has $n > 4$ connected components $C_1, \dots, C_n$. Then*

$$\min_{i=1, \dots, n} |C_i| \leq |K \cap K'| \max_{P \in \mathcal{P}_n} G(P). \quad (60)$$

Equality holds if and only if $K \cap K'$ is an affinely regular polygon with $n = 2m$ edges and, up to an affine transformation, K and K' are two congruent regular m -gons, and K' is K rotated by π/m about its center.

Proof We consider the family Γ of the closed arcs

$$\gamma_i = \partial C_i \cap \partial(K \cap K').$$

Since

$$|C_i| \leq |I(K \cap K', \gamma_i)|,$$

we can apply the previous theorem to $C = K \cap K'$. Equality holds if and only if C is an affinely regular $2m$ -gon and so K and K' have to be affinely regular m -gons. $\square$

We present another possible extension of the previous area functionals. Let C be a convex body. Let D be a convex body containing C and such that $D \setminus C$ consists of at least m connected components $D_1, \dots, D_m$. Let

$$H_m(C, D) = \min_{i=1, \dots, m} |D_i|.$$

Arguing as above, when we look for the maximizers of $H_m(C, D)|C|^{-1}$ it is easy to see that one can assume that C and D are m -gons. The number $\#(D \setminus C)$ of the connected components of $D \setminus C$ is m , and hence the edges of D support C at its vertices. We get the following result.

Theorem 5.3 *For $m \geq 3$ and for any two planar convex bodies C, D with $C \subset D$, $\#(D \setminus C) = m$, we have*

$$\frac{H_m(C, D)}{|C|} \leq \frac{1}{m} \tan^2 \frac{\pi}{m}. \quad (61)$$

Equality holds if and only if C, D are affinely regular m -gons and the vertices of C are the midpoints of the edges of D .

Proof Standard compactness arguments show that $H_m(C, D)|C|^{-1}$ has a maximum. As already stated in advance, we can assume that such a maximum is attained when C and D are two m -gons P and S , $P \subset S$. Let z_j be the vertices of P and v_j the vertices of S labeled so that z_j belongs to the segment $v_j v_{j-1}$.

First we prove that all triangles S_j with vertices z_j, v_j, z_{j+1} have equal area. Indeed, if $|S_k| > \min_{j=1, \dots, m} |S_j|$, then a small counterclockwise rotation of the side of S through z_k around z_k decreases the area of S_k and increases the area of S_{k-1} . Possible iterations of this procedure to different edges of S would permit to increase the value of our functional. The assumption about the maximality of the pair (P, S) implies

$$|S_k| = |S_{k-1}| \quad \text{for } k = 1, \dots, n. \quad (62)$$

Arguing as in the proof of Proposition 2.4, we consider the Lagrange multipliers system relative to $|P|$ as a function of z_j, v_j , under the constraints $|S_j| \geq \text{const}$, $z_j \in v_{j-1} v_j$. Denote by μ_j the parameters corresponding to the constraints $|S_j| \geq \text{const}$, and by ν_j that corresponding to $0 = \det(z_j - v_{j-1}, z_j - v_j) \equiv A_j$. Hence, we

obtain the system

$$\begin{aligned}\frac{\partial |P|}{\partial \bar{z}_j} &= \sum_{k=1}^m \mu_k \frac{\partial |S_k|}{\partial \bar{z}_j} + \sum_{k=1}^m v_k \frac{\partial A_k}{\partial \bar{z}_j} \quad \text{for } j = 1, \dots, n, \\ 0 &= \frac{\partial |P|}{\partial \bar{v}_j} = \sum_{k=1}^m \mu_k \frac{\partial |S_k|}{\partial \bar{v}_j} + \sum_{k=1}^m v_k \frac{\partial A_k}{\partial \bar{v}_j} \quad \text{for } j = 1, \dots, n.\end{aligned}$$

Since we are looking for minimizers of $|P|$ with constraints $|S_j| \geq \text{const}$, we have

$$\mu_j \geq 0 \quad \text{for } j = 1, \dots, n. \quad (63)$$

Taking into account the results in Sect. 2 we get:

$$(z_{j-1} - z_{j+1}) = \mu_{j-1}(v_{j-1} - z_{j-1}) + \mu_j(z_{j+1} - v_j) + v_j(v_{j-1} - v_j), \quad (64)$$

$$0 = \mu_j(z_j - z_{j+1}) + v_j(z_j - v_{j-1}) + v_{j+1}(v_{j+1} - z_{j+1}). \quad (65)$$

The constraint $A_j = 0$ can be written as $z_j = v_{j-1} + \rho_j(v_j - v_{j-1})$, where the variables ρ_j satisfy $0 < \rho_j < 1$. Consequently,

$$|S_j| = |(z_{j+1} - v_j) \times (z_j - v_j)| = \rho_{j+1}(1 - \rho_j)|U_j|,$$

where U_j denotes the triangle $v_{j-1}v_jv_{j+1}$. Identity (62) yields

$$\rho_{j+1}(1 - \rho_j)|U_j| = \rho_j(1 - \rho_{j-1})|U_{j-1}|. \quad (66)$$

Equations (64) and (65) become

$$\begin{aligned}(1 + \mu_{j-1})(1 - \rho_{j-1})(v_{j-2} - v_{j-1}) + (1 + \mu_j)\rho_{j+1}(v_j - v_{j+1}) \\ + (v_{j-1} - v_j)(1 - \rho_j) = 0,\end{aligned} \quad (67)$$

$$\begin{aligned}0 = (v_j - v_{j-1})(\rho_j(\mu_j + v_j) - \mu_j) \\ + (v_{j+1} - v_j)(-\rho_{j+1}(\mu_j + v_{j+1}) + v_{j+1}).\end{aligned} \quad (68)$$

Since $(v_j - v_{j-1})$ and $(v_{j+1} - v_j)$ are linearly independent, we obtain

$$\mu_j(1 - \rho_j) = \rho_j v_j, \quad (69)$$

$$\mu_j \rho_{j+1} = (1 - \rho_{j+1})v_{j+1}. \quad (70)$$

Taking the cross product of (67) with the vector $(v_j - v_{j-1})$ yields

$$(1 + \mu_{j-1})(1 - \rho_{j-1})U_{j-1} - (1 + \mu_j)\rho_{j+1}U_j = 0.$$

By means of (66) the latter reduces to

$$(1 + \mu_j)\rho_j = (1 + \mu_{j-1})(1 - \rho_j). \quad (71)$$

The systems (69), (70), (71) are $3n$ real equations in $3n$ real unknowns μ_j, ρ_j, v_j . From the first two equations we get

$$\mu_j(1 - \rho_j)^2 = \mu_{j-1}\rho_j^2. \quad (72)$$

A comparison with (71) implies

$$(1 + \mu_j)^2 \mu_j = (1 + \mu_{j-1})^2 \mu_{j-1}. \quad (73)$$

Since the function $y = (1 + x)^2 x$ is injective for $y > 0$, inequality (63) implies that $\mu_j = \mu$, for all j .

From (71) we deduce $\rho_j = 1/2$ and then, from (69), that $v_j = \mu$, for all j . Hence, (67) gives

$$(1 + \mu)v_{j-2} - 2\mu v_{j-1} + 2\mu v_j - (1 + \mu)v_{j+1} = 0,$$

which implies that $v_{j-2}v_{j+1}$ is parallel to $v_{j-1}v_j$ and their ratio is independent of j .

As in the proof of Theorem 1.6 (see (52)), we deduce that S is affinely regular. $\square$

Given a planar convex body K and a natural number $n \geq 3$, we denote by $\mathcal{P}_n^i(K)$ and $\mathcal{P}_n^c(K)$ the class of n -gons inscribed in K or with K inscribed in them, respectively. For any n distinct points $z_i \in \partial K$ we consider the polygon $P = \text{conv}\{z_1, \dots, z_n\} \in \mathcal{P}_n^i(K)$ and lines s_i supporting K at the vertices z_i of P . Let $S_i(K)$ be the triangle bounded by the lines s_{i-1} , s_i and the side at $z_{i-1}z_i$, and S be the polygon bounded by the lines s_i . Hence, S is an n -gon in $\mathcal{P}_n^c(K)$. This is equivalent to choosing two n -gons P, S , with $P \in \mathcal{P}_n^i(K)$, $S \in \mathcal{P}_n^c(K) \cap \mathcal{P}_n^c(P)$. In this case, for brevity, we say that the pair (P, S) belongs to $\mathcal{P}_n^{i,c}(K)$. Define

$$AL_n(K) = 2n \max \left\{ \min_{j=1, \dots, n} |S_j(K)|^{\frac{1}{3}} : (P, S) \in \mathcal{P}_n^{i,c}(K) \right\}.$$

Since $n \geq 3$, $AL_n(K)$ is finite for every K . The symbol we chose to denote these functionals is justified by some properties listed below, which present AL_n as a discretization of the affine length of the boundary of K . Recalling the functional $H_n(P, S)$ introduced above, we have

$$AL_n(K) = 2n \max \left\{ H_n(P, S)^{\frac{1}{3}} : (P, S) \in \mathcal{P}_n^{i,c}(K) \right\}.$$

As in the proof of Theorem 5.3 it turns out that the previous maximum is attained when all the $S_i(K)$ have the same area. In the sequel we denote by $\mathcal{E}_n$ the class of pairs of n -gons with the property that all the $S_i(K)$ have equal area.

Proposition 5.4 *For every planar convex body K and $n \geq 3$,*

$$AL_n(K) = 2n \max \left\{ H_n(P, S)^{\frac{1}{3}} : (P, S) \in \mathcal{P}_n^{i,c}(K) \cap \mathcal{E}_n \right\}.$$

Proof Clearly, $|S_j(K)|$ is a continuous function of z_i and s_i for all i and j . If z_i moves towards z_{i+1} , then $|S_{i+1}(P)|$ decreases and $|S_i(P)|$ increases, unless one or both of them remain equal to zero. Furthermore, if s_i rotates around z_i counterclockwise, then

$|S_{i+1}(P)|$ decreases and $|S_i(P)|$ increases. These facts easily imply that, if (P, S) does not belong to $\mathcal{E}_n$, then we can move the supporting lines s_i together with the points z_i increasing all $|S_j(K)|$. $\square$

By Proposition 5.4 the functional AL_n is then related to the arithmetic average of the cube root of $|S_j(K)|$, i.e.,

$$AL_n(K) = 2 \max_{(P, S) \in \mathcal{P}_n^{i,c}(K) \cap \mathcal{E}_n} \sum_{j=1}^n |S_j(K)|^{\frac{1}{3}}.$$

This further clarifies the connection with the affine length of K , $\Omega_1(K)$, as defined, for example, by Ludwig [18, Sect. 3]. There Ludwig proved that all upper (or lower) semicontinuous and equi-affine invariant valuations on the space of planar compact convex sets (endowed with the Hausdorff metric) are linear combinations of three basic valuations: the Euler characteristic, the area, and the affine length.

Such a characterization is the main ingredient in the following proof.

Proposition 5.5 *If K is a planar convex body, then*

$$\inf_{n \in \mathbb{N}} AL_n(K) = \Omega_1(K).$$

Proof We first notice that $AL_n(K) = 0$ when K is a polygon with less than n vertices. Moreover, these are the only convex sets where AL_n vanishes.

Set $AL(K) = \inf_n AL_n(K)$. Since the functionals AL_n are continuous and equi-affine invariant, it is easy to verify that AL is equi-affine invariant and upper semicontinuous. Following the arguments used by Ludwig [18, Theorem 2], it can be proved that AL is a valuation, i.e.,

$$AL(H \cup K) + AL(H \cap K) = AL(H) + AL(K),$$

for every pair of convex bodies such that $H \cup K$ is convex. By [18, Theorem 1], it follows that AL is a linear combination of the affine length, the area, and the Euler characteristic. Since $AL_n(P)$ vanishes when P is a polygon with less than n vertices, AL has to be a multiple of the affine length. A simple calculation of AL at the unit disc yields the result. $\square$

We now present some straightforward consequences of Theorem 5.3.

Theorem 5.6 *If K is a planar convex body and $n \geq 3$, then*

$$AL_n(K) \leq 2 \left(n \tan \frac{\pi}{n} \right)^{\frac{2}{3}} \max_{P \in \mathcal{P}_n^i(K)} |P|^{\frac{1}{3}},$$

and equality holds if and only if exists an affinely regular n -gon of maximal area in $\mathcal{P}_n^i(K)$.

Corollary 5.7 *If K is a planar convex body and $n \geq 3$, then*

$$AL_n(K) \leq 2 \left(n \tan \frac{\pi}{n} \right)^{\frac{2}{3}} |K|^{\frac{1}{3}},$$

and equality holds if and only if K is an affinely regular n -gon.

Corollary 5.7 and Proposition 5.5 yield the affine isoperimetric inequality

$$\Omega_1(K) \leq 2\pi^{\frac{2}{3}} |K|^{\frac{1}{3}},$$

without the equality conditions, which are known to characterize ellipses. Notice that equality in the formula of Theorem 5.6 implies that K has at least n points of intersection with a suitable ellipse, and so its Hausdorff distance from that ellipse decreases as n increases.

It is well known that the affine length of a planar convex body K is closely related to the approximation of K by polygons. We refer the interested reader to [9] for an extensive review. Theorem 5.6 implies that, for any planar convex body K and $n \geq 3$, there exists a polygon $P \in \mathcal{P}_n^i(K)$ such that

$$|P| \geq \frac{AL_n(K)^3}{8n^2 \tan^2(\frac{\pi}{n})}.$$

Proposition 5.5 allows $AL_n(K)$ to be replaced with $\Omega_1(K)$ in the previous inequality. This yields a sharp lower bound for the approximation of K with polygons from $\mathcal{P}_n^i(K)$, which also follows from the affine isoperimetric inequality and a result of Blaschke (see [9, Sect. 4]).

6 Applications to Tomography

The word tomography reminds people of the medical CAT scanner, where images are reconstructed from X-rays. Despite the common use of CAT scanners, the mathematical subject related to this reconstruction still deserves interesting and unsolved problems. One of them is stated in the question asked by P.C. Hammer in 1963: *How many parallel X-ray pictures of a convex body must be taken in order to permit its exact reconstruction?* The parallel X-ray of a planar convex body K in a direction θ provides the length of each chord of K parallel to θ .

The existence of finite sets of directions, with arbitrary large cardinality, such that the corresponding X-rays cannot determine a convex body among the others is shown with the following well-known example (see [8]).

Consider a regular n -gon Q centered at a fixed point o , and its rotation Q' by π/n about o . Let θ be a direction parallel to one of the edges of the convex hull of Q and Q' . It is easy to see that Q and Q' have the same parallel X-rays in the direction θ . This example arises in many papers, mainly related to Geometric and Discrete Tomography. We refer the interested reader to [7, 13].

Gardner and McMullen proved in [6] that *convex bodies are determined by X-rays taken in any set of directions that is not a subset of the directions of the edges of an*

affinely regular polygon. Since the cross ratio of any four directions of the edges of a regular polygon is an algebraic number, any set of four directions with a transcendental cross ratio uniquely determines a convex body by means of the corresponding X-rays.

Volčič [21] and Longinetti [16] show that the reconstruction of H is well posed when the set of directions guarantees uniqueness. Roughly speaking, if we know all the X-rays of H in such directions, and these X-rays contain errors ε as small as we want, then the corresponding reconstructions H_ε converge to H when ε goes to zero.

Now, consider the case when a finite number n of X-rays of H are exactly known, but the directions are determined up to an error δ . The error δ has to be small enough to distinguish the given n directions among them. For any positive δ , we cannot distinguish the set S of the given directions from the sets of non-uniqueness in the Gardner–McMullen theorem. Therefore, the results of well-posedness proved by Volčič cannot be used here. In [14] the following result is proved:

$$|K \Delta K'| \leq l^2 (8n)^{-1} \tan \frac{\pi}{n}, \quad (74)$$

where l is the length of the boundary of $K \cap K'$ and $K \Delta K'$ is the symmetric difference. Inequality (74) can be seen as a stability result and is optimal not only in the order but also in the constant, since equality holds if and only if the n directions, K and K' are chosen as in the above example. Inequality (74) is not affine invariant, while Hammer's problem is. An affine-invariant inequality in which l^2 is replaced by $|K \cap K'|$ is proved in [15] for sets of three directions. We show here that (55) can be used to generalize this result.

Let R be a connected component of $K \Delta K'$. For every direction $\theta \in S$, let θR be the connected component of $K \Delta K'$ different from R with the same X-ray of R in the direction θ . Let

$$W(R) = \bigcup_{h \in \mathbb{N}, \theta_{i_j} \in S} \theta_{i_h} \cdots \theta_{i_1} R.$$

In [14, Proposition 2] (see also [7, Lemma 1.2.6]) it is proved that $W(R)$ consists of a finite number h of components and they are at least $2n$.

Suppose that $K \Delta K' = W(R)$. Since all components in $W(R)$ have the same area, Corollary 5.2 yields

$$\frac{|W(R)|}{|K \cap K'|} \leq h \cdot \max_{P \in \mathcal{P}_h} G(P).$$

From (55) we obtain the explicit bound $2 \sin^2(\pi/h) / \cos(2\pi/h)$, a decreasing function of h . Since $h \geq 2n$, we get the following result.

Theorem 6.1 *If K and K' are two planar convex bodies with the same X-rays in n different directions ($n \geq 3$), and such that $K \Delta K'$ consists of a finite number of connected components of equal area, then*

$$|K \Delta K'| \leq |K \cap K'| \frac{1 - \cos(\pi/n)}{\cos(\pi/n)}. \quad (75)$$

Equality holds if and only if, up to an affine transformation, the directions are equally spaced, K and K' are congruent regular n -gons, and K' is K rotated by π/n about its center.

We remark that (75) is stronger than (74), via the classical isoperimetric inequality for n -gons.

In a forthcoming paper by Dulio, Longinetti, Peri and Venturi [4], Theorem 6.1 will be presented without the assumption on the connected components of $K \Delta K'$. There, also an improvement of (75) is given by using more information about the set of directions, as, for example, the cross ratio of four directions.

References

1. Barlotti, A.: Una proprietà degli n -agoni che si ottengono trasformando in una affinità un n -agone regolare. Boll. Unione Mat. Ital. **510**(3), 96–98 (1955)
2. Bianchi, G., Longinetti, M.: Reconstructing plane sets from projections. Discrete Comput. Geom. **5**, 223–242 (1990)
3. Coxeter, H.S.M.: Affinely regular polygons. Abh. Math. Sem. Univ. Hamburg **62**, 249–253 (1992)
4. Dulio, P., Longinetti, M., Peri, C., Venturi, A.: Sharp affine stability estimates for Hammer's problems. Adv. Appl. Math. (2007). doi:[10.1016/j.amm.2007.06.001](https://doi.org/10.1016/j.amm.2007.06.001)
5. Fisher, J.C., Jamison, R.E.: Properties of affinely regular polygons. Geom. Dedicata **69**, 241–259 (1998)
6. Gardner, R.J., McMullen, P.: On Hammer's X-ray problem. J. Lond. Math. Soc. **21**, 171–175 (1980)
7. Gardner, R.J.: Geometric Tomography. Cambridge University Press, New York (1995)
8. Giering, O.: Bestimmung von Eibeirichen und Eikörpern durch Steiner-Symmetrisierungen. Sber. Bayer. Akad. Wiss. München, Math.-Nat. Kl., 225–253 (1962)
9. Gruber, P.M.: Aspects of approximation of convex bodies. In: Gruber, P.M., Wills, J.M. (eds.) Handbook of Convex Geometry, pp. 319–345. North-Holland, Amsterdam (1993)
10. Hammer, P.C.: Problem 2. In: Proc. of Symposium in Pure Mathematics. Convexity, vol. VII. American Mathematical Society, Providence (1963)
11. Harel, G., Rabin, J.M.: Polygons whose vertex triangles have equal area. Am. Math. Mon. **110**, 606–619 (2003)
12. John, F.: Extremum problems with inequalities as subsidiary conditions. In: Courant Anniversary Volume, pp. 187–204. Interscience, New York (1948)
13. Kuba, A., Herman, G.T.: Discrete tomography: a historical overview. In: Herman, G.T., Kuba, A. (eds.) Discrete Tomography, pp. 3–34. Birkhäuser, Boston (1999)
14. Longinetti, M.: An isoperimetric inequality for convex polygons and convex sets with the same symmetrals. Geom. Dedicata **20**, 27–41 (1986)
15. Longinetti, M.: Una proprietà di massimo dei poligoni affinementemente regolari. Rend. Circ. Mat. Palermo **24**, 448–459 (1985)
16. Longinetti, M.: Some questions of stability in the reconstruction of plane convex bodies from projections. Inverse Probl. **1**, 87–97 (1985)
17. Lopez, M.A., Reisner, S.: Efficient approximation of convex polygons. Int. J. Comput. Geom. Appl. **10**, 445–452 (2000)
18. Ludwig, M.: A characterization of affine length and asymptotic approximation of convex discs. Abh. Math. Semin. Univ. Hamburg **169**, 75–88 (1999)
19. Reisner, S., Schütt, C., Werner, E.: Dropping a vertex or a facet from a convex polytope. Forum Math. **13**, 359–378 (2001)
20. Renyi, A., Sulanke, R.: Über die konvexe Hülle von n zufällig gewählten Punkten. Z. Wahrscheinlichkeitsthe. Verw. Geb. **2**, 75–84 (1963)
21. Volčič, A.: Well-posedness of the Gardner–McMullen reconstruction problem. In: Proceedings of Conference on Measure Theory, Oberwolfach, 1983. Lecture Notes in Mathematics, vol. 1089, pp. 199–210. Springer, Berlin (1984)