A Note on Projecting the Cubic Lattice

N.J.A. Sloane · Vinay A. Vaishampayan ·
Sueli I.R. Costa

Received: 7 April 2010 / Revised: 16 July 2010 / Accepted: 27 July 2010 /
Published online: 20 August 2010
© Springer Science+Business Media, LLC 2010

Abstract It is shown that, given any $(n - 1)$ -dimensional lattice Λ , there is a vector $v \in \mathbb{Z}^n$ such that the orthogonal projection of $\mathbb{Z}^n$ onto $v^\perp$ is, up to a similarity, arbitrarily close to Λ . The problem arises in attempting to find the largest cylinder anchored at two points of $\mathbb{Z}^n$ and containing no other points of $\mathbb{Z}^n$.

Keywords Projections · Shadows · Dense packings

1 Introduction

Let $\mathbb{Z}^n$ denote the cubic lattice with basis $e_1 := (1, 0, \dots, 0), \dots, e_n := (0, 0, \dots, 1)$. If we project $\mathbb{Z}^n$ onto the $(n - 1)$ -dimensional subspace

$$v^\perp := \{x \in \mathbb{R}^n : x \cdot v = 0\}$$

perpendicular to a vector $v \in \mathbb{Z}^n$, we obtain an $(n - 1)$ -dimensional lattice that we denote by Λ_v . We will show that, given any $(n - 1)$ -dimensional lattice Λ , we can choose $v \in \mathbb{Z}^n$ so that Λ_v is arbitrarily close to a lattice that is geometrically similar to Λ . More precisely, we will establish:

Theorem 1 *Let Λ be an $(n - 1)$ -dimensional lattice with Gram matrix A (with respect to some basis for $\mathbb{R}^{n-1}$). For any $\epsilon > 0$, there exist a nonzero vector $v \in \mathbb{Z}^n$,*

N.J.A. Sloane (✉) · V.A. Vaishampayan
AT&T Shannon Labs, 180 Park Avenue, Florham Park, NJ 07932-0971, USA
e-mail: njas@research.att.com

V.A. Vaishampayan
e-mail: vinay@research.att.com

S.I.R. Costa
University of Campinas, Campinas, SP 13083-970, Brazil
e-mail: sueli@ime.unicamp.br

a basis B for the $(n-1)$ -dimensional lattice Λ_v and a number c such that if A_v denotes the Gram matrix of B , then

$$\|A - cA_v\|_\infty < \epsilon. \quad (1)$$

The theorem is at first surprising, since A has $\binom{n+1}{2}$ degrees of freedom, whereas v has only n degrees of freedom (for the explanation see the remark following the proof of Theorem 2).

The problem arises from a question in communication theory (see Sect. 5), which calls for projections Λ_v with high packing density. Since both the determinant and minimal norm of a lattice are continuous functions of the entries in the Gram matrix, so is the packing density.¹ The theorem therefore implies that the packing density of Λ_v can be made arbitrarily close to that of Λ . So if we know a dense lattice in $\mathbb{R}^{n-1}$, we can find projections that converge to it in density.

Remark We know (see for example [2, Cor. 8]) that if Λ is a classically integral $(n-1)$ -dimensional lattice then Λ can be embedded in some odd unimodular lattice K of dimension $k \leq n+2$, although for $n \geq 7$ K need not be $\mathbb{Z}^k$. In any case this does not imply that Λ can be recovered as a projection of K .

Notation Λ^* denotes the dual lattice to Λ , A^{tr} is the transpose of A , and $\|A\|_\infty = \max_{i,j} |A_{i,j}|$. Our vectors are row vectors. For undefined terms from lattice theory see [3].

2 Proof of Theorem 1

We begin with some preliminary remarks about the projection lattice Λ_v and its dual Λ_v^* . For simplicity we will only consider projections that use vectors of the form $v = (1, v_1, v_2, \dots, v_{n-1}) \in \mathbb{Z}^n$. Let $\hat{v} := (v_1, v_2, \dots, v_{n-1})$, $M := \|v\|^2 = 1 + \sum v_i^2$.

The matrix that orthogonally projects $\mathbb{R}^n$ onto $v^\perp$ is $P := I_n - \frac{1}{M}v^{\text{tr}}v$. As a generator matrix G for $\mathbb{Z}^n$ (expressed in terms of $e_1, \dots, e_n$) we take I_n with its first row replaced by v . Let G_v be obtained by omitting the first (zero) row of GP . Then G_v is an $(n-1) \times n$ generator matrix for the projection lattice Λ_v , and $A_v := G_v G_v^{\text{tr}} = I_{n-1} - \frac{1}{M}\hat{v}^{\text{tr}}\hat{v}$ is its Gram matrix, with $\det \Lambda_v = \det A_v = \frac{1}{M}$.

It is often easier to work with the dual lattice Λ_v^* . This is the intersection of $\mathbb{Z}^n$ with the subspace $v^\perp$, and has generator matrix

$$\begin{bmatrix} -v_1 & 1 & 0 & \dots & 0 \\ -v_2 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -v_{n-1} & 0 & 0 & \dots & 1 \end{bmatrix}, \quad (2)$$

Gram matrix $A_v^* = I_{n-1} + \hat{v}^{\text{tr}}\hat{v}$, and determinant M .

¹The minimal norm μ of a d -dimensional lattice with Gram matrix A is the minimum over all $z \in \mathbb{Z}^d$, $z \neq 0$, of the quadratic form zAz^{tr} . It is enough to consider the finite set of z in some ball around the origin. For a given $z \neq 0$, zAz^{tr} is a continuous function of the entries of A , and since the minimum of a finite set of continuous functions is continuous, μ is a continuous function of the entries of A .

If a sequence of matrices T_i converges in the $\|\cdot\|_\infty$ norm to a positive-definite matrix T , then T_i^{-1} converges to T^{-1} . So the following theorem is equivalent to Theorem 1.

Theorem 2 *Let Λ be an $(n-1)$ -dimensional lattice with Gram matrix A (with respect to some basis for $\mathbb{R}^{n-1}$). For any $\epsilon > 0$, there exist a nonzero vector $v \in \mathbb{Z}^n$, a basis B for the $(n-1)$ -dimensional lattice Λ_v^* and a number c such that if A_v^* denotes the Gram matrix of B , then*

$$\|A - cA_v^*\|_\infty < \epsilon. \quad (3)$$

Proof of Theorem 2 We may write $A = LL^T$ where $L = [L_{i,j}]$ is an $(n-1) \times (n-1)$ lower triangular matrix. For $w = 1, 2, \dots$ let us form the $(n-1) \times n$ matrix

$$\begin{aligned} L_w &:= -\begin{bmatrix} \lfloor wL \rfloor & \mathbf{0} \end{bmatrix} + \begin{bmatrix} \mathbf{0} & I_{n-1} \end{bmatrix} \\ &= \begin{bmatrix} -\lfloor wL_{1,1} \rfloor & 1 & 0 & \dots & 0 & 0 \\ -\lfloor wL_{2,1} \rfloor & -\lfloor wL_{2,2} \rfloor & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ -\lfloor wL_{n-1,1} \rfloor & -\lfloor wL_{n-1,2} \rfloor & -\lfloor wL_{n-1,3} \rfloor & \dots & -\lfloor wL_{n-1,n-1} \rfloor & 1 \end{bmatrix}, \end{aligned} \quad (4)$$

where $\mathbf{0}$ denotes a column of $n-1$ zeros. We call L_w a “lifted” version of L .

We apply elementary row operations to L_w so as to put it in the form

$$\tilde{L}_w := \begin{bmatrix} -v_1 & 1 & 0 & \dots & 0 & 0 \\ -v_2 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ -v_{n-2} & 0 & 0 & \dots & 1 & 0 \\ -v_{n-1} & 0 & 0 & \dots & 0 & 1 \end{bmatrix}, \quad (5)$$

and take $v = (1, v_1, \dots, v_{n-1})$. Then Λ_v^* has generator matrix $\tilde{L}_w$. But $\tilde{L}_w$ and L_w generate the same lattice. It follows that Λ_v^* has a Gram matrix

$$A_v^* = L_w L_w^T = w^2 A + B = w^2 \left(A + \frac{1}{w^2} B \right), \quad (6)$$

using (4), where the entries in B are of order $O(w)$ as $w \rightarrow \infty$. This implies (3) (with $c = 1/w^2$). $\square$

Remark The apparent paradox mentioned in Sect. 1 is explained by the fact that we use $\binom{n}{2}$ degrees of freedom in going from (4) to (5).

3 Examples

3.1 The Lattice $2\mathbb{Z} \oplus \mathbb{Z}$

We start with a concrete example. If we take $v = (1, 1, 0)$ then Λ_v has Gram matrix $\frac{1}{2} \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$, and is geometrically similar to $\sqrt{2}\mathbb{Z} \oplus \mathbb{Z}$. Similarly $v = (1, 1, 1)$ produces the hexagonal (or A_2) lattice, and in general $v = (1, 1, \dots, 1)$ gives A_{n-1} . On the other hand, there is no $v = (1, v_1, v_2) \in \mathbb{Z}^3$ such that Λ_v is geometrically similar to $2\mathbb{Z} \oplus \mathbb{Z}$ (see Proposition 3). However, we can find projections which converge to a lattice that is geometrically similar to $2\mathbb{Z} \oplus \mathbb{Z}$. Since any two-dimensional lattice is geometrically similar to its dual, we can apply Theorem 2 with $\Lambda = 2\mathbb{Z} \oplus \mathbb{Z}$. Then $L = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$, the lifted generator matrix is $L_w = \begin{bmatrix} -2w & 1 & 0 \\ 0 & -w & 1 \end{bmatrix}$, $\tilde{L}_w = \begin{bmatrix} -2w & 1 & 0 \\ -2w^2 & 0 & 1 \end{bmatrix}$, $v = (1, 2w, 2w^2)$, and a Gram matrix for Λ_v^* is $I_2 + \tilde{v}^{\text{tr}} \tilde{v} = \begin{bmatrix} 4w^2+1 & 4w^3 \\ 4w^3 & 4w^4+1 \end{bmatrix}$. If we subtract w times the first generator from the second, this becomes

$$\begin{bmatrix} 4w^2+1 & -w \\ -w & w^2+1 \end{bmatrix} = w^2 \begin{bmatrix} 4+1/w^2 & -1/w \\ -1/w & 1+1/w^2 \end{bmatrix},$$

which converges to $w^2 \begin{bmatrix} 4 & 0 \\ 0 & 1 \end{bmatrix}$ as $w \rightarrow \infty$.

Proposition 3 *There is no vector $v = (1, a, b) \in \mathbb{Z}^3$ such that Λ_v^* is geometrically similar to $2\mathbb{Z} \oplus \mathbb{Z}$.*

Proof From (2), Λ_v^* has Gram matrix $A := \begin{bmatrix} a^2+1 & ab \\ ab & b^2+1 \end{bmatrix}$. If Λ_v^* is geometrically similar to $2\mathbb{Z} \oplus \mathbb{Z}$ then there is a matrix $T := \begin{bmatrix} r & s \\ t & u \end{bmatrix} \in SL_2(\mathbb{Z})$ and $\lambda \in \mathbb{R}$ such that

$$A = \lambda T \begin{bmatrix} 4 & 0 \\ 0 & 1 \end{bmatrix} T^{\text{tr}} = \lambda \begin{bmatrix} 4r^2+s^2 & 4rt+su \\ 4rt+su & 4t^2+u^2 \end{bmatrix}.$$

This implies $\lambda \in \mathbb{Q}$, and taking the determinant and trace of both sides we obtain $a^2+b^2+1 = 4\lambda^2$, $a^2+b^2+2 = 4\lambda^2+1 = \lambda\sigma$, where $\sigma := 4r^2+4t^2+s^2+u^2 \in \mathbb{Z}$. Hence the discriminant of the quadratic for λ , $\sigma^2 - 16$, is a perfect square, so $\sigma = 4$ or 5 , $\lambda = \frac{1}{2}$, $\frac{1}{4}$ or 1 , $a^2+b^2+1 = 1$, $\frac{1}{4}$ or 3 , none of which are possible. $\square$

3.2 The Lattice 5_1

For an example where the floor operations in (4) are actually needed, consider the lattice Λ with Gram matrix $\begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix}$ and determinant 5 (this is the lattice 5_1 in the notation of [1]). Again there is no $v = (1, v_1, v_2) \in \mathbb{Z}^3$ such that Λ_v is geometrically similar to Λ . We take $L = \begin{bmatrix} \sqrt{3} & 0 \\ \frac{1}{\sqrt{3}} & \sqrt{\frac{5}{3}} \end{bmatrix}$, and find that

$$v = (1, \lfloor \sqrt{3}w \rfloor, \lfloor \sqrt{3}w \rfloor \lfloor \sqrt{5/3}w \rfloor + \lfloor w/\sqrt{3} \rfloor).$$

3.3 The Lattice D_m , $m \geq 3$

As generator matrix for D_m^* we take ([3, p. 120])

$$\begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \\ 1/2 & 1/2 & 1/2 & \dots & 1/2 & 1/2 \end{bmatrix}. \quad (7)$$

We set $w = 2t$, $t \in \mathbb{Z}$, and obtain

$$v = \left(1, 2t, (2t)^2, \dots, (2t)^{m-1}, t \frac{(2t)^m - 1}{2t - 1} \right).$$

In particular, when $m = 3$, we have

$$v = (1, 2t, 4t^2, 4t^3 + 2t^2 + t), \quad (8)$$

for which Λ_v^* converges to the body-centered cubic lattice D_3^* and Λ_v to the face-centered cubic lattice D_3 .

3.4 The Lattice E_8

Using the generator matrix given in [3, p. 121], we find that $v = (1, v_1, v_2, \dots, v_8)$ is given by $v_1 = 2w$, $v_2 = 2w^2 - w$, $v_i = w(v_{i-1} - v_{i-2})$ for $i = 3, 4, \dots, 7$, and $v_8 = (w/2)(1 + \sum_{i=1}^7 v_i)$, where w is even.

3.5 The Leech Lattice

Using [3, Fig. 4.12], we find that $v = (1, v_1, v_2, \dots, v_{24})$ is given by

$$\begin{aligned} v_1 &= 8w, \\ v_i &= 4w(v_{i-1} + 1), \quad \text{for } i = 2, \dots, 7, 9, 10, 11, 13, 17, \\ v_i &= 2w \left(\sum_{j \in S_i} v_j + 1 \right), \quad \text{for } i = 8, 12, 14, 15, 16, 18, 19, 20, \\ v_i &= 2w \sum_{j \in S_i} v_j, \quad \text{for } i = 21, 22, 23, \\ v_{24} &= w \left(\sum_{i=1}^{23} v_i - 3 \right), \end{aligned} \quad (9)$$

where $S_8 = \{1, 2, \dots, 7\}$, $S_{12} = \{1, 2, 3, 8, 9, 10, 11\}$, $S_{14} = \{1, 4, 5, 8, 9, 12, 13\}$, $S_{15} = \{2k, 1 \leq k \leq 7\}$, $S_{16} = \{3, 4, 7, 8, 11, 12, 15\}$, $S_{18} = \{2, 4, 7, 8, 9, 16, 17\}$, $S_{19} = \{3, 4, 5, 8, 10, 16, 18\}$, $S_{20} = \{1, 4, 6, 8, 11, 16, 19\}$, $S_{21} = \{1, 2, 3, 4, 8, 12, 16, 20\}$, $S_{22} = \{8, 9, 12, 13, 16, 17, 20, 21\}$, $S_{23} = \{2k, k = 4, 5, \dots, 11\}$.

4 Faster Convergence

The construction in Theorem 2 produces a vector v of length $\|v\| = O(w^n)$, while from (6) we have $\|A - \frac{1}{w^2}A_v^*\|_\infty = O(\frac{1}{w}) = O(\frac{1}{\|v\|^{1/n}})$. It is sometimes possible to obtain a faster rate of convergence. Suppose A is D_3 , and instead of (7) let us take the following generator matrix for D_3^* :

$$L := \begin{bmatrix} -1 & -1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$$

Let

$$\begin{aligned} L_w &= \begin{bmatrix} w-1 & w+1 & -w & 0 \\ -w-1 & w & -w+1 & 0 \\ -w & -w & -w & 1 \end{bmatrix} \\ &= [-wL \quad \mathbf{0}] + [\mathbf{0} \quad I_3] + \begin{bmatrix} -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \end{aligned} \quad (10)$$

with

$$A_v^* = L_w L_w^{\text{tr}} = \begin{bmatrix} 3w^2+2 & w^2+1 & -w^2 \\ w^2+1 & 3w^2+2 & w^2 \\ -w^2 & w^2 & 3w^2+1 \end{bmatrix}. \quad (11)$$

The last matrix in (10) is chosen so that there are no terms of order w in (11). Let H_w denote the 3×3 matrix formed by the last three columns of L_w , and define v_1, v_2, v_3 by

$$H_w^{-1}L_w = \begin{bmatrix} -v_1 & 1 & 0 & 0 \\ -v_2 & 0 & 1 & 0 \\ -v_3 & 0 & 0 & 1 \end{bmatrix}.$$

Then

$$v = (1, v_1, v_2, v_3) = (1, 2w^2 - w + 1, 2w^2 + w + 1, 4w^3 + 3w) \quad (12)$$

has $\|v\| = O(w^3)$, and now $\|A - \frac{1}{w^2}A_v^*\|_\infty = O(\frac{1}{\|v\|^{2/3}})$, which is a faster convergence than we found in Sect. 3.3. We do not know if similar speedups are always possible. Incidentally, we first found (12)—before Theorem 1 was proved—by a combination of computer search and guesswork.

5 The Fat Strut Problem

The problem studied in this paper arose when constructing codes for a certain analog communication channel [6]. The codes require that one finds a curved tube in the sphere S^{2n-1} which does not intersect itself, has a specified length and as large a

volume as possible. The method used in [6] is based on finding a vector $v \in \mathbb{Z}^n$ with a specified value of $\|v\|$, such that there is a cylinder of large volume with axis $\overrightarrow{0v}$ which contains no points of $\mathbb{Z}^n$ other than 0 and v . The cross-section of the cylinder is an $(n - 1)$ -dimensional ball, and 0 and v are the centers of the two end-faces. The radius of the cylinder is taken to be as large as possible subject to the condition that the interior contains no point of $\mathbb{Z}^n$. The problem is to choose v , for a given length $\|v\|$, so that the volume of the resulting cylinder is maximized. We call a cylinder which achieves the maximal volume a *fat strut*.

A fat strut has the property that the projection of the cylinder onto $v^\perp$ does not contain the image of any nonzero point of $\mathbb{Z}^n$. The radius of the cylinder is therefore equal to the radius of the largest $(n - 1)$ -dimensional sphere in the projection lattice Λ_v which contains no nonzero point of Λ_v . In other words, finding a fat strut for a given length $\|v\|$ is equivalent to maximizing the density of the projection lattice Λ_v .

It is worth contrasting the fat strut problem with the result of [4] and [5] that for any lattice sphere packing in dimension three or higher there is always an *infinite* cylinder of nonzero radius (obviously not passing through the origin) which does not touch any of the spheres.

Acknowledgement We thank the referee for some very helpful comments.

References

1. Conway, J.H., Sloane, N.J.A.: Low-dimensional lattices I: Quadratic forms of small determinant. Proc. R. Soc. Lond. A **418**, 17–41 (1988)
2. Conway, J.H., Sloane, N.J.A.: Low-dimensional lattices V: Integral coordinates for integral lattices. Proc. R. Soc. Lond. A **425**, 211–232 (1989)
3. Conway, J.H., Sloane, N.J.A.: Sphere Packings, Lattices and Groups, 3rd edn. Springer, New York (1998)
4. Heppes, A.: Ein satz über gitterförmige kugelpackungen. Ann. Univ. Sci. Budapest. Eötvös Sect. Math. **3–4**, 89–90 (1960/1961)
5. Horváth, J.: Über die durchsichtigkeit gitterförmiger kugelpackungen. Stud. Sci. Math. Hung. **5**, 421–426 (1970)
6. Vaishampayan, V.A., Costa, S.I.R.: Curves on a sphere, shift-map dynamics, and error control for continuous alphabet sources. IEEE Trans. Inf. Theory **49**, 1658–1672 (2003)