

Neighborly Cubical Polytopes*

M. Joswig and G. M. Ziegler

Fachbereich Mathematik, MA 7-1, Technische Universität Berlin,
10623 Berlin, Germany
{joswig,ziegler}@math.tu-berlin.de

Abstract. Neighborly cubical polytopes exist: for any $n \geq d \geq 2r + 2$, there is a cubical convex d -polytope C_d^n whose r -skeleton is combinatorially equivalent to that of the n -dimensional cube. This solves a problem of Babson, Billera, and Chan.

Kalai conjectured that the boundary ∂C_d^n of a neighborly cubical polytope C_d^n maximizes the f -vector among all cubical $(d - 1)$ -spheres with 2^n vertices. While we show that this is true for polytopal spheres if $n \leq d + 1$, we also give a counterexample for $d = 4$ and $n = 6$.

Further, the existence of neighborly cubical polytopes shows that the graph of the n -dimensional cube, where $n \geq 5$, is “dimensionally ambiguous” in the sense of Grünbaum. We also show that the graph of the 5-cube is “strongly 4-ambiguous.”

In the special case $d = 4$, neighborly cubical polytopes have $f_3 = (f_0/4) \log_2(f_0/4)$ vertices, so the facet–vertex ratio f_3/f_0 is not bounded; this solves a problem of Kalai, Perles, and Stanley studied by Jockusch.

1. Introduction

In Chapter 12 of his famous book [12] Grünbaum discusses the concept of k -equivalence of polytopes. A d -polytope P is k -equivalent to a d' -polytope P' if the k -skeleta of P and P' are combinatorially equivalent. An interesting case occurs when $\dim P \neq \dim P'$. In this situation Grünbaum calls the k -skeleton \mathcal{S} of either polytope *dimensionally ambiguous*. The k -skeleton \mathcal{S} of a d -polytope is called *strongly d -ambiguous* if there is another d -polytope Q , not combinatorially equivalent to P , whose k -skeleton is also combinatorially equivalent to \mathcal{S} . From the existence of neighborly polytopes, such as the cyclic polytopes, it follows that the $(\lfloor d/2 \rfloor - 1)$ -skeleton of the d -simplex is dimensionally ambiguous for $d \geq 5$.

* This work was supported by a DFG Gerhard-Hess-Forschungsförderungspreis (Zi 475/1-2) and by the German Israeli Foundation (G.I.F.) Grant I-0309-146.06/93.

Motivated by a problem of Kalai [18, Problem 3.4(iv)*], we first constructed a 4-polytope with the graph of the 5-cube, thus showing that the graph of the five-dimensional cube is dimensionally ambiguous. This turns out to be a special case of a d -polytope with the r -skeleton of the $(d + 1)$ -cube, for $r = \lfloor d/2 \rfloor - 1$. We describe two constructions in Section 4. The resulting polytopes are cubical, and hence they appear in the combinatorial classification of the cubical polytopes with 2^{d+1} vertices due to Blind and Blind [6], [7].

More generally, in this paper we construct *neighborly cubical polytopes* in the sense of Babson et al. [3]: for every $n \geq d \geq 2$ there is a cubical d -polytope with the r -skeleton of the n -cube, for $r = \lfloor d/2 \rfloor - 1$. In particular, this yields a 4-polytope with the graph of the n -cube for every $n \geq 4$. The neighborly cubical polytopes C_d^n are constructed as linear projections of “deformed” cubes, see Section 6. The combinatorics of the moment curve is involved in an essential way; this is reminiscent of the construction of neighborly (simplicial) polytopes. We also give an explicit combinatorial description of C_d^n which can be seen as a “cubical Gale evenness criterion.”

A result of Grünbaum [12, Theorem 12.2.1] implies that no d -polytope can have the r -skeleton of the n -cube for $r > \lfloor d/2 \rfloor - 1$ with $n > d$; this is reviewed in Section 2.

Our construction specializes to a known phenomenon for $d = 2$: there are n -cubes whose two-dimensional “shadows” have 2^n vertices (“projections that preserve the 0-skeleton”). These were first constructed by Murty [15] and more explicitly by Goldfarb [11]. They amount to linear programs for which the shadow boundary pivot rule takes an exponential number of steps. In [2] the Goldfarb cubes were interpreted as a special case of a construction of “deformed cubes,” and indeed the neighborly cubical polytopes constructed in this paper are projections of deformed cubes $C^n(\varepsilon)$ as well.

An interesting new phenomenon occurs in the case $d = 4$: Jockusch [13] had constructed examples of cubical 4-polytopes for which the facet/vertex ratio f_3/f_0 was higher than previously expected, namely, arbitrarily close to $5/4$. The neighborly cubical polytopes show that indeed the ratio f_3/f_0 is not bounded for cubical 4-polytopes: these polytopes have $f_0 = 2^n$ vertices and $f_3 = (n-2)2^{n-2} = f_0 \log(f_0/4)/4 = \Theta(f_0 \log f_0)$ facets.

In particular, we get that a 4-polytope with f_0 vertices can have more than $O(f_0)$ nonsimplicial facets, which solves a problem of Erickson [9]. Also, any Schlegel diagram of a neighborly 4-polytope is a cubical complex (a “hexahedral mesh” in Computational Geometry terminology) with $f_0 = 2^n$ vertices and $f_3 - 1 = f_0 \log(f_0/4)/4 - 1 = \Theta(f_0 \log f_0)$ cubical facets that subdivides a cube. This answers a question of Bern and Eppstein [8], who had asked whether a hexahedral mesh could have more than linearly many facets.

Kalai’s cubical upper bound conjecture [3, Conjecture 4.2] claimed that among all cubical $(d - 1)$ -spheres with 2^n vertices, the boundaries of cubical neighborly polytopes simultaneously maximize all components of the f -vector. In Section 7 we give a counterexample, but we prove the claim in the special case $n = d + 1$ for *polytopal* $(d - 1)$ -spheres.

A major part of the research on cubical polytopes is guided by the comparison with simplicial polytopes. In this paper we extend this analogy by providing cubical analogs to the cyclic (neighborly simplicial) polytopes. However, we also offer a surprising instance where the cubical case differs from the simplicial case: While even-dimensional

neighborly polytopes are always simplicial, it is not true that a 4-polytope with the graph of the 5-cube must necessarily be cubical—we construct an explicit example in Section 4.4. Together with the existence of the unique cubical 4-polytope with the graph of the 5-cube this implies that the graph of the 5-cube is strongly 4-ambiguous.

2. Neighborly Cubical Polytopes

We refer to [12] and [18] for general introductions to polytopes and polytopal complexes. Two polytopes or polytopal complexes are *combinatorially equivalent* if their posets of faces are isomorphic. In the following, a *d-cube* is any polytope that is combinatorially isomorphic to the standard d -cube $C_d = [0, 1]^d \subseteq \mathbb{R}^d$. A *combinatorial cube* is such a d -cube, for any d . A *cubical polytope* is any polytope all of whose proper faces are combinatorial cubes. The *k-skeleton* of a polytope is the polytopal complex given by all faces of dimension k or less. We say that P *has the k-skeleton of a cube* if its k -skeleton is combinatorially equivalent to that of a combinatorial cube. A *neighborly cubical polytope* is a cubical d -polytope (with 2^n vertices for some $n \geq d$) which has the $(\lfloor d/2 \rfloor - 1)$ -skeleton of a cube. This notion was introduced in [3], where neighborly cubical spheres were constructed, and the question about the existence of neighborly cubical polytopes was raised. We start by explaining the choice of parameters in this definition.

Proposition 1 (Characterization of Cubes [6]). *Any cubical d -polytope has at least 2^d vertices. If a cubical d -polytope has exactly 2^d vertices, then it is a combinatorial d -cube.*

Corollary 2. *If all the k -faces of a d -polytope have 2^k vertices, for all $0 \leq k \leq d - 1$, then the polytope is cubical. If in addition the polytope has 2^d vertices, then it is a combinatorial cube.*

It is well known that the f -vector of a cubical polytope is subject to restrictions that are similar to the Dehn–Sommerville equations for simplicial/simple polytopes.

Proposition 3 (Cubical Dehn–Sommerville Equations [12, Theorem 9.4.1]). *Let (f_0, \dots, f_{d-1}) be the f -vector of a cubical d -polytope. Then, for $0 \leq k \leq d - 2$,*

$$\sum_{i=k}^{d-1} (-1)^i 2^{i-k} \binom{i}{k} f_i = (-1)^{d-1} f_k.$$

Lemma 3 (Simple Cubical Polytopes [18, Exercise 0.1, p. 23]). *Every simple cubical d -polytope with $d > 2$ is a d -cube. (Every 2-polytope is simple and cubical.)*

Proposition 4. *If a d -polytope P has the r -skeleton of the n -dimensional cube, then all k -faces of P are cubes for $k \leq 2r$.*

Proof. Let F be a k -face of P and $k \leq 2r$. By induction on k we can assume that F is cubical. If F is simple, then F is a cube by Lemma 3. Thus assume that F has a vertex v of degree $k' > k$. Let a_1, \dots, a_{k+1} be $k+1$ distinct vectors such that $v + a_i$ is a neighbor of v in F . As $\dim F = k$, the vectors a_1, \dots, a_{k+1} are linearly dependent, i.e., we can choose $\lfloor (k+1)/2 \rfloor$ vectors among them which do not span a proper face of F . However, $\lfloor (k+1)/2 \rfloor \leq \lfloor (2r+1)/2 \rfloor = r$. This contradicts the assumption that P has the r -skeleton of a cube. \square

We note here that in the simplicial case more is true: If a d -polytope P has the r -skeleton of the n -dimensional simplex, then all k -faces of P are simplices for $k \leq 2r+1$. In particular, if a d -polytope P has the r -skeleton of the n -dimensional simplex for $r \geq \lceil d/2 \rceil$, then P is a d -simplex.

Corollary 5. *If a d -polytope P has the r -skeleton of the n -dimensional cube for $r \geq d/2$, then P is a d -cube.*

However, this is not good enough to establish the analogy to the simplicial case. What if P is a $2k$ -polytope with the $(k-1)$ -skeleton of the $(2k+1)$ -cube? By Proposition 4 all $(2k-2)$ -faces are cubes, but what about the $(2k-1)$ -faces, i.e., the facets? It turns out that the result of Proposition 4 is sharp in the sense that there are $2k$ -polytopes which have the $(k-1)$ -skeleton of a $(2k+1)$ -cube but which are *not* cubical: in Section 4.4 we present an example of a noncubical 4-polytope with the graph of the 5-cube.

The proof of the following is based on a theorem of van Kampen and Flores, see Theorem 11.1.3 and Section 11.3 of [12].

Proposition 6 [12, Theorem 11.2.1]. *Let P be a d -polytope with the r -skeleton of an n -polytope, with $n > d$. Then $r \leq \lfloor d/2 \rfloor - 1$.*

3. Projections

We discuss the effect of orthogonal projections on polytopes. Everything in this section is well known; it is included for the sake of completeness.

Let P be a full-dimensional polytope. A vector \mathbf{n} is *normal* with respect to a facet F of P if it is orthogonal to F and it points “to the outside,” that is, if the linear functional corresponding to \mathbf{n} and restricted to P attains its maximum at the points in F . A vector is called *normal* with respect to a face G if it is a positive linear combination of normal vectors of all facets of P containing G . Equivalently, the linear functional corresponding to a normal vector of G attains its maximum at the points in G . Obviously, every facet has a unique normal vector of length 1, while a face of higher codimension does not.

Consider an orthogonal projection π onto some proper affine subspace.

Lemma 7. *If the face G has a normal vector orthogonal to the direction of projection, then the image $\pi(G)$ is a face of the polytope $\pi(P)$. Conversely, if \overline{G} is a face of $\pi(P)$,*




Fig. 1. Orthogonal projection of a 2-polytope.

then the full preimage $\pi^{-1}(\overline{G})$ is a face of P with a normal vector orthogonal to the direction of projection.

The lemma above characterizes the “shadow boundary,” that is, it describes the faces that are mapped to the boundary of the projection. Note that this may include faces of G that are mapped to faces of lower dimension, see Fig. 1.

Lemma 8. *The restriction of π to a face G is injective if and only if G has a normal vector which is not orthogonal to the direction of projection.*

Combining the two lemmas above gives the following characterization.

Corollary 9. *A face G is mapped onto a face $\pi(G)$ of the same dimension if and only if it has a normal vector which is orthogonal to the direction of projection and another one which is not.*

We can also say something about the shape of the projection in this case.

Lemma 10. *A face G is mapped onto a face $\pi(G)$ of the same dimension if and only if the faces G and $\pi(G)$ are affinely isomorphic.*

4. The Case $n = d + 1$

4.1. First Construction

Let $d = 2r$ be even, $Q := [-1, +1]^r$, and

$$\tilde{P} := \text{conv}(Q \times 2Q \times \{-1\} \cup 2Q \times Q \times \{1\}) \subseteq \mathbb{R}^{d+1}.$$

This is clearly a combinatorial $(d + 1)$ -cube, with the complete linear description

$$\tilde{P} = \left\{ \begin{pmatrix} \mathbf{x} \\ x_{d+1} \end{pmatrix} \in \mathbb{R}^{d+1}: \begin{array}{l} -1 \leq x_{d+1} \leq 1, \\ \pm 2x_i \leq 3 - x_{d+1} \text{ for } 1 \leq i \leq r, \\ \pm 2x_i \leq 3 + x_{d+1} \text{ for } r < i \leq d \end{array} \right\}.$$

The projection $\pi: \mathbb{R}^{d+1} \rightarrow \mathbb{R}^d$ that deletes the last coordinate yields the d -polytope

$$P := \pi(\tilde{P}) = \text{conv}(Q \times 2Q \cup 2Q \times Q) \subseteq \mathbb{R}^d.$$

One Fourier–Motzkin elimination step [18, Section 1.2] shows that P can also be described in terms of its facets by

$$P = \{ \mathbf{x} \in \mathbb{R}^d: \begin{array}{l} \pm x_i \leq 2 \text{ for } 1 \leq i \leq 2r, \\ \pm x_i \pm x_j \leq 3 \text{ for } 1 \leq i \leq r < j \leq 2r. \end{array} \}.$$

This P is a cubical $2r$ -polytope with $f_0 = 2^{d+1}$ vertices and $f_{d-1} = 4r + (2r)^2 = d(d+2)$ facets.

It is also easy to see that P has the $(r - 1)$ -skeleton of a $(d + 1)$ -cube, either using the criteria of Section 3, or by direct verification from the complete description of the polytope in terms of facets and vertices.

This example of a polytope with the $d/2$ -skeleton of a cube is amazing because of its simplicity, and also because of its symmetry: it has a vertex-transitive symmetry group, and only two orbits of facets.

4.2. Second Construction

Blind and Blind completed a classification of the (combinatorial types) of cubical d -polytopes with 2^{d+1} vertices in [7]. From this classification, we derive below that, for even d , there is exactly one combinatorial type of a d -polytope with the $d/2$ -skeleton of the $(d + 1)$ -cube; for odd d , there are precisely two combinatorial types. The description of the polytopes given by Blind and Blind also implies the construction given above.

A (cubical) d -polytope P whose boundary complex ∂P is isomorphic to a subcomplex of the d -skeleton of some (higher-dimensional) cube is called *liftable*. We get all distinct combinatorial types of liftable d -polytopes (and $(d - 1)$ -spheres) with at most 2^{d+1} vertices as follows:

The cube C^{d+1} has $d + 1$ pairs of opposite facets F_i^\pm ($1 \leq i \leq d + 1$). For $k, m \geq 0$ and $l \geq 1$ with $k + l + m = d + 1$ let $B(k, l, m)$ be the cubical d -ball in the boundary of C^{d+1} formed by F_i^\pm ($1 \leq i \leq k$) and F_i^+ ($k + 1 \leq i \leq k + l$).

The combinatorial types $P(k, l, m)$ of liftable d -polytopes with at most 2^{d+1} vertices are given by the boundary complexes of the cubical balls $B(k, l, m)$. The number of

vertices of $P(k, l, m)$ equals 2^{d+1} if and only if $k, m \geq 1$. Note that $P(k, l, m)$ is combinatorially equivalent to $P(m, l, k)$; thus in the following let $k \geq m$. There are $\lfloor d^2/4 \rfloor$ suitable triples (k, l, m) with $k \geq m \geq 1$.

Theorem 11 (Classification for $n = d + 1$ [7, Theorem 3]). *For $d \geq 4$, all the combinatorial types of cubical d -polytopes with 2^{d+1} vertices are given by the liftable polytopes $P(k, l, m)$ with $k, l, m \geq 1, k+l+m = d+1$, and $k \geq m$, plus in addition the “two-fold nonlinearly capped” d -polytope P_{NLC}^d .*

A polytope $P(k, l, m)$ is r -neighborly if, for every r -face H of C^{d+1} , there is some facet of $B(k, l, m)$ containing H , and some facet of the complement $C^{d+1} \setminus B(k, l, m)$ containing H as well. Every r -face of C^{d+1} is contained in exactly $d + 1 - r$ facets of C^{d+1} . Thus $P(k, l, m)$ is r -neighborly if and only if $k + l \leq (d + 1 - r) - 1$ and $l + m \leq (d + 1 - r) - 1$, that is $r + 1 \leq m$ and $r + 1 \leq k$, where the first condition implies the second because of $k \geq m$.

Now we consider the case $r = \lfloor d/2 \rfloor - 1$.

Corollary 12. *If d is even, then $P(d/2, 1, d/2)$ is the unique $(d/2 - 1)$ -neighborly cubical polytope with 2^{d+1} vertices. If d is odd, then there are precisely two $((d - 1)/2 - 1)$ -neighborly cubical polytopes with 2^{d+1} vertices, namely, $P((d + 1)/2, 1, (d - 1)/2)$ and $P((d - 1)/2, 1, (d - 1)/2)$.*

Now we compute the f -vector of $P(k, l, m)$, denoted by $f(P(k, l, m)) = (f_0, \dots, f_{d-1})$. With the same reasoning as above, an i -face of C^{d+1} is a face of $P(k, l, m)$ if it lies in some facet of $B(k, l, m)$, and also in some facet not in $B(k, l, m)$. We deduce that

$$f_{d+1-i} = \binom{d+1}{i} 2^i - \sum_{j=0}^i \binom{l}{i-j} \left\{ \binom{k}{j} + \binom{m}{j} \right\} 2^j.$$

4.3. The Cubical Upper Bound Conjecture

From the analogy to the simplicial case one is tempted to expect that the neighborly cubical polytopes achieve equality for the cubical upper bound conjecture. For an explicit formula see Corollary 20.

Conjecture 13 (Cubical Upper Bound Conjecture (Kalai) [3, Conjecture 4.2]). *Let P be a cubical $(d - 1)$ -sphere with $f_0(P) = 2^n$ vertices. Then its number of facets is bounded by that of a neighborly cubical polytope/sphere C_d^n . Moreover,*

$$f_i(P) \leq f_i(C_d^n) \quad \text{for } 1 \leq i \leq d - 1.$$

There is an obvious upper bound for the f -vector of a cubical sphere. We owe this observation to both Kalai [14] and, independently, the anonymous referee.

Proposition 14. *Let P be a cubical $(d - 1)$ -sphere with $f_0(P) = n$ vertices. Then*

$$\sum_{k=0}^{d-1} f_k(P) \cdot 2^k \leq n^2.$$

Proof. Due to reflection symmetry each vertex of a cube has a unique antipode. Now let p be the number of antipodal pairs of vertices in all nonempty faces of P . Clearly, $p \leq n^2$. Moreover, p is the number of pairs (v, F) , where v is a vertex of P which is contained in the nonempty face F . The face F is a k -cube for some k , hence it has 2^k vertices. We obtain $p = \sum_{k=0}^{d-1} f_k(P) \cdot 2^k$. □

Theorem 15. *In the special case $n = d + 1$, Conjecture 13 is true if restricted to cubical polytopes. However, it is false for spheres even for $d = 4$ and $n = 6$.*

Proof. (1) The proof for polytopes relies on the classification of Theorem 11. Here we can disregard the “two-fold nonlinearly capped” d -polytope P_{NLC}^d , since it has the same f -vector as the “two-fold linearly capped” d -polytope, which is $P(d - 1, 1, 1)$, see Figure 1 of [7]. Thus our problem is to minimize, for fixed d and i , the function

$$\delta_i(k, l, m) = \sum_{j=0}^i \binom{l}{i-j} \left\{ \binom{k}{j} + \binom{m}{j} \right\} 2^j$$

subject to the restrictions $k, l, m \geq 1, k + l + m = d + 1$, and $k \geq m$. For this, we note the simple properties and inequalities

$$\begin{aligned} \delta_i(k, l, m) &\geq \delta_i(k - 1, l, m + 1) && \text{for } k > m, \\ \delta_i(k, l, m) &\geq \delta_i(k + 1, l - 2, m + 1) && \text{for } l \geq 2, \\ \delta_i(k, 2, m) &\geq \delta_i(k, 1, m + 1) && \text{for } k > m, \quad \text{and} \\ \delta_i(k, 2, k) &= \delta_i(k + 1, 1, k), \end{aligned}$$

from which the claim immediately follows.

(2) The second claim is verified via an explicit construction in Section 7. □

For even d , the first part of this result also follows from Babson et al. [3, Theorem 4.3], who used Adin’s “cubical h -vector” [1].

4.4. A Noncubical Polytope

Here we give an example of a noncubical 4-polytope with the same graph as the 5-cube. For this it is helpful to have yet another coordinate representation of the cubical 4-polytope with the graph of the 5-cube.

Let P be the polytope defined as the convex hull of the following 32 points in \mathbb{R}^4 .

Note that each row corresponds to four distinct points due to arbitrary variation of the signs.

- $(\pm 1 \quad \pm 1 \quad 1 \quad 1),$
- $(\pm 1 \quad \pm 1 \quad 4 \quad 1),$
- $(\pm 2 \quad \pm 2 \quad 3 \quad \frac{4}{5}),$
- $(\pm 2 \quad \pm 2 \quad 2 \quad \frac{4}{5}),$
- $(\pm 3 \quad \pm 3 \quad 2 \quad \frac{1}{2}),$
- $(\pm 3 \quad \pm 3 \quad 3 \quad \frac{1}{2}),$
- $(\pm 4 \quad \pm 4 \quad 0 \quad 0),$
- $(\pm 4 \quad \pm 4 \quad 5 \quad 0).$

All these points are vertices. Moreover, the last eight vertices span a facet $F = \{x \in P : x_4 = 0\}$, which is a 3-cube. The graph of P is isomorphic to the graph of the 5-cube.

Projecting the polytope P onto the facet F from a point beyond F yields a polytopal complex, the *Schlegel diagram* of P with respect to F , which is essentially equivalent to the boundary complex of P [12, Section 5.3], [18, Section 5.2]. For our example, the whole Schlegel diagram has the same symmetry group of order 16 as a prism over a square. Thus it is sufficient to consider a diagonal section as indicated in Fig. 2.

Now remove the 2-face spanned by the four vertices $(\pm 1, \pm 1, 4, 1)$ and merge the two facets containing it. This yields a regular cell complex C whose 1-skeleton is the same as before, that is, it is isomorphic to the graph of the 5-cube. Clearly, $\|C\|$ is homeomorphic to the 3-sphere. However, as realized, C is not polytopal because the angle α in Fig. 2 exceeds π .

It is fairly obvious that the modified cell complex C can be realized as a polytopal complex by an appropriate change of the coordinates, such that finally α becomes less than π . It may be less obvious that—for a special choice of coordinates—the transformed




Fig. 2. Schlegel diagram and a section.

diagram with $\alpha < \pi$ can be lifted to a 4-polytope P' . Here is a realization:

$$\begin{pmatrix} \pm 1 & \pm 1 & 1 & 0 \\ \pm 2 & \pm 2 & 4 & 0 \\ \pm 3 & \pm 3 & 3 & 1 \\ \pm 3 & \pm 3 & 2 & \frac{5}{4} \\ \pm 4 & \pm 4 & 2 & \frac{21}{20} \\ \pm 4 & \pm 4 & 3 & \frac{9}{5} \\ \pm \frac{56}{13} & \pm \frac{56}{13} & 0 & \frac{779}{260} \\ \pm 5 & \pm 5 & 16 & 0 \end{pmatrix}.$$

The polytope P' has one facet that has 12 vertices, namely,

$$(\pm 1, \pm 1, 1, 0), \quad (\pm 2, \pm 2, 4, 0), \quad (\pm 5, \pm 5, 16, 0).$$

Therefore P' is a 4-polytope with the graph of the 5-cube which is not cubical.

5. Alternating Oriented Matroids and Cyclic Polytopes

In Section 6 we construct a class of polytopes which are ‘‘cubical relatives’’ of the cyclic polytopes. The combinatorial structure of the cyclic polytopes is well known [12, Section 4.6], [18, Example 0.6]. We give a brief account in the framework of oriented matroids, which also captures the ‘‘interior combinatorial structure’’ of the cyclic polytopes.

Let $C_d(n)$ be the cyclic d -polytope on n vertices. It can be realized as the convex hull of n points on any curve of order d , such as the moment curve $t \mapsto (t, t^2, t^3, \dots, t^d)$ in \mathbb{R}^d .

Any point configuration in \mathbb{R}^d , and thus any polytope via its vertices, gives rise to an oriented matroid [18, Section 6.4], [4]. The positive cocircuits of the oriented matroid bijectively correspond to the facets of the polytope.

For a cyclic polytope $C_d(n)$ the associated oriented matroid is known as the *alternating oriented matroid* $\mathcal{C}(n, d + 1)$ of rank $d + 1$ on n points [4, Sections 3.4 and 9.4]. Homogenizing the vertices of $C_d(n)$ it can be represented by the (rows of the) matrix

$$\begin{pmatrix} 1 & t_1 & t_1^2 & \cdots & t_1^d \\ 1 & t_2 & t_2^2 & \cdots & t_2^d \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & t_n & t_n^2 & \cdots & t_n^d \end{pmatrix}$$

for $t_1 < t_2 < \dots < t_n$.

There is a notion of duality for oriented matroids which generalizes duality of projective spaces. In the case of the alternating oriented matroids, the dual of $\mathcal{C}(n, d + 1)$ is obtained from $\mathcal{C}(n, n - d - 1)$ by reorienting every other row in the representation above,

$$\mathcal{C}(n, d + 1)^* = \overline{\{2,4,6,\dots\}}\mathcal{C}(n, n - d + 1),$$

(see pp. 108–109 of [5] and Section 2 of [17]), that is, by the rows of the $n \times (n - d + 1)$ -matrix

$$\begin{pmatrix} 1 & t_1 & t_1^2 & \cdots & t_1^{n-d-1} \\ -1 & -t_2 & -t_2^2 & \cdots & -t_2^{n-d-1} \\ 1 & t_3 & t_3^2 & \cdots & t_3^{n-d-1} \\ -1 & -t_4 & -t_4^2 & \cdots & -t_4^{n-d-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ (-1)^{n+1} & (-1)^{n+1}t_n & (-1)^{n+1}t_n^2 & \cdots & (-1)^{n+1}t_n^{n-d-1} \end{pmatrix}.$$

From the representation of the alternating matroid and its dual given above it is obvious that that a deletion (omitting a row in the primal oriented matroid) or a contraction of the first element (omitting the first row and the first column in the primal oriented matroid, which amounts to omitting the first row in the dual) again gives an alternating matroid (on fewer points and, in the second case, of smaller rank).

The number of facets of $C_d(n)$, that is, the number of positive cocircuits of the alternating matroid $\mathcal{C}(n, d + 1)$, is known to be

$$\binom{n - \lceil d/2 \rceil}{\lfloor d/2 \rfloor} + \binom{n - 1 - \lceil (d - 1)/2 \rceil}{\lfloor (d - 1)/2 \rfloor},$$

by Gale’s evenness criterion [12, p. 63], [18, Theorem 0.7].

6. The General Case

The following is our main theorem: “Neighborly cubical polytopes exist!”—they can be obtained as projections of deformed cubes. As mentioned in the Introduction, the case $d = 2$ and the case $n = d + 1$ were known previously.

Theorem 16. *For any $n \geq d \geq 2r + 2$, there exists a combinatorial n -cube $C^n \subseteq \mathbb{R}^n$ and a linear projection map $\pi: \mathbb{R}^n \rightarrow \mathbb{R}^d$ such that $C_d^n := \pi(C^n)$ is a cubical d -polytope whose r -skeleton is isomorphic to that of C^n (via π).*

Proof. We first construct a combinatorial n -cube $C^n(\varepsilon) \subseteq \mathbb{R}^n$ that depends on a parameter $\varepsilon > 0$; then we verify that for ε sufficiently small the projection $\pi: \mathbb{R}^n \rightarrow \mathbb{R}^d$ to the last d coordinates preserves the r -skeleton; and finally we argue that $\pi(C^n(\varepsilon)) = C_d^n$ is cubical.

(A) For $0 < \varepsilon \leq 1$ define $C^n(\varepsilon) \subseteq \mathbb{R}^n$ as the solution set of

$$\varepsilon |x_k| \leq \frac{2 \binom{k}{2}}{\varepsilon^{k-1}} - (-1)^k \sum_{j=1}^{k-1} \binom{k-2}{j-1} x_j \quad \text{for } 1 \leq k \leq n. \tag{1}$$

This set is a combinatorial n -cube. To see this, we verify by induction on k that all solutions to the first k conditions of (1) satisfy

$$|x_k| < \frac{2^{\binom{k}{2}+1}}{\varepsilon^k}. \tag{2}$$

In fact, the upper bound of (2) increases with k , and for $k = 1$ we have $\varepsilon|x_1| \leq 1$, so $|x_1| < 2/\varepsilon$ is surely satisfied. Thus we may use induction, and for $k \geq 2$ estimate

$$\begin{aligned} \varepsilon|x_k| &\leq \frac{2^{\binom{k}{2}}}{\varepsilon^{k-1}} - (-1)^k \sum_{j=1}^{k-1} \binom{k-2}{j-1} x_j \\ &< \frac{2^{\binom{k}{2}}}{\varepsilon^{k-1}} + \sum_{j=1}^{k-1} \binom{k-2}{j-1} \frac{2^{\binom{k-1}{2}+1}}{\varepsilon^{k-1}} \\ &\leq \frac{2^{\binom{k}{2}}}{\varepsilon^{k-1}} + 2^{k-2} \frac{2^{\binom{k-1}{2}+1}}{\varepsilon^{k-1}} = \frac{2^{\binom{k}{2}+1}}{\varepsilon^{k-1}}. \end{aligned}$$

In this computation the second term is always smaller in absolute value than the first, that is,

$$\frac{2^{\binom{k}{2}}}{\varepsilon^{k-1}} - \left| \sum_{j=1}^{k-1} \binom{k-2}{j-1} x_j \right| > 0 \quad \text{for } 1 \leq k \leq n,$$

and from this we see that $C^n(\varepsilon)$ is a combinatorial cube. (It is an iterated deformed product in the sense of [2].)

(B) Now let $\pi: \mathbb{R}^n \rightarrow \mathbb{R}^d$ be the projection to the last d coordinates, $(x_1, \dots, x_n) \mapsto (x_{n-d+1}, \dots, x_n)$.

Claim 1. For sufficiently small $\varepsilon > 0$ and $n \geq d \geq 2r + 2$, the orthogonal projection $\pi: C^n(\varepsilon) \rightarrow \pi(C^n(\varepsilon)) =: C_d^n(\varepsilon)$ preserves the r -skeleton. That is, π restricts to an isomorphism

$$\pi: C^n(\varepsilon)^{[r]} \rightarrow C_d^n(\varepsilon)^{[r]}$$

of polytopal complexes.

To verify this, we need to see that every r -face F of $C^n(\varepsilon)$ is mapped bijectively to an r -face $\pi(F)$ of $C^n(\varepsilon)$. By Corollary 9 this is equivalent to the condition that for every r -face F there is a normal vector which is orthogonal to the direction of projection and another one which is not.

In our specific situation, let F be an r -face of $C^n(\varepsilon)$, and let $\mathbf{v} \in C^n(\varepsilon)$ be a vertex of F . Then there are unique signs $\sigma_1, \dots, \sigma_n \in \{+1, -1\}$ such that \mathbf{v} is determined by

$$(-1)^k \sum_{j=1}^{k-1} \binom{k-2}{j-1} v_j + \sigma_k \varepsilon v_k = \frac{2^{\binom{k}{2}}}{\varepsilon^{k-1}} \quad \text{for } 1 \leq k \leq n,$$

Claim 3.

- (i) If $\varepsilon = 0$, then all maximal minors of $\overline{A}(\Sigma)$ have nonzero determinant.
- (ii) If $\varepsilon > 0$ is so small that all maximal minors of $\overline{A}(\Sigma)$ have the same sign as for $\varepsilon = 0$, then every alternating subset of $n - d + 1$ rows of $\overline{A}(\Sigma)$ positively spans \mathbb{R}^{n-d} .
- (iii) For $n \geq d \geq 2r + 1$, every subset of $n - 1 - r$ rows of $\overline{A}(\Sigma)$ contains an alternating subset of size $n - d + 1$.

To see (i), we note that $\{1, t, \binom{t}{2}, \dots, \binom{t}{n-d+1}\}$ and $\{1, t, t^2, \dots, t^{n-d+1}\}$ are two different bases for the vector space of rational polynomials of degree at most $n - d + 1$. Thus the matrix $\overline{A} = \overline{A}(\Sigma)$ arises by invertible row operations from the matrix

$$\overline{B} = ((-1)^k (k - 1)^{j-1})_{2 \leq k \leq n, 1 \leq j \leq n},$$

whose maximal minors are Vandermonde determinants.

For (ii), observe the following: It can be seen from the oriented matroid of the vector configuration whether there is a positive linear relation between a set of vectors, such as the rows of \overline{A} . The condition (in order to obtain that it positively spans) is that the configuration must be a positive circuit, or, equivalently, *totally cyclic* [4, Section 3.4], [18, Section 6.4]. For ε small enough (as indicated), we have the same oriented matroid as for $\varepsilon = 0$, and hence as for \overline{B} . The oriented matroid $\mathcal{M}(\overline{B})$ of rank $n - d$ determined by the rows of \overline{B} is the dual of the alternating oriented matroid, which differs from the alternating oriented matroid by a reorientation of $\{3, 5, 7, \dots\}$, see Section 5. In $\mathcal{M}(\overline{B})$, the positive circuits are all subsets of size $n - d + 1$. In particular, the ground set is a positive vector, that is, there is a positive linear relation among all the rows of \overline{B} .

For part (iii), start with the alternating index set $\{2, 3, \dots, n\}$ for the rows of \overline{A} . Now successively delete any r rows from \overline{A} , but whenever a row is deleted, we remove also the next row above or below that has not yet been deleted. Thus in each of the (at most) r deletion steps, we remove two adjacent rows of \overline{A} , and hence the index set is kept to be alternating. After all this, we are left with a submatrix $\overline{\overline{A}}$ of \overline{A} that has at least $n - 1 - 2r$ rows and whose index set is alternating. Since $n - d + 1 \geq n - 1 - 2r$, we may take the first $n - d + 1$ rows of $\overline{\overline{A}}$.

(C) *The polytopes $C_d^n(\varepsilon)$ are indeed cubical.*

Let F be a facet of $C_d^n(\varepsilon)$. Its preimage $F' = \pi^{-1}(F)$ is a face of $C^n(\varepsilon)$ of dimension at least $d - 1$. Suppose $\dim F' \geq d$. From Lemma 7 we infer that some $n - d$ rows of the $n \times (n - d)$ -matrix A are (positively) linearly dependent. This is a contradiction to (i) of Claim 3. We conclude that $\dim F' = d - 1 = \dim F$. The claim now follows from Lemma 10. □

It follows from the discussion in the last paragraph that each facet of $C_d^n(\varepsilon)$ is an affinely isomorphic image of some $(d - 1)$ -face of $C_d^n = C_d^n(\varepsilon)$.

combination with positive coefficients for which the last $n - d - (p - 1)$ components vanish. We get the answer by comparison with $\{\pm(p + 1)\} \cup \alpha^{(p)}$:

- If the gap between $(p + 1)$ and $\alpha^{(p)}$ is *even*, then $\{\pm(p + 1)\} \cup \alpha^{(p)}$ is a positive circuit of its contraction by the Gale evenness criterion, for any small enough $\varepsilon > 0$, and hence also for $\varepsilon = 0$. Then the “ $(-1)^{p+1}$ ” component of $\pm(p + 1)$ can be replaced by the “ $(-1)^{p+1}\varepsilon$ ” component of $(-1)^{p+1} \cdot p$. Hence

*If the gap between p and $\alpha^{(p)}$ is **odd**, then we need $\sigma_p = (-1)^{p+1}$.*

- If the gap between $(p + 1)$ and $\alpha^{(p)}$ is *odd*, then $\{\pm(p + 1)\} \cup \alpha^{(p)}$ is a circuit with negative element “ $\pm(p + 1)$ ” and all other elements positive, since it *violates* Gale’s evenness criterion. However, then if we replace the “ $(-1)^{p+1}$ ” component of $\pm(p + 1)$ by a “ $(-1)^p\varepsilon$ ” component of $(-1)^p\varepsilon$, we get a positive circuit supported on $\{(-1)^p p\} \cup \alpha^{(p)}$. Hence

*If the gap between p and $\alpha^{(p)}$ is **even**, then we need $\sigma_p = (-1)^p$.*

The rest is easy: looking at the circuit on

$$\{\sigma_1 \cdot 1, \sigma_2 \cdot 2, \dots, \sigma_p p\} \cup \alpha^{(p)}$$

that is obtained by extending the linear combination of $\alpha^{(p)}$, we see that the coefficient of the row $\sigma_p \cdot p$ is very large (order of $O(1/\varepsilon)$) compared with the coefficients on $\alpha^{(p)}$. To compensate this, we need

$$\begin{aligned} \sigma_{p-1} &= -(-1)^p \text{ with coefficient of order } O(1/\varepsilon^2), \\ \sigma_{p-2} &= -(-1)^{p-1} \text{ with coefficient of order } O(1/\varepsilon^3), \end{aligned}$$

and so on, until

$$\sigma_1 = -(-1)^2 \text{ with coefficient of order } O(1/\varepsilon^p).$$

The case $p = 0$ is included in this. The case $p = n - d + 1$ ($\alpha^{(p)} = \emptyset$) is easy. □

The “cubical Gale evenness criterion” also allows us to count the facets of C_d^n .

Corollary 19. *For $n \geq d \geq 2$ and for ε sufficiently small, then the number $f(n, d) := f_{d-1}(C_d^n)$ of facets of C_d^n is given by*

$$\begin{aligned} f(n, d) &= 2d + 4 \sum_{p=0}^{n-d-1} \left(\binom{\lfloor d/2 \rfloor + p + 1}{p + 2} + \binom{\lfloor (d+1)/2 \rfloor + p}{p + 2} \right) 2^p \\ &= 2d + \sum_{p=2}^{n-d+1} \left(\binom{\lfloor d/2 \rfloor + p - 1}{p} + \binom{\lfloor (d-1)/2 \rfloor + p - 1}{p} \right) 2^p \\ &= \begin{cases} 2d + \sum_{p=2}^{n-d+1} \binom{k + p - 1}{p} 2^{p+1} & \text{for } d = 2k, \\ 2d + \sum_{p=2}^{n-d+1} \frac{p + 2k - 2}{p + k - 1} \binom{k + p - 1}{p} 2^p & \text{for } d = 2k + 1. \end{cases} \end{aligned}$$

We evaluate the function $f(n, d)$ for some particularly interesting choices of (n, d) :

$$\begin{aligned}
 d = 2: & \quad f(n, 2) = 2^n, \\
 d = 3: & \quad f(n, 3) = 2^n - 2, \\
 d = 4: & \quad f(n, 4) = (n - 2)2^{n-2}, \\
 d = 5: & \quad f(n, 5) = (n - 4)2^{n-2} + 2, \\
 n = d: & \quad f(d, d) = 2d, \\
 n = d + 1: & \quad f(d + 1, d) = d^2 + d + 2\lfloor d/2 \rfloor.
 \end{aligned}$$

For d even, the number $f(n, d)$ is already determined by n and d , together with the fact that we have a cubical d -polytope with the r -skeleton of the n -cube—using the cubical Dehn–Sommerville equations, see Proposition 3.

7. A Counterexample to the Cubical Upper Bound Conjecture

The following construction of a cubical 3-sphere starts with the cubical 4-polytope C_4^6 , whose f -vector is $(64, 192, 192, 64)$. By local “surgery” (the cubical equivalent of a bistellar flip) we will obtain a cubical 3-sphere with f -vector $(64, 196, 198, 66)$: thus we have a counterexample to Kalai’s cubical upper bound conjecture, verifying the second half of Theorem 15. The verification that the cubical flip can indeed be performed relies heavily on the description of C_4^6 given by the “cubical Gale evenness criterion” (Theorem 18).

Adopting standard oriented matroid notation we denote a k -face of the 6-cube by a sign vector in $\{+, -, 0\}^6$ with k zeros. Each k -face of C_4^6 corresponds to a k -face of the 6-cube. Thus the nonempty faces of C_4^6 also correspond to *certain* sign vectors in $\{+, -, 0\}^6$.

We show that C_4^6 does not maximize the f -vector among cubical 3-spheres with 64 vertices, as follows. Consider the following chain of three cubical facets $A = (-+00+0)$, $B = (-00++0)$, $C = (- -0+00)$. Consider the cubical 3-ball $\Phi = A \cup B \cup C$. Its boundary $\partial\Phi$ is a cubical 2-sphere.

Lemma 20. *Each facet of C_4^6 , except for $A, B,$ or $C,$ intersects $\partial\Phi$ in precisely one (possibly empty) face.*

Proof. Consider the following three pairs of 2-faces: $(A \setminus B, B \setminus A), (B \setminus C, C \setminus B),$ and $(A \setminus B, C \setminus B)$. We prove that for each such pair (X, Y) none of the vertices of X is on a common facet with any vertex of Y . The claim then follows. We proceed case by case.

Assume that F is a facet that contains any vertex from $A \setminus B = (-+0-+0)$ and any vertex from $B \setminus A = (- -0++0)$. Then $F = (-0u0+v)$, where $u, v \in \{+, -, 0\}$ are to be determined. Either $u = 0$ or $v = 0$. The “cubical Gale evenness criterion,” Theorem 18, case $p = 1$, implies that the initial minus sign must be followed by an even number of zeros, which is impossible.

In the second case $B \setminus C = (-+0++0)$ and $C \setminus B = (- -0+-0)$. Hence a presumptive facet would have coordinates $(-0w+0x)$ with either $w = 0$ or $x = 0$. By the “cubical Gale evenness criterion,” Theorem 18, case $p = 1$, the initial minus sign is

8. Comments

(1) A cubical d -polytope is k -stacked if it has a cubical subdivision without interior $(d - k - 1)$ -faces [3, Definition 5.3]. Thus every k -stacked cubical polytope is also $(k + 1)$ -stacked.

Proposition 21. *The neighborly cubical d -polytopes C_d^n are $\lfloor (d + 1)/2 \rfloor$ -stacked.*

Proof. The projection $\mathbb{R}^{d+1} \rightarrow \mathbb{R}^d$ that deletes the last coordinate maps C_{d+1}^n to C_d^n . The “upper faces” of the cubical polytope C_{d+1}^n (those with a normal vector whose last coordinate is positive) thus define a cubical subdivision B_d^n of C_d^n . Since, for $r = \lfloor d/2 \rfloor - 1$, all r -faces of C_{d+1}^n get mapped to the boundary of C_d^n , we conclude that C_d^n has no interior r -faces. Thus C_d^n is k -stacked, for $k = d - 1 - r = d - \lfloor d/2 \rfloor = \lfloor (d + 1)/2 \rfloor$. \square

In the setting of Babson et al. [3, Section 5] this yields a new extreme ray for the cubical g -cone for even d .

Corollary 22. *For even d and $k := d/2$, the neighborly cubical d -polytopes C_d^n form a sequence of polytopes for which the cubical g -vector is dominated by its k th component, that is,*

$$\lim_{n \rightarrow \infty} \frac{g_i^c(C_d^n)}{g_k^c(C_d^n)} = 0 \quad \text{for all } i \neq k.$$

Thus “the ray $\mathbb{R}^+ e_k$ lies in the closure of the Adin g -cone,” for even d and $k = d/2$, in the terminology of Section 5 of [3]. In particular, for $d = 4$ this implies that the closure of the Adin g -cone, i.e., of

$$\text{cone} \{(g_1^c, g_2^c) = (f_0 - 16, 4f_3 - 3f_0 + 16): \text{cubical 4-polytopes}\}$$

is the complete positive orthant in \mathbb{R}^2 .

(2) One is tempted to ask whether for any polytope P of dimension $n \geq d$ there is a d -polytope Q that has isomorphic r -skeleton, for $r = \lfloor d/2 \rfloor - 1$. In general, this is known to be false, due to a construction by Klee and Grünbaum, see Grünbaum [12, Theorem 12.2.2]. However, the case of the cross polytopes, $P = C_n^\Delta$, seems to be open and particularly interesting.

(3) Is there a construction of (even-dimensional) neighborly cubical polytopes that have all vertices on a sphere? Note that the trigonometric moment curve [12, p. 67], [18, p. 75] yields this in the simplicial case. By the explicit construction of a Schlegel diagram as the Delaunay subdivisions of a finite point set, Seidel [16] obtained such a construction for $d = 4$ and $n \leq 7$.

(4) The polytope $C_4^5 \cong P$ and the noncubical polytope P' of Section 4.4, as well as the polytope C_4^6 and the counterexample of Section 7, were first constructed using the

POLYMAKE system [10]. Corresponding POLYMAKE input files are available

<http://www.math.tu-berlin.de/diskregeom/polymake/examples/NCP/>.

The sequence of facets A , B , C used in Section 7 was found in an exhaustive computer search. The same test revealed that a similar construction does not work with the smaller C_4^5 polytope.

Acknowledgments

Thanks to Raimund Seidel for inspiring e-discussions. We are indebted to Roswitha Blind for a correction that led to a substantial modification in the construction of the counterexample of Section 7. Thanks to Gil Kalai and the anonymous referee for sharing with us their observation which led to Proposition 14.

References

1. R. M. Adin: A new cubical h -vector, *Discrete Math.* **157** (1996), 3–14.
2. N. Amenta and G. M. Ziegler: Deformed products and maximal shadows, in: *Advances in Discrete and Computational Geometry* (B. Chazelle, J. E. Goodman, and R. Pollack, eds.), Contemporary Mathematics, Vol. 223, American Mathematical Society, Providence, RI, 1998, pp. 57–90.
3. E. K. Babson, L. J. Billera, and C. S. Chan: Neighborly cubical spheres and a cubical lower bound conjecture, *Israel J. Math.* **102** (1997), 297–315.
4. A. Björner, M. Las Vergnas, B. Sturmfels, N. White, and G. M. Ziegler: *Oriented Matroids*, Encyclopedia of Mathematics, Vol. 46, Cambridge University Press, Cambridge, 1993; second edn., 1999.
5. R. Bland and M. Las Vergnas: Orientability of matroids, *J. Combin. Theory Ser. B* **24** (1978), 94–123.
6. G. Blind and R. Blind: Convex polytopes without triangular faces, *Israel J. Math.* **71** (1990), 129–134.
7. G. Blind and R. Blind: The almost simple cubical polytopes, *Discrete Math.* **184** (1998), 25–48.
8. H. Edelsbrunner: Size of hexahedral meshes, Problem for a course on “Geometry and Topology of Grid Generation,” University of Illinois at Urbana Champaign, April 2, 1999.
9. J. Erickson: Problem, Open Problem Session (E. Welzl, ed.) at “Discrete and Computational Geometry: Ten Years Later”, Mt. Holyoke College, July 15, 1996.
10. E. Gawrilow and M. Joswig: POLYMAKE—a framework for analyzing convex polytopes, TU Berlin 1997–1999: <http://www.math.tu-berlin.de/diskregeom/polymake/>
11. D. Goldfarb: On the complexity of the simplex algorithm, in: *Advances in Optimization and Numerical Analysis*, Proc. 6th Workshop on Optimization and Numerical Analysis, Oaxaca, Mexico, January 1992, Kluwer, Dordrecht, 1994, pp. 25–38.
12. B. Grünbaum: *Convex Polytopes*, Interscience, London, 1967.
13. W. Jockusch: The lower and upper bound problems for cubical polytopes, *Discrete Comput. Geom.* **9** (1993), 159–163.
14. G. Kalai: Email communication, March 1999.
15. K. G. Murty: Computational complexity of parametric linear programming, *Math. Programming* **19** (1980), 213–219.
16. R. Seidel: Email communication, July 1998.
17. B. Sturmfels: Neighborly polytopes and oriented matroids, *European J. Combin.* **9** (1988), 537–546.
18. G. M. Ziegler: *Lectures on Polytopes*, Graduate Texts in Mathematics, Vol. 152, Springer-Verlag, New York, 1995.

Received December 30, 1998. Online publication May 8, 2000.