Colorectal resection, both open and laparoscopic-assisted, in patients with benign indications is associated with proangiogenic changes in plasma angiopoietin 1 and 2 levels

**H. M. C. Shantha Kumara · A. Hoffman ·
I. Y. Kim · D. Feingold · N. Dujovny · M. Kalady ·
M. Luchtefeld · R. L. Whelan**

Published online: 27 November 2008
© Springer Science+Business Media, LLC 2008

Erratum to: Surg Endosc
DOI 10.1007/s00464-008-0132-5

The correct spelling of the sixth author is M. Kalady.

The online version of the original article can be found under
doi:[10.1007/s00464-008-0132-5](https://doi.org/10.1007/s00464-008-0132-5).

H. M. C. Shantha Kumara · A. Hoffman · D. Feingold ·
R. L. Whelan (✉)
Section of Colon & Rectal Surgery, Columbia University
College of Physicians & Surgeons, New York, USA
e-mail: rlw3@columbia.edu

H. M. C. Shantha Kumara
e-mail: sk2675@columbia.edu

I. Y. Kim
Department of Surgery, The Wonju College of Medicine,
Yonsei University, Wonju, South Korea

N. Dujovny · M. Luchtefeld
Division of Colon and Rectal Surgery,
Ferguson Clinic, Grand Rapids, MI, USA

M. Kalady
Department of Colon & Rectal Surgery,
Cleveland Clinic, Cleveland, OH, USA